

Bangladesh

Health Facility Survey 2022



Final Report











Bangladesh Health Facility Survey 2022

Final Report

National Institute of Population Research and Training (NIPORT) Medical Education and Family Welfare Division Ministry of Health and Family Welfare Dhaka, Bangladesh

> icddr,b Dhaka, Bangladesh

> > Funded by

Government of the People's Republic of Bangladesh U.S. Agency for International Development (USAID), Bangladesh

July 2024







This report presents final findings of the 2022 Bangladesh Health Facility Survey (2022 BHFS), which was implemented by the National Institute of Population Research and Training (NIPORT). icddr,b (International Centre for Diarrhoeal Disease Research, Bangladesh) provided technical assistance at every stages of the survey including monitoring the field work and data collection. Associates for Community and Population Research (ACPR), a private research agency, collected the field survey data. As a part of periodic assessment of health systems and services provided by various health facilities, the 2022 BHFS collected data to monitor and evaluate population, health, and nutrition programs under the Training, Research and Development (TRD) operational plan of the 4th Health, Population, and Nutrition Sector Program (HPNSP). The survey was funded by the government of Bangladesh and the U.S. Agency for International Development (USAID) provided financial support for technical assistance.

Additional information about the 2022 BHFS may be obtained from the NIPORT, 13/1 Sheikh Shaheb Bazar Road, Azimpur, Dhaka -1205 Bangladesh. Telephone: 88-02-5861-1206; Internet: www.niport.gov.bd or from icddr,b, 68 Shaheed Tajuddin Ahmed Sharani Mohakhali, Dhaka 1212 (GPO Box 128, Dhaka 1000) Bangladesh. Telephone: +(880-2) 8860523-32; Email: info@icddrb.org; Internet: http://www.icddrb.org

Cover inspiration: The Community Clinic represents the gateway towards formal healthcare services in Bangladesh which symbolizes promoting healthcare pathways as well as promoting human-centered care and experience regarding selective key health services, including Family Planning Services, Antenatal Care, Delivery and Newborn Care, and Child Health Services. This report systematically outlines these four key health services.

Cover design: Nuruzzaman Lucky, Senior Graphic Designer, MCHD, icddr,b Cover illustration: Sanzida, Freelance Illustrator and Designer

Recommended citation:

National Institute of Population Research and Training (NIPORT), and icddr,b (International Centre for Diarrhoeal Disease Research, Bangladesh). 2024. *Bangladesh Health Facility Survey 2022*. Dhaka, Bangladesh: NIPORT and icddr,b.

This publication was produced with the support of the United States Agency for International Development (USAID) under the terms of the Data for Impact (D4I) associate award 7200AA18LA00008, which is implemented by the Carolina Population Center at the Universi ty of North Carolina at Chapel Hill, in partnership with Palladium International, LLC; ICF Macro, Inc.; John Snow, Inc.; and Tulane University. The views expressed in this publication do not necessarily reflect the views of USAID or the United States government. TR-23-549 D4I

CONTENTS

| TABLE | ES AND | FIGUR | ES | vii |
|-------|------------|-----------|---|--------|
| FORE | WORD | ••••• | | xiii |
| PREFA | ACE | ••••• | | XV |
| ACKN | OWLED | OGMEN | Т | xvii |
| STAKE | EHOLDI | ER ADV | /ISORY COMMITTEE (SAC) FOR 2022 BHFS | xix |
| TECH | NICAL V | WORKI | ING COMMITTEE AND SAMPLING COMMITTEE FOR 2022 BHFS | xxi |
| CONT | RIBUTC | ORS TO | THE REPORT | xxiii |
| ACRO | NYMS A | AND AB | BREVIATIONS | XXV |
| READ | ING AN | D UNDI | ERSTANDING TABLES FROM THE 2022 BHFS | xxix |
| MAP C | OF BANC | GLADE | SH | xxxvi |
| 1 | OVER | VIEW C |)F THE HEALTH SYSTEM IN BANGLADESH | 1 |
| - | 1.1 | Health | Status in Bangladesh | |
| | 1.2 | Health. | Population, and Nutrition Sector Program | 1 |
| | 1.3 | Health | Service Delivery System of Bangladesh | 2 |
| | | 1.3.1 | Management Structure and Health Facilities under DGHS | 2 |
| | | 1.3.2 | Management Structure and Health Facilities under DGFP | 4 |
| | | 1.3.3 | Urban Primary Health Care Services Delivery Project (UPHCSDP) | 4 |
| | | 1.3.4 | NGO Health Programs | 5 |
| | | 1.3.5 | Private Health Sector | 5 |
| | 1.4 | Health | Financing | 5 |
| | 1.5 | Health | Stewardship | 6 |
| 2 | METH | | YCV | 7 |
| Z | | Oportic | /G1 | |
| | 2.1 | Instituti | ional Framework and Objectives of the 2022 BHFS | / Q |
| | 2.2 | 221 | Institutional Framework | 8 |
| | | 2.2.1 | Objectives of the 2022 BHFS | |
| | 23 | Data Co | objectives of the 2022 DTH 5 | 8 |
| | 2.5 | Survey | Implementation | 0 |
| | 2.1 | 2.4.1 | Ouestionnaire Adaptation | 9 |
| | | 2.4.2 | Pre-test | 9 |
| | | 2.4.3 | Main Training | 9 |
| | | 2.4.4 | Data Collection | |
| | | 2.4.5 | Data Management | |
| | | 2.4.6 | Data Analysis and Report Preparation | 11 |
| | 2.5 | Sampli | ng | 11 |
| | | 2.5.1 | Sampling Methodology | 11 |
| | | 2.5.2 | Sampling Frame | 12 |
| | | 2.5.3 | 2022 BHFS Sample Size and Sample Allocation | 12 |
| | | 2.5.4 | Sample of Health Facilities and Outcomes | 12 |
| | | 2.5.5 | Sample of Health Service Providers | 12 |

| FAC | FACILITY-LEVEL INFRASTRUCTURE, RESOURCES, MANAGEMENT, AND SUPPORT | | | | |
|---------------|--|-------------------|--|--|--|
| ······ 3 1 | Background | 19 20 | | | |
| 3.1 | Availability of Services | 20 | | | |
| 5.2 | 3.2.1 Overall Availability of Specific Client Services | 20 | | | |
| | 3.2.1 Availability of Basic Client Services | | | | |
| | 3.2.2 General Service Prenaredness | | | | |
| 33 | Basic Management Systems to Support and Maintain Quality Services and A | npropriate | | | |
| 5.5 | Client Utilization | 36 | | | |
| | 3.3.1 Supportive Management Practices for Providers | 36 | | | |
| | 3.3.2 Quality Assurance | | | | |
| 31 | Availability of Human Resources for Health | | | | |
| 3.5 | Management of Medical Waste in Health Care Facilities | | | | |
| CHI | LD HEALTH SERVICES | 64 | | | |
| 4.1 | Background | 65 | | | |
| 4.2 | Availability of Child Health Services | 65 | | | |
| | 4.2.1 Provision of Outpatient Curative Care, Child Growth Monitoring, C | hild Vaccination, | | | |
| | and Nutrition Services | 65 | | | |
| | 4.2.2 Availability of Guidelines, Trained Staff, and Basic Equipment for | Child Curative | | | |
| | Care | 66 | | | |
| | 4.2.3 Availability of Medicines and Commodities for Child Health Care | 73 | | | |
| | 4.2.4 Infection Control in Sick Child Services | 77 | | | |
| | 4.2.5 Laboratory Diagnostic Capacity | | | | |
| 4.3 | Readiness of Health Facilities to Provide Child Curative Care | | | | |
| 4.4 | Basic Management and Administrative System | 80 | | | |
| | 4.4.1 Personal Supervision and Training | | | | |
| | 4.4.2 In-Service Training by Topic | | | | |
| FAM | AILY PLANNING SERVICES | 94 | | | |
| 5.1 | Background | 95 | | | |
| 5.2 | Family Planning Services | 95 | | | |
| | 5.2.1 Availability of Family Planning Services | 96 | | | |
| | 5.2.2 Facilities Providing Modern Family Planning Methods and Services | 97 | | | |
| | 5.2.3 Frequency of Availability of Family Planning Services | | | | |
| 5.3 | Availability of FP Commodities on the Day of Assessment | | | | |
| | 5.3.1 Availability of Oral Pills | | | | |
| | 5.3.2 Availability of Injectables | | | | |
| | 5.3.3 Availability of Condoms | | | | |
| | 5.3.4 Availability of Intrauterine Contraceptive Devices | | | | |
| | 5.3.5 Availability of Implants (One Rod or Two Rod) | | | | |
| 5.4 | Availability of Service Guidelines, Trained Staff, and Equipment | | | | |
| | 5.4.1 Availability of Guidelines for Family Planning Services | | | | |
| | 5.4.2 Availability of Trained Staff in Family Planning | | | | |
| | 5.4.3 Availability of Equipment for Family Planning Services | | | | |
| | 5.4.4 Infection Control During Provision of Family Planning | | | | |
| 5.5 | Readiness of Health Facilities to Provide Family Planning Services | | | | |
| 5.6 | Basic Management and Administrative Systems | | | | |
| | 5.6.1 Training and Supervision | | | | |

| | | 5.6.2 | In-Service Training for Family Planning Service Providers by Topic | 110 |
|---|------|---------|--|-----|
| 6 | ANTH | ENATAL | CARE SERVICES | 123 |
| | 6.1 | Backgr | ound | 124 |
| | 6.2 | Availat | bility of Antenal Care (ANC) Services | 124 |
| | | 6.2.1 | ANC Service Provision | 124 |
| | | 6.2.2 | Availability of ANC Guidelines, Trained Staff, and Equipment | 125 |
| | | 6.2.3 | Availability of Laboratory Diagnostic Tests | 129 |
| | | 6.2.4 | Availability of Medicines for Routine ANC | 132 |
| | | 6.2.5 | Availability of Infection Control Items | 133 |
| | 6.3 | Readin | ess of Health Facilities to Provide ANC Services | 134 |
| | 6.4 | Basic N | Management and Administrative Systems | 135 |
| | | 6.4.1 | Recent In-service Training and Supervision | 135 |
| | | 6.4.2 | In-service Training by Topic | 136 |
| 7 | DELI | VERY A | ND NEWBORN CARE | 147 |
| | 7.1 | Backgr | ound | 148 |
| | 7.2 | Availat | bility of Maternal Health Services | 149 |
| | | 7.2.1 | Service Provision | 149 |
| | | 7.2.2 | Trained Staff, Guidelines, and Equipment for Delivery Services | 151 |
| | | 7.2.3 | Medicines and Commodities for Delivery Care | 158 |
| | | 7.2.4 | Items for Infection Control during Provision of Delivery Care | 161 |
| | 7.3 | Signal | Functions for Emergency Obstetric Care | 162 |
| | 7.4 | Readin | ess of Health Facilities to Provide Normal Delivery Services | 164 |
| | 7.5 | Newbo | rn Care Practices | 165 |
| | | 7.5.1 | Routine Newborn Care | 165 |
| | | 7.5.2 | Essential Medicines for Newborn Care | 166 |
| | | 7.5.3 | Availability of Equipment for Newborn Care Services | 168 |
| | 7.6 | Newbo | rn Signal Functions | 171 |
| | 7.7 | Basic N | Management and Administrative Systems | 175 |
| | | 7.7.1 | Recent In-Service Training and Supervision | 175 |
| | | 7.7.2 | In-Service Training in Newborn and Delivery Care by Topic | 175 |
| 8 | NON- | COMM | JNICABLE DISEASES | 194 |
| | 8.1 | Backgr | ound | 196 |
| | 8.2 | Availat | bility of Services for Diabetes | 197 |
| | | 8.2.1 | Service Provision | 197 |
| | | 8.2.2 | Availability of Diagnostic Capacity and Medicines for Diabetes | 199 |
| | 8.3 | Availat | bility of Services for Cardiovascular Disease | 200 |
| | | 8.3.1 | Service Provision | 200 |
| | | 8.3.2 | Availability of Medicines and Commodities Related to Cardiovascular Diseas | 202 |
| | 8.4 | Availat | bility of Services for Chronic Respiratory Diseases | 203 |
| | | 8.4.1 | Service Provision | 203 |
| | | 8.4.2 | Availability of Medicines and Commodities for Chronic Respiratory Diseases | 205 |
| | 8.5 | Availal | bility of Services for Hypertension | 206 |
| | | 8.5.1 | Service Provision | 206 |
| | | 8.5.2 | Availability of Medicines and Commodities for Hypertension | 208 |
| | 8.6 | Availat | bility of Services for Cervical Cancer | 210 |
| | | 8.6.1 | Availability of Screening Services for Cervical Cancer | 210 |

| | | 8.6.2 Service Guidelines | 211 |
|-------|--------|--|-----|
| | | 8.6.3 Trained Staff | |
| | | 8.6.4 Equipment | 211 |
| 9 | TUBE | RCULOSIS | |
| | 9.1 | Background | |
| | 9.2 | Availability of TB Services | |
| | | 9.2.1 Service Provision | |
| | | 9.2.2 Availability of Diagnostic Capacity and Medicines for TB Treatment | |
| | 9.3 | Readiness of Health Facilities to Provide TB Services | 230 |
| REFER | RENCES | S | 237 |
| APPEN | DIX A | PERSONS INVOLVED IN THE SURVEY | 241 |
| APPEN | DIX B | SUMMARY INDICATORS | 245 |
| APPEN | DIX C | QUESTIONNAIRES | 249 |

TABLES AND FIGURES

| 1 | OVERVIEW OF THE HEALTH SYSTEM IN BANGLADESH | | | | | |
|---|---|--|----------|--|--|--|
| | Table 1.1 | Secondary and primary health care facilities in public sector | 3 | | | |
| 2 | METHODOL | METHODOLOGY | | | | |
| | Table 2.1 | Bangladesh health facility distribution by division and health facility type | 15 | | | |
| | Table 2.2 | Health facility sample allocation by division and health facility type | 15 | | | |
| | Table 2.3 | Results of facility contacts by background characteristics | 16 | | | |
| | Table 2.4 | Distribution of surveyed facilities by background characteristics | 17 | | | |
| | Table 2.5 | Distribution of interviewed providers | 18 | | | |
| 3 | FACILITY-L | EVEL INFRASTRUCTURE, RESOURCES, MANAGEMENT, AND SU | PPORT | | | |
| | Table 3.1 | Availability of specific services | 43 | | | |
| | Table 3.2 | Availability of basic client services | 44 | | | |
| | Table 3.3 | Availability of basic amenities for client services | 45 | | | |
| | Table 3.4 | Availability of basic equipment | 46 | | | |
| | Table 3.5 | Standard precautions for infection control | 47 | | | |
| | Table 3.6 | Capacity for processing equipment for reuse | 48 | | | |
| | Table 3.7.1 | Availability of laboratory diagnostic testing by facility type | 49 | | | |
| | Table 3.7.2 | Laboratory diagnostic capacity by facility type | 50 | | | |
| | Table 3.7.3 | Laboratory diagnostic capacity by location and division | 51 | | | |
| | Table 3.8.1 | Availability of essential medicines by facility type | 52 | | | |
| | Table 3.8.2 | Availability of essential medicines by location and division | 53 | | | |
| | Table 3.8.3 | Availability of medicines in community clinics | 54 | | | |
| | Table 3.8.4 | Availability of DDS kit medicines | 55 | | | |
| | Table 3.8.5 | Availability of medicines and equipment of delivery kit | 56 | | | |
| | Table 3.9 | Supportive management practices at the facility level | 57 | | | |
| | Table 3.10 | Quality assurance and client opinion | 58 | | | |
| | Table 3.11.1 | Staffing pattern in surveyed facilities | 59 | | | |
| | Table 3.11.2 | Staffing pattern in surveyed facilities (medical officer and nurse) | 60 | | | |
| | Table 3.12.1 | Sharps waste management | 61 | | | |
| | Table 3.12.2 | Non-sharps waste management | 62 | | | |
| | Table 3.13 | Key indicators for general service preparedness in MCWCs | 63 | | | |
| | Figure 3.1 | Availability of basic client services in public health facilities | 21 | | | |
| | Figure 3.2 | Availability of electricity from national grid/polli-biddut, by facility type | 23 | | | |
| | Figure 3.3 | Availability of regular electricity, by facility type | 24 | | | |
| | Figure 3.4 | Availability of client latrine, by facility type | 25 | | | |
| | Figure 3.5 | Availability of communication equipment (land-line/mobile phone), by faci | lity | | | |
| | Figure 3.6 | type Availability of a functioning computer with Internet, by facility type | 26 27 | | | |
| | Figure 3.7 | Availability of emergency transport, by facility type | 28 | | | |
| | | i i and i i i i i i i i i i i i i i i i i i i | | | | |

| | Figure 3.8 | Average score of basic amenities (out of 6 amenities), by facility type | 29 |
|---|-------------|--|-------------|
| | Figure 3.9 | Average score of availability of all infection control items (out of 16), by facility type | 30 |
| | Figure 3.10 | Average capacity to conduct basic diagnostic tests (out of five), by facility t | уре |
| | | | 32 |
| | Figure 3.11 | Functioning x-ray machine in health facilities, by facility type | 33 |
| | Figure 3.12 | Functioning ultrasound machines in health facilities, by facility type | 34 |
| | Figure 3.13 | Quality assurance, by facility type | 38 |
| | Figure 3.14 | Sharps waste management, by facility type | 40 |
| | Figure 3.15 | Non-sharps waste management, by facility type | 41 |
| 4 | CHILD HEA | LTH SERVICES | |
| | Table 4.1 | Availability of child health services | 84 |
| | Table 4.2 | Guidelines and equipment for outpatient child curative care services | 85 |
| | Table 4.2.1 | Equipment for inpatient child curative care services | 86 |
| | Table 4.3 | Trained staff for child curative care services | |
| | Table 4.4 | Availability of essential and priority medicines | 88 |
| | Table 4.5 | Infection control and laboratory diagnostic capacity | 00 89 |
| | Table 4.6 | Readiness of health facilities to provide child curative care services | رہ ۵0 |
| | Table 4.0 | Supportive management for providers of shild health services | 90 01 |
| | Table 4.7 | Supportive management for providers of child health services | 19 م |
| | Table 4.8 | I raining for child health service providers | 92 |
| | Table 4.9 | Key indicators for child curative care services in MCWCs | 93 |
| | Figure 4.1 | Availability of child health services in 2022, by facility type | 66 |
| | Figure 4.2 | Availability of IMCI guidelines in health facilities for outpatient child curati | ive |
| | | care, by facility type | 67 |
| | Figure 4.3 | Availability of IMCI trained staff (at any time) in health facilities for outpat | ient |
| | e | child curative care, by facility type | 68 |
| | Figure 4.4 | Availability of functioning child weighing scale for outpatient child curative | • |
| | 118010 | care hy facility type | 69 |
| | Figure 4.5 | Availability of functioning thermometer for outpatient child curative care. | 07 v |
| | Tigure 4.5 | facility type | י 70 |
| | Figure 16 | Availability of growth chart in health facilities for outpatient child curative | 70 |
| | Figure 4.0 | Availability of growth chart in health facilities for outpatient child curative | 71 |
| | E | Care, by facinity type | / 1 |
| | Figure 4.7 | Average score of equipment (child weigning scale, length or height board, | |
| | | thermometer, and stethoscope) for child curative care, by facility type | 72 |
| | Figure 4.8 | Availability of ORS in health facilities for outpatient child curative care, by | |
| | | facility type | 73 |
| | Figure 4.9 | Availability of zinc tablets or syrup in health facilities for outpatient child | |
| | | curative care, by facility type | 74 |
| | Figure 4.10 | Availability of amoxillin in health facilities for outpatient child curative care | e, |
| | | by facility type | 75 |
| | Figure 4.11 | Availability of paracetamol syrup or suspension in health facilities for | |
| | C C | outpatient child curative care, by facility type | 76 |
| | Figure 4.12 | Availability of mebendazole/albendazole in health facilities for outpatient cl | nild |
| | | curative care, by facility type | |
| | Figure 4 13 | Average score of infection control items (out of 6 items) for child curative c | are |
| | 115010 7.15 | hy facility type | יייי, דד |
| | | <i>by</i> memory type | / / |

| | Figure 4.14 | Average readiness score of health facilities (out of 10 items) to provide child | 70 |
|---|--------------------------|---|------------|
| | Figure 4 15 | Drovidar training on tonias related to shild health in the last 24 months. | 19 00 |
| | Figure 4.15 | Provider training on topics related to child health actual in the last 24 months | 80 |
| | Figure 4.16 | before the survey | 01 |
| | Eigung 4 17 | Denviden training on tenios related to shild health in multip health facilities | 81 |
| | Figure 4.17 | Provider training on topics related to child health in public health facilities | 02 |
| | | during the last 24 months | 82 |
| 5 | FAMILY PL | ANNING SERVICES | ••••• |
| | Table 5.1 | Availability of family planning services | . 112 |
| | Table 5.2 | Methods of family planning provided | . 113 |
| | Table 5.3 | Methods of family planning provided | . 114 |
| | Table 5.4 | Frequency of availability of family planning services | . 115 |
| | Table 5.5 | Availability of family planning commodities | . 116 |
| | Table 5.6 | Guidelines, trained staff, and basic equipment for family planning services | . 117 |
| | Table 5.7 | Items for infection control during provision of family planning | . 118 |
| | Table 5.8 | Readiness of health facilities to provide family planning services | . 119 |
| | Table 5.9 | Supportive management for providers of family planning services | . 120 |
| | Table 5.10 | Training for family planning service providers | . 121 |
| | Table 5.11 | Key indicators for family planning services in MCWCs | 122 |
| | Figure 5.2 Figure 5.3 | Facilities providing LARC/PMs, by facility type Facilities providing male or female sterilization services, by facility type | 98 99 |
| | Figure 5.4 | Availability of oral pills, by facility type | . 100 |
| | Figure 5.5 | Availability of injectables, by facility type | . 101 |
| | Figure 5.6 | Availability of condoms, by facility type | . 102 |
| | Figure 5.7 | Availability of IUCDs, by facility type | . 103 |
| | Figure 5.8 | Availability of implants in health facilities, by facility type | . 104 |
| | Figure 5.9 | Availability of FP guidelines in health facilities, by facility type | . 105 |
| | Figure 5.10 | Availability of staff trained in FP at any time in health facilities, by facility type | . 106 |
| | Figure 5.11 | Availability of functioning blood pressure apparatus in heath facilities, by facility type | . 107 |
| | Figure 5.12 | Average number of items (out of six) available for infection control in family planning service areas, by facility type | y . 108 |
| | Figure 5.13 | Average readiness score of health facilities (out of 8 items) to provide FP services, by facility type | . 109 |
| 6 | ANTENATA | L CARE SERVICES | 1 |
| | Table 6.1 | Availability of antenatal care services | . 138 |
| | Table 6.2 | Guidelines, trained staff, and basic equipment for antenatal care services | . 139 |
| | Table 6.3 | Diagnostic capacity | . 140 |
| | Table 6.4 | Availability of medicines for routine antenatal care | . 141 |
| | Table 6.5 | Items for infection control during provision of antenatal care | . 142 |
| | Table 6.6 | Readiness of health facilities to provide antenatal care services | . 143 |
| | Table 6.7 | Supportive management for providers of antenatal care services | . 144 |
| | Table 6.8 | Training for antenatal care service providers | . 145 |
| | Table 6.9 | Key indicators for antenatal care services in MCWCs | . 146 |

| Figure 6.1 | Availability of ANC services in health facilities, by facility type | 125 |
|--------------|--|-------|
| Figure 6.2 | Availability of ANC guidelines in health facilities, by facility type | 126 |
| Figure 6.3 | Availability of ever trained staff for ANC in health facilities, by facility type | 127 |
| Figure 6.4 | Availability of blood pressure (BP) apparatus to support ANC services | 128 |
| Figure 6.5 | Average score of equipments (out of 6 items) to provide ANC services, by | |
| | facility type | . 129 |
| Figure 6.6 | Availability of blood hemoglobin (Hb) test in health facilities to provide AN | С |
| | services, by facility type | 130 |
| Figure 6.7 | Availability of urine protein test in health facilities to provide ANC services, | |
| | by facility type | . 131 |
| Figure 6.8 | Average score of laboratory diagnostic tests (out of 4 tests) to provide ANC | |
| | services, by facility type | . 132 |
| Figure 6.9 | Availability of iron or folic acid in health facilities to provide ANC services. | . 133 |
| Figure 6.10 | Average score of infection control items (out of 6 items) for ANC services, b | y |
| | facility type | 134 |
| Figure 6.11 | Average readiness score of health facilities (out of 6 items) to provide ANC | |
| C | services, by facility type | 135 |
| | | |
| DELIVERY A | AND NEWBORN CARE | |
| Table 7.1 | Availability of maternal health services | . 179 |
| Table 7.2 | Guidelines, trained staff, and equipment for delivery services | . 180 |
| Table 7.3 | Availability of staff trained on normal delivery or newborn care | . 181 |
| Table 7.4 | Medicines and commodities for delivery | . 182 |
| Table 7.5 | Items for infection control during provision of delivery care | . 183 |
| Table 7.6 | Signal functions for emergency obstetric care | . 184 |
| Table 7.7 | Readiness of health facilities to provide normal delivery services | . 185 |
| Table 7.8 | Newborn care practices | . 186 |
| Table 7.9 | Essential medicines for newborn care | . 187 |
| Table 7.10 | Availability of equipment for newborn care services | . 188 |
| Table 7.10.1 | Newborn signal functions | . 189 |
| Table 7.11 | Supportive management for providers of delivery care | . 190 |
| Table 7.12 | Training for providers of normal delivery services: Immediate newborn care | . 191 |
| Table 7.13 | Training for providers of normal delivery services: Immediate newborn care | . 192 |
| Table 7.14 | Key indicators for delivery and newborn care services in MCWCs | . 193 |
| | | |
| Figure 7.1.1 | Availability of normal delivery services in health facilities, by facility type | . 150 |
| Figure 7.1.2 | Availability of cesarean delivery services in health facilities, by facility type | . 151 |
| Figure 7.2.1 | Availability of trained staff in delivery care (at any time) in health facilities, | by |
| | facility type | . 152 |
| Figure 7.2.2 | Availability of guidelines on maternal health in health facilities that offer | |
| | normal delivery services, by facility type | . 153 |
| Figure 7.2.3 | Availability of emergency transport in health facilities, by facility type | 154 |
| Figure 7.2.4 | Availability of delivery beds in health facilities, by facility type | . 154 |
| Figure 7.2.5 | Availability of functioning examination light in health facilities. by facility | |
| 6 | type | 155 |
| Figure 726 | Availability of delivery pack in health facilities by facility type | 156 |
| Figure 7 2 7 | Availability of partograph in health facilities by facility type | 156 |
| Figure 7.2.7 | Availability of gloves in health facilities by facility type | 157 |
| Figure 7.2.1 | Availability of avutacin in health facilities, by facility type | 150 |
| rigule 7.3.1 | Availability of oxytochi in health facilities, by facility type | 139 |

| Figure 7.3.2 | Availability of injectable antibiotics in health facilities, by facility type 159 |
|-----------------|--|
| Figure 7.3.3 | Availability of magnesium sulphate in health facilities, by facility type 159 |
| Figure 7.3.4 | Availability of skin disinfectant in health facilities, by facility type |
| Figure 7.3.5 | Availability of intravenous fluids with infusion set in health facilities, by |
| 8 | facility type |
| Figure 7.3.6 | Average items available (out of six) for infection control in delivery service |
| 118010 / 1010 | area 162 |
| Figure 7.4 | Trends in REmOC signal functions by facility type 163 |
| Figure 7.5 | Trends in CEmOC signal functions, by facility type |
| Figure 7.5 | Average readiness seere of health facilities (out of 12 items) to provide normal |
| Figure 7.0 | delivery services by facility type |
| E | denivery services, by facility type |
| Figure $7.7.1$ | Availability of amoxicillin in health facilities, by facility type |
| Figure 7.7.2 | Availability of gentamicin in health facilities, by facility type |
| Figure 7.7.3 | Availability of 7.1% chlorhexidine in health facilities, by facility type |
| Figure 7.8.1 | Availability of functioning infant weighing scale for newborn care services, by |
| | facility type168 |
| Figure 7.8.2 | Availability of functioning fetal stethoscope for newborn care services, by |
| | facility type |
| Figure 7.8.3 | Availability of functioning newborn bag and mask for newborn care services, |
| | by facility type170 |
| Figure 7.8.4 | Availability of functioning suction apparatus with catheter in health facilities, |
| | by facility type 170 |
| Figure 7.8.5 | Availability of functioning suction bulb or penguin sucker for newborn care |
| | services, by facility type |
| Figure 7.9.1 | Primary newborn signal functions, by facility type, 2022 BHFS 172 |
| Figure 7.9.2 | Basic newborn signal functions, by facility type, 2022 BHFS |
| Figure 7.9.3 | Comprehensive newborn signal functions, by facility type, 2022 BHFS 173 |
| Figure 7.9.4 | Advanced newborn signal functions, by facility type, 2022 BHFS |
| Figure 7.10.1 | Training on emergency triage assessment for newborn care services, by facility |
| U | type |
| Figure 7.10.2 | In-service training on different services* for newborn care, by facility type 177 |
| 118010 /11012 | |
| NON-COMM | UNICABLE DISEASES |
| Table 8.1 | Availability of diabetes services and guidelines, trained staff, and equipment 214 |
| Table 8.2 | Diagnostic capacity and essential medicines for diabetes |
| Table 8.3 | Availability of cardiovascular disease services and guidelines, trained staff, and |
| | equipment |
| Table 8.4 | Availability of essential medicines and commodities for |
| | cardiovascular diseases 217 |
| Table 8 5 | Availability of chronic respiratory disease services and guidelines trained |
| 1 4010 0.5 | staff and equipment 218 |
| Table 96 | Availability of assortial medicines and commodities for abronia |
| 1 able 8.0 | Availability of essential medicines and commodities for chrome |
| T 11 07 | |
| adle 8./ | Availability of hypertension services and guidelines, trained starr, and |
| T 11 0.0 | equipment |
| Table 8.8 | Availability of essential medicines and commodities for hypertension |
| Table 8.9 | Availability of cervical cancer diagnosis/screening services and guidelines, |
| | trained staff, and equipment |

| Figure 8.1 | Availability of diabetes services in health facilities, by facility type | 198 |
|-------------|---|------|
| Figure 8.2 | Trained staff, guidelines, and basic equipment to provide diabetes services | in |
| | public facilities | 199 |
| Figure 8.3 | Diagnostic capacity and medicines to provide diabetes services in public | |
| | facilities | 200 |
| Figure 8.4 | Availability of cardiovascular disease services in health facilities, by facility | у |
| - | type | 201 |
| Figure 8.5 | Availability of trained staff, guidelines, and basic equipment to provide | |
| - | cardiovascular disease services in public facilities | 202 |
| Figure 8.6 | Availability of commodities and medicines to provide cardiovascular diseas | se |
| U | services in public facilities | 203 |
| Figure 8.7 | Availability of chronic respiratory disease services in health facilities, by | |
| C | facility type | 204 |
| Figure 8.8 | Availability of trained staff, guidelines, and basic equipment to provide chr | onic |
| C | respiratory disease-related services among public facilities | 205 |
| Figure 8.9 | Availability of commodities and medicines to provide chronic respiratory | |
| C | disease services among public facilities | 206 |
| Figure 8.10 | Availability of hypertension services in health facilities, by facility type | 207 |
| Figure 8.11 | Availability of trained staff, guidelines, and basic equipment to provide | |
| C | hypertension services among public facilities | 208 |
| Figure 8.12 | Availability of medicines to provide services for hypertension among public | с |
| - | facilities | 209 |
| Figure 8.13 | Availability of cervical cancer screening/diagnosis services in health faciliti | ies, |
| - | by facility type | 211 |
| Figure 8.14 | Availability of trained staff, guidelines, and basic equipment to | |
| C | diagnose/screen cervical cancer in public facilities | 212 |
| | | |
| TUBERCUL | OSIS | |
| Table 9.1 | Availability of tuberculosis services, guidelines, and trained staff for | |
| | tuberculosis services | 233 |
| Table 9.2 | Diagnostic capacity and availability of medicines for tuberculosis treatment | 234 |
| Table 9.3 | Readiness of health facilities to provide TB services | 235 |
| | | |
| Figure 9.1 | Availability of TB services (TB diagnostic, treatment and/or treatment follo | W- |
| | up services), by facility type | 226 |
| Figure 9.2 | Availability of any guideline for TB services, by facility type | 227 |
| Figure 9.3 | Availability of staff trained at any time for TB services, by facility type | 228 |
| Figure 9.4 | Availability of TB microscopy in health facilities, by facility type | 229 |
| Figure 9.5 | Availability of first-line TB medications in health facilities, by facility type | 230 |
| Figure 9.6 | Average readiness score of health facilities for providing TB services | 231 |

FOREWORD





Md. Azizur Rahman Secretary Medical Education and Family Welfare Division Ministry of Health and Family Welfare Government of the People's Republic of Bangladesh

This is my pleasure to congratulate the National Institute of Population Research and Training (NIPORT) on publishing the fifth Bangladesh Health Facility Survey (2022 BHFS) with financial support from the Government of Bangladesh and U.S. Agency for International Development (USAID). International Centre for Diarrheal Disease Research, Bangladesh (icddr,b) provided overall technical assistance. In addition, Data for Impact (D4I) provided support for finalizing the final report.

The information was gathered on the availability of basic and essential health care services and the readiness of health facilities to provide quality services in child health, maternal and newborn care, family planning, tuberculosis, and non-communicable diseases. Additionally, the survey assessed the functionality of delivery and newborn care services. The survey provides important information for policymakers and program personnel in addressing the monitoring and evaluation needs of the Health, Population and Nutrition Sector Programs (HPNSP) of the Ministry of Health Family Welfare (MOHFW).

The findings of the survey revealed challenges and weakness in facility preparedness that is impeding service delivery. We hope that policymakers and program managers will focus on the major gaps identified in the 2022 BHFS that require prompt action for improving the quality of care. Moreover, there is a dire need for developing a concrete and evidence-based plan, as well as regular surveillance of the service delivery.

We urge all stakeholders to play active roles in implementing national priorities to achieve goal of HPNSP and Sustainable Development Goals (SDGs). I hope that academicians, researchers, and program personnel will conduct further comprehensive analyses of 2022 BHFS data to provide additional in-depth knowledge for the future direction and effective implementation of the sector programs.

I express my sincere gratitude to all those who contributed to the successful completion of the survey. I wish to formally acknowledge the invaluable contributions of the representatives of Stakeholder Advisory Committee (SAC); Technical Working Committee (TWC) members; officials of Directorate General of Health Services (DGHS); Directorate General of Family Planning (DGFP); icddr,b; D4I; ACPR; the field staffs: data processing team and specially survey respondents.

Finally, I would like to thank NIPORT for their leadership in conducting the survey. I appreciate icddr,b, D4I, and ACPR for their collaborative effort. I deeply acknowledge the invaluable financial support provided by the Government of Bangladesh and USAID, which played a crucial role in ensuring the ultimate success of this important national survey.

(Md. Azizur Rahman)

PREFACE





Dr. Ashrafi Ahmad Director General National Institute of Population Research and Training (NIPORT) Medical Education and Family Welfare Division Ministry of Health and Family Welfare

The 2022 Bangladesh Health Facility Survey (BHFS) is the fifth round of nationally representative health facility survey conducted in Bangladesh. The series started in 2009 and was followed by surveys in 2011, 2014, and 2017. The 2022 BHFS was implemented by the National Institute of Population Research and Training (NIPORT). International Centre for Diarrheal Disease Research, Bangladesh (icddr,b) provided overall technical assistance for the survey. Additionally, Data for Impact (D4I) provided support for finalizing the final report. The survey was funded by the Government of Bangladesh and the U.S. Agency for International Development (USAID).

The 2022 BHFS assessed government, non-government, and private healthcare facilities across all eight administrative divisions of Bangladesh. This survey collected data from six types of public facilities, private hospitals with at least 10 beds, and NGO clinics or hospitals, totaling 1,557 health facilities and 6,209 healthcare providers. The 2022 BHFS serves as one of the crucial resources for assessing the performance of the 4th Health, Population, and Nutrition Sector Program (HPNSP) and guiding the development of the 5th HPNSP. The BHFS 2022 provides estimates for four indicators of the Results Framework of the 4th HPNSP. Furthermore, the survey serves as a key data source for tracking the progress of Bangladesh's health system. We hope that the data will support policymakers and program managers in monitoring and developing programs and strategies to enhance health, family planning, and nutrition services in Bangladesh.

This report is a result of relentless effort, dedication, support, and involvement of a large number of institutions and individuals. A Technical Working Committee (TWC)—with representatives from NIPORT; Ministry of Health and Family Welfare (MOHFW); University of Dhaka; icddr,b; D4I; and USAID/Bangladesh provided technical guidance in all aspects to the implementation of the survey. A Stakeholder Advisory Committee (SAC) was formed consisting of experts from governmental, non-governmental, and international entities, along with professionals and researchers working in health, nutrition, and population sectors. Their expert contributions were absolutely vital in significantly improving the quality of the survey. I am immensely grateful to all the dedicated professionals whose tireless efforts led to the successful completion of the survey.

My heartfelt thanks go out to the staff of health facilities who willingly participated and provided the information for this survey. I extend my sincere gratitude to all the organizations that played a vital role in the successful completion of the 2022 BHFS. I would like to commend the professionals of NIPORT's Research Unit for successful completion of the survey. We are indebted to icddr,b for their diligent field monitoring, ensuring the quality of the data collected. I am also thankful to ACPR for completing the data collection in a professional manner. I wish to extend my gratitude to the honorable Secretary of the Medical Education and Family Welfare Division, Ministry of Health and Family Welfare, Government of Bangladesh, for their invaluable guidance and support throughout every stage of the survey's implementation. Finally, USAID/Bangladesh deserves special thanks for providing financial support for the survey.

(Dr. Ashrafi Ahmad)

ACKNOWLEDGMENT





Mohammad Ahsanul Alam Director (Research) & Line Director, OP-TRD National Institute of Population Research and Training (NIPORT) Medical Education and Family Welfare Division Ministry of Health and Family Welfare

It is with great pleasure that I announce the publication of the final report of the fifth Bangladesh Health Facility Survey (2022 BHFS). This accomplishment would not have been possible without the collaborative efforts and contributions of all the individuals and organizations involved in this survey. First and foremost, I would like to extend my gratitude to all the professionals at the National Institute of Population Research and Training (NIPORT) for their outstanding leadership in conducting the survey. The constant support and guidance from the Government of Bangladesh were pivotal in achieving the success of this national survey.

I am deeply grateful to the International Centre for Diarrheal Disease Research, Bangladesh (icddr,b), for providing comprehensive technical assistance in every aspect of the survey, from questionnaire development to the publication of the final report. I would like to acknowledge their meticulous field monitoring and quality control efforts, which were vital in maintaining the integrity of the data collection. I also wish to express my sincere thanks to Data for Impact (D4I) for their crucial role in finalizing the report.

A Technical Working Committee (TWC) consisting of representatives from NIPORT, Ministry of Health and Family Welfare (MOHFW), University of Dhaka, icddr,b, and USAID/Bangladesh provided comprehensive technical guidance throughout the survey's implementation. Additionally, a Stakeholder Advisory Committee (SAC) comprised of experts from government, non-government, and international organizations, along with researchers, statisticians, epidemiologists, demographers, and health system specialists in the health, nutrition, and population sectors, contributed their valuable insights during the survey. Their expert opinions were crucial in the implementation process and in finalizing the report. I would like to extend my sincere appreciation to all the individual researchers and experts who contributed to the final report of the 2022 BHFS. Special thanks are due to the Directorate General of Health Services (DGHS), Directorate General of Family Planning (DGFP) for their active involvement and support.

I also wish to express my sincere appreciation to the healthcare providers and staff of the 1,557 health facilities who participated in the survey. Their cooperation and willingness to share information were essential in painting a comprehensive picture of Bangladesh's healthcare system. Additionally, I commend the field staff, data collection teams, quality control officers and all other personnel involved in the survey for their relentless efforts and professionalism.

Finally, I would like to extend my deepest thanks to the Government of Bangladesh and USAID/Bangladesh for their financial support. The findings of the 2022 BHFS will undoubtedly serve as a valuable resource for evaluating and monitoring the Health, Population, and Nutrition Sector Programs (HPNSP) in Bangladesh. I hope these findings will guide policymakers, program managers, and researchers in enhancing health, family planning, and nutrition services in Bangladesh.

(Mohammad Ahsanul Alam)

STAKEHOLDER ADVISORY COMMITTEE (SAC) FOR 2022 BHFS

| Director General, NIPORT | Chairperson |
|--|--|
| Additional Secretary (Development), Medical Education and Family Welfare Division, MOHFW (or Representative) | Member |
| M.M. Reza, Chief Technical Advisor, PMMU, Azimpur | Member |
| Director General, Health Economics Unit (HEU), MOHFW (or Representative) | Member |
| Joint Secretary (Planning), Medical Education and Family Welfare Division, MOHFW | Member |
| Joint Secretary (Planning), Health Services Division, MOHFW (or Representative) | Member |
| Director (Training), NIPORT, Azimpur, Dhaka (or representative) | Member |
| Deputy Secretary, Population Planning Wing, Planning Commission | Member |
| Project Director, Urban Primary Health Care Project, Nagar Bhaban (or Representative) | Member |
| Line Director, Maternal, Newborn, Child, and Adolescent Health (MNCAH), Directorate General of Health Services (or Representative) | Member |
| Line Director, Hospital Service Management, DGHS (or Representative) | Member |
| Line Director, Planning, Monitoring and Research (PMR), Directorate General of Health Services (or Representative) | Member |
| Director, Maternal and Child Health (MCH) Services and Line Director, Maternal, Child, Reproductive and Adolescent Health (MCRAH), DGFP (or Representative) | Member |
| Line Director, Clinical Contraception Services Delivery Program (CCSDP), DGFP Services (or Representative) | Member |
| Director, Census Wing, Bangladesh Bureau of Statistics | Member |
| Project Director, Monitoring the Situation of Vital Statistics of Bangladesh (MSVSB), Project, Bangladesh Bureau of Statistics Prof. M. Nurul Islam, Pro-Vice Chancellor, World University of Bangladesh | Member |
| Prof. Sved Shahadat Hossain, ISBT, University of Dhaka | Member |
| Prof. Nitai Chakraborty, Dept. of Statistics, University of Dhaka | Member |
| Prof. Sheikh Giash Uddin, Dent. of Statistics, Jagannath University, Dhaka | Member |
| Prof. Aminul Hague Department of Population Sciences, University of Dhaka | Member |
| Dr. Mohammad Shahraar Farid, Brajaat Managament Specialist (Monitoring, Evaluation | Member |
| and Learning), OPHN, USAID Dr. Kanta Jamil, Ex. Senior M&E and Research Advisor, OPHN, USAID Dr. Ahmed Al-Sabir, Consultant, D4I, USA Dr. Shams El Arifeen, Senior Director and Senior Scientist, Maternal and Child Health | Member Member Member |
| Division, icddr,b Dr. Ahmed Ehsanur Rahman, Scientist, Maternal and Child Health Division, icddr,b President, Obstetrical and Gynaecological Society of Bangladesh (OGSB), Dhaka (or Representative) | Member Member |
| Representative, WHO, Dhaka, Bangladesh Representative, World Bank, Dhaka, Bangladesh Representative, JICA, Dhaka, Bangladesh Representative, CIDA, Dhaka, Bangladesh Representative, Concern Data Collection/Implementing Agency | Member Member Member Member Member |
| Mr. Giyasuddin Ahamed, Ex. Sr. Research Associate, NIPORT | Member |
| Mrs. Shahin Sultana, Ex. Sr. Research Associate, NIPORT | Member |
| Mr. Mohammed Ahsanul Alam, Director (Research), NIPORT | Member- Secretary |

TECHNICAL WORKING COMMITTEE (TWC) AND SAMPLING COMMITTEE FOR 2022 BHFS

Technical Working Committee (TWC)

| Mr. Mohammed Ahsanul Alam, Director (Research), NIPORT | Chairman |
|--|------------------|
| Joint Secretary (Planning), Medical Education and Family Welfare Division (or, representative) | Member |
| Prof. Syed Shahadat Hossain, ISRT, University of Dhaka | Member |
| Prof. Nitai Chakraborty, Dept. of Statistics, University of Dhaka | Member |
| Prof. Sheikh Giash Uddin, Dept. of Statistics, Jagannath University, Dhaka | Member |
| Representative Line Director, Maternal, Newborn, Child, and Adolescent Health (MNCAH), Directorate General of Health Services | Member |
| Representative Line Director, Maternal, Child, Reproductive and Adolescent Health (MCRAH), Directorate General of Family Planning | Member |
| Representative Line Director, Clinical Contraception Services Delivery Program (CCSDP), Directorate General of Family Planning | Member |
| Dr. Mohammad Shahrear Farid, Project Management Specialist (Monitoring, Evaluation and Learning), OPHN, USAID | Member |
| Dr. Kanta Jamil, Ex. Senior M&E and Research Advisor, OPHN, USAID | Member |
| Dr. Ahmed Al-Sabir, Consultant, D4I, USA | Member |
| Dr. Shams El Arifeen, Senior Director and Senior Scientist, Maternal and Child Health Division, icddr,b | Member |
| Dr. Ahmed Ehsanur Rahman, Scientist, Maternal and Child Health Division, | Member |
| icddr,b | |
| Representative, Concern Data Collection/Implementing Agency | Member |
| Mrs. Shahin Sultana, Ex. Sr. Research Associate, NIPORT | Member |
| Mr. Giyasuddin Ahamed, Ex. Sr. Research Associate, NIPORT | Member-Secretary |

Sampling Committee (SRC)

| Mr. Mohammed Ahsanul Alam, Director (Research), NIPORT | Chairman |
|---|----------|
| Dr. Ahmed Ehsanur Rahman, Scientist, Maternal & Child Health Division, icddr,b | Member |
| Prof. Nitai Chakraborty, Dept. of Statistics, University of Dhaka | Member |
| Prof. Syed Shahadat Hossain, ISRT, University of Dhaka | Member |
| Dr. Ahmed Al-Sabir, Consultant, D4I, USA | |
| Md. Abu Bakkar Siddique, Assistant Scientist, Maternal & Child Health Division, | Member |
| icddr,b | |

CONTRIBUTORS TO THE REPORT

NIPORT

Md. Shafiqul Islam Director General

Mr. Mohammed Ahsanul Alam Director (Research)

Mr. Giayasuddin Ahamed Ex. Sr. Research Associate

Mr. Zahirul Islam Statistician

DIRECTORATE GENERAL OF FAMILY PLANNING (DGFP)

Dr. Nurun Nahar Director, Maternal and Child Health Training Institute (MCHTI)

UNIVERSITY OF DHAKA

Dr. Syed Shahadat Hossain Professor of Applied Statistics and Data Science, ISRT

Ms. Gaylan Peyari Tarannum Dana Professor Department of Population Science

Ms. Jahida Gulshan Professor of Applied Statistics and Data Science, ISRT

Ms. Shahnaz Nilima Assistant Professor Department of Statistics

Ms. Eashrat Jahan Eyemoon Assistant Professor Department of Sociology

JAGANNATH UNIVERSITY

Dr. Sheikh Giash Uddin Professor Department of Statistics

USAID/BANGLADESH

Dr. Mohammad Shahrear Farid Project Management Specialist (Monitoring, Evaluation and Learning) Office of Population, Health and Nutrition

DATA FOR IMPACT

Dr. Ahmed Al-Sabir Consultant

Ms. Shusmita Khan Knowledge Management and Communication Specialist

icddr,b

Dr. Ahmed Ehsanur Rahman Scientist Maternal and Child Health Division

Md. Abu Bakkar Siddique Assistant Scientist Maternal and Child Health Division

Dr. Shafiqul Ameen Assistant Scientist Maternal and Child Health Division

Dr. Twaha Mansurun Haque Program Manager Maternal and Child Health Division

Dr. Ashiquzzaman Project Research Physician Maternal and Child Health Division

Ms. Anindita Saha Statistical officer Maternal and Child Health Division

Professor Nitai Chakraborty Consultant

ACRONYMS AND ABBREVIATIONS

| ACE | Angiotensin-converting Enzyme |
|-----------|--|
| ACPR | Associates for Community and Population Research |
| ADB | Asian Development Bank |
| AHI | Assistant Health Inspectors |
| AMTSL | Active Management of the Third Stage of Labor |
| ANC | Antenatal care |
| ARB | Angiotensin receptor Blocked |
| ARI | Acute Respiratory Infection |
| BCC | Behavioral Change Communication |
| BDHS | Bangladesh Demographic and Health Survey |
| BDR | Border Guard Bangladesh |
| BEmOC | Basic Emergency Obstetric Care |
| BENAP | Bangladesh Every Newborn Action Plan |
| BHFS | Bangladesh Health Facility Survey |
| BMDC | Bangladesh Medical & Dental Council |
| BMMS | Bangladesh Maternal Mortality and Health Care Survey |
| BNC | Bangladesh Nursing Council |
| BNHA | Bangladesh National Health Accounts |
| BP | Blood Pressure |
| BRAC | Bangladesh Rural Advancement Committee |
| CARE | Cooperative for Assistance and Relief Everywhere |
| CBHC | Community Based Health Care |
| CC | Community Clinic |
| CEmOC | Comprehensive Emergency Obstetric Care |
| CHCP | Community Health Care Providers |
| CHX | Chlorhexidine |
| CPR | Contraceptive Prevalence Rate |
| C-section | Cesarean Section |
| СТ | Computed Tomography |
| CVD | Cerebrovascular Disease |
| D&C | Dilation and Curettage |
| D4I | Data for Impact |
| DD-FP | Deputy Director-Family Planning |
| DDS | Drugs and Dietary Supply |
| DGDA | Directorate General of Drug Administration |
| DGFP | Directorate General of Family Planning |
| DGHED | Directorate General of Health Engineering Department |
| | |

| DGHEU | Directorate General of Health Economics Unit |
|---------|--|
| DGHS | Directorate General of Health Services |
| DH | District Hospital |
| DHS | Demographic and Health Survey |
| DNS | Directorate of Nursing Services |
| DOTS | Directly Observed Treatment, Short-course |
| DPT | Diptheria-Partussis-Tetanus |
| DS | Double Strength |
| ЕМО | Emergency Medical Officer |
| EmOC | Emergency Obstetric Care |
| EmONC | Emergency Obstetric and Neonatal Care |
| EPI | Expanded Immunization Program |
| ETAT | Emergency Triage Assessment and Treatment |
| FDC | Fixed-dose Combination |
| FNAC | Fine Needle Aspiration Cytology |
| FP | Family Planning |
| FPI | Family Planning Inspector |
| FWA | Family Welfare Assistant |
| FWV | Family Welfare Visitor |
| GDP | Gross Domestic Product |
| GK | Gonoshashthaya Kendra |
| GOB | Government of Bangladesh |
| HA | Health Assistants |
| Hb | Hemoglobin |
| HFS | Health Facility Survey |
| HI | Health Inspectors |
| HIV | Human Immunodeficiency Virus |
| HLD | High Level Disinfection |
| HPN | Health, Population and Nutrition |
| HPNSDP | Health, Population and Nutrition Sector Development Programme |
| HPNSP | Health, Population and Nutrition Sector Programme |
| HPV | Human Papilloma Virus |
| HSS | Health System Strengthening |
| icddr,b | International Center for Diarrhoeal Disease Research, Bangladesh |
| IHFAN | International Health Facility Assessment Network |
| IMCI | Integrated Management of Childhood Illness |
| IMO | Indoor Medical Officer |
| IMPAC | Integrated Management of Pregnancy and Childbirth |
| IUCD | Intrauterine Contraceptive Device |

| KMC | Kangaroo Mother Care |
|---------|---|
| LAM | Lactational Amenorrhea Method |
| LARC/PM | Long-acting Reversible Contraceptives/Permanent Methods |
| LBW | Low Birth Weight |
| LED | Light Emitting Diode |
| LGD | Local Government Division |
| MA | Medical Assistant |
| MAM | Moderate Acute Malnutrition |
| MBBS | Bachelor of Medicine, Bachelor of Surgery |
| MBDC | Mycobacterium Disease Control |
| MCH-FP | Maternal, Child Health, and Family Planning |
| MCHTI | Maternal and child Health Training Institute |
| MCWC | Mother and Child Welfare Center |
| MDG | Millennium Development Goal |
| MDR-TB | Multidrug-Resistant Tuberculosis |
| MMR | Maternal Mortality Rate |
| MNC&AH | Maternal, Neonatal, Child, and Adolescent Health |
| MO | Medical Officer |
| MOHFW | Ministry of Health and Family Welfare |
| MOLGRDC | Ministry of Local Government, Rural Development and Cooperatives |
| MSH | Management Science for Health |
| MUAC | Mid Upper Arm Circumference |
| MVA | Manual Vacuum Aspiration |
| NCD | Non-communicable Disease |
| NEMEW | National Electro-medical & Engineering Workshop |
| NGO | Non-Governmental Organization |
| NIPORT | National Institute of Population Research and Training |
| NMR | Neonatal Mortality Rate |
| NNHP | National Newborn Health Program |
| NSF | Newborn Signal Function |
| NTP | National Tuberculosis Control Program |
| OBS/GYN | Obstetrics/Gynecology |
| ORS | Oral Rehydration Solution |
| PAC | Post-abortion Care |
| PH | Private Hospital |
| PHC | Primary Health Care |
| PMMU- | |
| MOHFW | Program Management and Monitoring Unit of Ministry of Health and Family Welfare |
| PNC | Postnatal care |

| PPFP | Postpartum Family Planning |
|---------|---|
| PPP | Private Public Partnership |
| | |
| QA | Quality Assurance |
| QC | Quality Control |
| | |
| RCHCIB | Revitalization of Community Health Care Initiatives in Bangladesh |
| RD | Rural Dispensary |
| RDM | Research for Decision Makers |
| RMO | Resident Medical Officer |
| RR-TB | Rifampicin-Resistant Tuberculosis |
| RSE | Relative Standard Error |
| | |
| SAC | Stakeholders Advisory Committee |
| SACMO | Sub-assistant Community Medical Officer |
| SAM | Severe Acute Malnutrition |
| SARA | Service Availability and Readiness Assessment |
| SBA | Skilled Birth Attendant |
| SDG | Sustainable Development Goal |
| SMF | State Medical Faculty |
| SPA | Service Provisional Assessment |
| STI | Sexually Transmitted Infection |
| SVRS | Sample Vital Registration System |
| | |
| TAB | Tablet computer |
| TB | Tuberculosis |
| TEMO | Transport and Equipment Maintenance Organization |
| TLCA | TB leprosy Control Assistant |
| TRD | Training, Research and Development |
| TWC | Technical Working Committee |
| | |
| UFPO | Upazila Family Planning Officer |
| UH&FPO | Upazila Health & Family Planning Officer |
| UHC | Upazila Health Complex |
| UHFWC | Union Health & Family Planning Center |
| ULB | Urban Local Bodies |
| UPHCP | Urban Primary Health Care Service Project |
| UPHCSDP | Urban Primary Health Care Services Delivery Project |
| US | United States |
| USAID | United States Agency for International Development |
| USC | Union Sub-center |
| USC/RD | Union sub-center/Rural dispensary |
| | |
| WHO | World Health Organization |

READING AND UNDERSTANDING TABLES FROM THE 2022 BANGLADESH HEALTH FACILITY SURVEY (BHFS)

The Bangladesh Health Facility Survey (BHFS) 2022 final report is based on approximately 84 tables of data. For quick reference, they are located at the end of each chapter and can be accessed through links in the pertinent text (electronic version). Additionally, this reader-friendly version features 109 figures that clearly highlight trends, subnational patterns, and background characteristics. The text has been simplified to highlight key points in bullets.

While the text and figures featured in each chapter highlight some of the most important findings from the tables, not every finding can be discussed or displayed graphically. For this reason, 2022 BHFS data users should be comfortable reading and interpreting tables.

The following pages introduce the organization of 2022 BHFS tables, the presentation of background characteristics, and a brief summary of sampling and understanding denominators. In addition, this section provides some exercises for users as they practice their new skills in interpreting 2022 BHFS tables.



Example 1: Availability of Basic Client Services

Among all facilities, the percentages offering indicated basic client services and all basic client services, by background characteristics, 2022 BHFS

| Table 3.2: Availability of basic client services | | | | | | | | | | |
|--|--|--|--|--|--|--|--|--|--|---|
| Among all facilities, the percentages offering the indicated basic client services and all basic client services, by background characteristics, Bangladesh HFS 2022 | | | | | | | | | | |
| 3 Background characteristic | Child curative care | 2 Child growth monitoring services | Child vaccination services | Any modern methods of family planning | Antenatal care services | Normal delivery | All basic client services with normal delivery ¹ | All basic client services without normal delivery | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 99.1 100.0 95.1 100.0 | 92.9 98.4 68.1 98.1 | 90.9 95.2 57.0 98.5 | 97.1 93.5 95.1 98.2 | 99.8 100.0 99.0 100.0 | 97.4 100.0 87.1 99.5 | 85.1 88.7 40.1 95.5 | 85.8 88.7 42.0 96.0 | 41 4 7 30 | 319 62 100 157 |
| Union-level public facilities UHFWC USC/RD | 95.3 94.2 98.1 | 74.2 78.2 63.6 | 79.1 79.2 79.1 | 83.5 96.0 50.3 | 98.3 100.0 93.7 | 49.4 59.2 23.4 | 32.6 39.2 15.2 | 52.3 59.3 33.7 | 310 225 85 | 434 293 141 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 94.5 88.4 56.6 | 92.4 81.9 22.5 | 97.5 71.9 3.0 | 82.1 74.8 25.0 | 99.8 100.0 72.4 | 4.7 33.5 89.0 | 3.6 16.6 1.1 | 70.4 50.5 1.1 | 994 29 183 | 488 127 189 |
| Location Urban Rural | 65.7 94.6 | 36.9 88.1 | 22.4 92.7 | 38.6 82.6 | 79.3 99.3 | 81.7 16.6 | 11.2 11.6 | 16.2 66.2 | 235 1,322 | 514 1,043 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Svlbet | 92.7 97.4 83.8 76.7 95.1 95.7 88.4 | 87.6 87.5 75.7 76.2 73.9 84.7 82.7 | 84.5 85.1 80.0 70.9 82.5 87.3 84.0 | 82.0 80.9 70.7 70.3 75.4 79.3 74.4 | 98.8 97.4 95.4 90.5 97.8 97.7 98.0 | 19.2 30.6 30.1 30.4 26.4 17.1 31.6 | 11.1 16.0 11.2 8.8 10.2 8.0 20.4 | 63.6 71.5 49.1 50.1 55.4 64.5 59.2 | 111 291 317 210 214 193 99 | 152 244 264 208 212 190 |
| Mymensingh | 94.7 | 78.8 | 87.2 | 78.6 | 97.5 | 18.2 | 7.1 | 59.3 | 122 | 151 |
| Total | 90.2 | 80.4 | 82.1 | 75.9 | 96.3 | 26.5 | 11.5 | 58.7 | 1,557 | 1,557 |
| Total excluding CCs | 82.7 | 59.1 | 54.9 | 65.0 | 90.1 | 65.0 | 25.4 | 38.0 | 563 | 1,069 |
| Total public | 94.8 | 88.2 | 93.1 | 82.9 | 99.5 | 17.8 | 12.8 | 66.7 | 1,345 | 1,241 |
| I otal public excluding CCs | 95.7 | 76.4 | 80.5 | 85.1 | 98.5 | 55.0 | 38.8 | 56.2 | 351 | 753 |

¹ Basic client services include outpatient curative care for sick children, child growth monitoring, facility-based child vaccination services, any modern methods of family planning, antenatal care, and normal delivery.

Step 1: Read the title and subtitle, highlighted in orange in the table above. They provide a brief description of the information contained in the table. In this case, the table is about the provision of basic client services in health facilities assessed in the 2022 BHFS.

Step 2: Scan the column headings—highlighted in green in Example 1. They describe how the information is categorized. In this table, there are ten columns of data. The first six columns represent one basic client service— child curative care, child growth monitoring, child vaccination, any modern methods of family planning, antenatal care, and normal delivery. The seventh column shows what percent of facilities have ALL six basic client services, while the eighth column shows the percent of facilities with all basic client services EXCEPT normal delivery. Note that the two last columns, in gray, list the weighted and unweighted numbers of facilities surveyed. The weighted numbers are the denominators, that is, the total number of facilities surveyed for each topic and each background characteristic. In this case, 1,557 facilities were surveyed. Of these 1,557 facilities,

41 are district and upazila level-public facilities (319 unweighted), 310 are union-level public facilities (434 unweighted), 994 are community clinics (488 unweighted), 29 are NGO clinics/hospitals (127 unweighted), and 183 are private hospitals (189 unweighted). For more on weighting, see Example 3.

Step 3: Scan the row headings—the first vertical column highlighted in **teal** in Example 1. These show the different ways the data are divided into categories based on background characteristics. In this case, the table presents availability of basic client services by facility type, urban-rural location, and division. (Note that facility type is further divided into subcategories). Most of the tables in the 2022 BHFS will be divided into these same categories.

Step 4: Look at the row at the bottom of the table highlighted in red. These figures represent the totals of public facilities, or the percent of public facilities that offer each of the six services, and the percent of public facilities that offer ALL services. This table shows that services for child curative care (94.8%) are widely available in public facilities in Bangladesh, while 12.8% of all public facilities provide ALL six basic client services.

Note that there are three additional rows below the total which present what the total would be if community clinics (CCs) are excluded, total for public facilities, and total for public facilities excluding CCs. The reason for showing the total public excluding CCs is that CCs are the lowest level of facilities that are supported by community health care providers, and often are least likely to offer all of the health services and/or to have the items necessary for providing a service if they offer it.

Step 5: To find out what percent of health facilities in Rangpur division offer any modern methods of family planning, draw two imaginary lines, as shown in the table above in purple. This shows that 79.3% of health facilities in Rangpur division offer any modern methods of family planning.

By looking at patterns by background characteristics, we can see how the provision of basic client services varies across Bangladesh. Knowing how client services vary among different types of health facilities can help program planners and policy makers determine which interventions need to be prioritized where.

*For the purpose of this document data are presented exactly as they appear in the table including decimal places. However, the text in the remainder of this report rounds data to the nearest whole percentage point.

Practice: Use the table in Example 1 above to answer the following questions:

a) Are normal delivery services more likely to be offered in urban or rural facilities?

b) In which division are child curative care services LEAST widely available?

c) Facilities in which facility type category are most likely to offer ALL six basic client services?

basic client services with normal delivery at 95.5%.

c) District and upazila public facilities—85.1%. Within this category, upazila health complexes (UHC) are most likely to offer all

b) Khulna division—76.7%.

a) Urban-81.7% of urban facilities offer normal delivery services, compared to 16.6% of rural facilities.

Answers:

Example 2: Availability of Antenatal Care (ANC) Services

Based on all health facilities and a subset of health facilities

| Table 6.1: Availability of antenatal care services | | | | | | | | | |
|---|---------------|--------------------------|-------------------------|---|------------------------------------|------------|------------------------------|--|--|
| Among all facilities, the percentage offering antenatal care (ANC) services and, among facilities offering ANC services, the percentage offering the service on the indicated number of days per week, by background characteristics, Bangladesh HFS 2022 | | | | | | | | | |
| 3 | 2 | | | Percentage of facilities offering ANC where ANC services are | | Number of | Number of | | |
| | Percentage of | Number of | Number of | offered on indicated days ¹ | | facilities | facilities | | |
| Background | offer ANC | facilities (weighted) | facilities (unweighted) | every day ¹ | Provides every day ¹ | (weighted) | offering ANC (unweighted) | | |
| Facility type | | | | | | | | | |
| District and upazila public facilities | 99.8 | 41 | 319 | 9.6 | 90.4 | 41 | 318 | | |
| DH | 100.0 | 4 | 62 | 6.5 | 93.5 | 4 | 62 | | |
| MCWC | 99.0 | 7 | 100 | 22.4 | 77.6 | 7 | 99 | | |
| UHC | 100.0 | 30 | 157 | 7.0 | 93.0 | 30 | 157 | | |
| Union-level public facilities | 98.0 | 310 | 434 | 35.4 | 64.6 | 305 | 427 | | |
| UHFWC | 100.0 | 225 | 293 | 35.8 | 64.2 | 225 | 293 | | |
| USC/RD | 93.7 | 85 | 141 | 34.1 | 65.9 | 80 | 134 | | |
| Community clinic (CC) | 99.8 | 994 | 488 | 16.7 | 83.3 | 992 | 487 | | |
| NGO clinic/hospital | 100.0 | 29 | 127 | 3.6 | 96.4 | 29 | 127 | | |
| Private nospital | 72.4 | 183 | 189 | 16.1 | 83.9 | 132 | 141 | | |
| Location | | | | | | | | | |
| Urban | 79.3 | 235 | 514 | 13.0 | 87 | 187 | 467 | | |
| Rural | 99.3 | 1,322 | 1,043 | 21.0 | 79 | 1,312 | 1,033 | | |
| Division | | | | | | | | | |
| Barishal | 98.8 | 111 | 152 | 13.9 | 86.1 | 110 | 148 | | |
| Chattogram | 97.4 | 291 | 244 | 15.5 | 84.5 | 283 | 238 | | |
| Dhaka | 95.4 | 317 | 264 | 23.3 | /6./ | 303 | 254 | | |
| Knuina | 90.5 | 210 | 108 | 12.3 | 86.8 | 190 | 197 | | |
| Rajsnani | 97.8 | 214 | 212 | 24.0 | 75.5 | 209 | 205 | | |
| Sulbot | 97.5 | 193 | 130 | 24.0 | 79.0 | 07 | 122 | | |
| Mymensingh | 90.0 | 122 | 150 | 22.0 | 78.6 | 118 | 1/6 | | |
| Mymenangn | 57.5 | 122 | 101 | 21.4 | 10.0 | 110 | 140 | | |
| Total | 96.3 | 1,557 | 1,557 | 20.0 | 80.0 | 1,499 | 1,500 | | |
| Total excluding CCs | 90.1 | 563 | 1,069 | 26.4 | 73.6 | 507 | 1,013 | | |
| Total public | 99.5 | 1,345 | 1,241 | 20.7 | 79.3 | 1,338 | 1,232 | | |
| Total public excluding CCs | 98.5 | 351 | 753 | 32.3 | 67.7 | 346 | 745 | | |
| ¹ Every day refers to all working days when the facility is open. | | | | | | | | | |

Step 1: Read the title and subtitle. In this case, the table is about two separate groups: (a) all health facilities and (b) facilities that offer antenatal care (ANC) services.

Step 2: Identify the two panels. First, identify the columns that refer to all facilities (a). Then, isolate the columns that refer only to facilities that offer ANC services (b).

Step 3: Scan the row headings to identify the background characteristics. In this table, availability of ANC services is presented by facility type, urban-rural location, and division.

Step 4: Find the total for public facilities and denominators for each panel. What percent of public facilities in Bangladesh offer ANC services? It's 99.5%. Now look at the (weighted) denominator for the second panel (b). How many public facilities in Bangladesh surveyed in the 2022 BHFS offer ANC services? It's 1,338, which represents 99.5% of the 1,345 public facilities in the survey sample. The second panel is a subset of the first panel.

When reading and using the 2022 BHFS tables, be sure to identify which group of facilities is being displayed. For example, look at the first column in panel b. It is NOT correct to say that 79.3% of public facilities provide ANC services every day. It IS correct to say that 79.3% of *public facilities offering ANC services* provide these services every day.

Practice: Use the table in Example 2 above to answer the following questions:

a) What category of health facility is least likely to offer ANC services?

b) What percent of public health facilities offering ANC services provide ANC services but not every day? c) How many union-level public facilities offer ANC services? (Hint: the weighted gray columns show the denominators, or the number of health facilities in the sample or subsample.)

d) What percent of community clinics (CCs) offering ANC services provide ANC services every day?

e) In which division are ANC services most likely to be provided every day by facilities offering ANC services?

e) ANC services are most likely to be offered every day of the week by facilities offering ANC services in Khulna division (86.8%).

day. 3% of CCs offering ANC services offer these services every day.

c) 305 union-level public facilities offer AUC services.

b) 20.7% of public facilities offering ANC services do not offer these services every day.

a) Private hospitals are least likely to offer ANC services (69.4%).

Answers:
Example 3: Understanding Sampling Weights in 2022 BHFS Tables

In the 2022 BHFS, the sample is a group of facilities that have been selected from a list of all health facilities in the country. The sample represents the entire health system, that is, all health facilities in Bangladesh. Most countries want to collect data and report information that represents facilities in the entire country as well as facilities in subnational areas like divisions.

In the case of the BHFS, researchers also want to know about health facilities of different types, as well as facilities run by different managing authorities (for example, public, NGO, or private). We want the sample of district hospitals (DHs) surveyed to represent all district hospitals in the country, and we want the community clinics (CCs) sampled to represent all CCs. However, there are many more CCs than DHs in Bangladesh. If we chose a simple random sample of health facilities, we will only get a few DHs but hundreds of CCs. A simple random sample would likely not include enough DHs for meaningful analysis. To solve this

| Percent distribution and number of surveyed facilities, by background characteristics, Bangladesh HFS 2022 | | | | | | |
|--|---|---|--|--|--|--|
| Background | Weighted percent | Number of facilities surveyed | | | | |
| characteristic | surveyed facilities | Weighted | Unweighted | | | |
| Facility type District and upazila public facilities DH MCWC UHC | 2.6 0.3 0.4 1.9 | 41 4 7 30 | 319 62 100 157 | | | |
| Union-level public facilities UHFWC USC/RD | 3 19.9 14.5 5.4 | 2 ³¹⁰ ²²⁵ ⁸⁵ | 1 434 293 141 | | | |
| Community clinic (CC) | 63.8 | 994 | 488 | | | |
| NGO clinic/hospital ¹ Private hospital | 1.9 11.8 | 29 183 | 127 189 | | | |
| Location Urban Rural | 15.1 84.9 | 235 1,322 | 514 1,043 | | | |
| Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 7.1 18.7 20.4 13.5 13.7 12.4 6.4 7.8 | 111 291 317 210 214 193 99 122 | 152 244 264 208 212 190 136 151 | | | |
| Total | 100 | 1,557 | 1,557 | | | |
| Total excluding CCs | - | 563 | 1,069 | | | |
| Total public | - | 1,345 | 1,241 | | | |
| Total public excluding CCs | - | 351 | 753 | | | |
| ¹ The NGO category includes facilities run by local government. "-" means not applicable | | | | | | |

Table 2.4: Distribution of surveyed facilities, by background characteristics

problem, some types of health facilities are oversampled.

For example, let's say that we have enough money to visit 1,557 health facilities for a survey and you want to produce results that are representative of health facilities in Bangladesh as a whole and representative of each different type of health facility (as in the table above). The types of health facilities are not evenly distributed: there are many more CCs than DHs, for instance.

A sampling statistician can determine how many facilities of each type should be surveyed in order to get reliable statistics for the specific indicators the country wants. In the case of the 2022 BHFS, the teal column (1) shows the actual number of facilities selected and interviewed in each type, ranging from 62 DHs to 488 CCs. The sampling statistician assures us that these are enough facilities to get reliable results for each type of facility.

But now there is a new challenge. With this distribution of facilities by type, some facility types are overrepresented and some types are underrepresented. For example, the unweighted column tells us that 62 DHs were surveyed, which equals 4% of all facilities in the sample (1,557). In reality, DHs comprise less than 1% of all health facilities in Bangladesh. On the other hand, 488 CCs were surveyed, which equals 31% of the facilities in the sample. In actuality, about 64% of health facilities in Bangladesh are CCs. Would our survey show the true state of health facilities in Bangladesh if we used this sample distribution? No; this unweighted distribution of facilities does not accurately represent the true state of health facilities.

In order to get statistics that are representative of the entire health system in Bangladesh, the distribution of different types of health facilities in our sample needs to resemble the true distribution of health facility types in Bangladesh. DHs, for example, should only contribute a small amount to the national total. Likewise, CCs should contribute much more to the total. Statisticians mathematically calculate a "weight" which is used to adjust the number of health facilities from each facility type, so that each health facility type's contribution to the total is proportionate to the actual distribution of health facilities in Bangladesh. The numbers in the **purple column (2)** represent the "weighted" values. The weighted values can be smaller or larger than the unweighted values. The total sample size of 1,557 facilities has not changed, but the distribution of the facilities by facility type has been adjusted to represent their contribution to the total number of facilities in the country.

How do statisticians weigh each category? They recalculate the categories to reflect the real distribution of facilities in the country. If you were to compare the **green column (3)** to the actual distribution of facilities in Bangladesh, you would see that facilities of each type surveyed are contributing to the total sample with the same weight that they contribute to the total number of facilities in Bangladesh. The weighted number of health facilities in the survey now accurately represents how many facilities are CCs—64% of the facilities in Bangladesh.—and how few facilities are DHs—less than 1% of all health facilities in Bangladesh.

With sampling and weighting, it is possible to survey enough facilities to provide reliable statistics at both the national and division level, without distorting the overall distribution of facilities within the country. Both weighted and unweighted numbers are shown in each of the BHFS tables. Don't be distressed if the weighted numbers seem low—they may actually represent a larger number of facilities as shown in the unweighted number column.

Note: Data from the actual, unweighted number of facilities are used for analysis. For example, even though the weighted number of district and upazila public facilities is only 41, the data collected from all 319 of this facility type are used for analysis. The only difference is that the results are weighted after analysis to represent information from district and upazila public facilities in the proportion that they exist in the country

BANGLADESH



BANGLADESH



1.1 HEALTH STATUS IN BANGLADESH

B angladesh has made remarkable progress in the health sector over the past decades. Health status has improved substantially in Bangladesh, leading to a reduction in neonatal, infant, and under-five mortality, maternal deaths, and an increase in life expectancy of six years for men and eight years for women between 2005 and 2022 (Sample Vital Registration System [SVRS] 2022). The under-five mortality dropped from 88 deaths per 1,000 live births in 2004 to 31 deaths per 1,000 live births in 2022 (Bangladesh Demographic and Health Survey [BDHS] 2022). Maternal mortality also declined by 52%, from 322 deaths per 100,000 live births to 156 deaths per 100,000 live births between 2001 (BMMS 2001) and 2015 (SVRS 2022). The country has achieved the Millennium Development Goal (MDG) 4 target for under-five mortality (48 deaths per 1,000 live births) and is expected to achieve its Sustainable Development Goal (SDG) 3 targets for under-five mortality (less than 25 deaths per 1,000 live births) and maternal mortality (less than 70 maternal deaths per 1,000 live births) by 2030. The decline in mortality is attributed to changing fertility behavior, which has led to a decrease in the total fertility rate from 3.3 children per woman in 1999-2000 to 2.3 children per woman in 2022 (BDHS 2022).

The country also made significant strides in reducing the burden of communicable diseases, such as tuberculosis, malaria, and HIV/AIDS, and has increased access to essential medicines and vaccines. The country's response to the COVID-19 pandemic has been commendable, with the government taking proactive measures to contain the spread of the virus and ensure access to healthcare services for all.

1.2 HEALTH, POPULATION, AND NUTRITION SECTOR PROGRAM

The 4th Health, Population and Nutrition Sector Program (4th HPNSP) of Bangladesh is a comprehensive health program that aims to improve the health and well-being of all citizens of Bangladesh by expanding access to quality and equitable healthcare in a healthy and safe living environment. This sector program was implemented by Ministry of Health and Family Welfare (MOHFW) in January 2017 and has been extended to June 2024. The program has several objectives, including improving access to and utilization of evidence-based high-impact health, population and nutrition-related services, strengthening systems to support service delivery, and effective stewardship and governance. The program also aims to promote healthy lifestyles and practices at community and facility levels. The program is contributing significantly to the achievement of the SDGs in Bangladesh.

Strategic Priorities for the 4th HPNSP

The 4th HPNSP in Bangladesh has formulated eight strategic objectives to cover all aspects of the health sector, including governance and stewardship, health systems, health services, and the wider determinants of health. The program's implementation plan outlines the following sector-specific strategic priorities:

- Strengthen governance and stewardship of the public and private health sectors
- Undertake institutional development for improved performance at all levels of the system
- Provide sustainable financing for equitable access to healthcare for the population and accelerated progress toward universal health coverage
- Strengthen the capacity of the MOHFW's core health systems (financial management, procurement, and infrastructure development)

- Establish a quality health workforce available to all through public and private health service providers
- Improve health measurement and accountability mechanisms and build a robust evidence base for decision making
- Improve equitable access to and utilization of quality health, nutrition, and family planning services
- Promote healthy lifestyle choices and a healthy environment

1.3 HEALTH SERVICE DELIVERY SYSTEM OF BANGLADESH

The health service delivery system of Bangladesh is a complex network of public health departments, nongovernmental organizations (NGOs), and private institutions. The MOHFW formulates national policies, plans, and decisions for health care and education. These policies are implemented by authorities at the national and community levels. The MOHFW was divided into two divisions in March 2017: the Health Services Division and the Medical Education and Family Welfare Division. The Health Services Division is responsible for policy making, management, and maintenance of nursing care, health financing, and other health-related matters. The Medical Education and Family Welfare Division is responsible for policies regarding medical education, family planning, medical colleges and universities, and registration of births and deaths.

There are 10 implementing authorities under the MOHFW that deliver HPN services through various operational plans under HPNSDP. Those are the Directorate General of Health Services (DGHS), DGFP, Directorate of Nursing Services (DNS), NIPORT, Directorate General of Drug Administration (DGDA), Directorate General of Health Engineering Department (DGHED), Directorate General of Health Economics Unit (DGHEU), Transport and Equipment Maintenance Organization (TEMO), and the National Electro-medical & Engineering Workshop (NEMEW).

The MOHFW and its relevant regulatory bodies (Bangladesh Medical & Dental Council [BMDC]; Bangladesh Nursing Council [BNC]; State Medical Faculty [SMF]; Homeo, Unani and Ayurvedic Board; and Bangladesh Pharmacy Council) have indirect control over the system of public, NGO, and private sector health care providers. The delivery of health services in urban areas, including primary health care services, is mandated by the Ministry of Local Government, Rural Development and Cooperatives (MOLGRDC).

1.3.1 Management Structure and Health Facilities Under DGHS

The DGHS is the largest implementing authority under the MOHFW in Bangladesh, with over 100,000 officers and staff members. In addition to operating health care delivery systems across the country, the DGHS provides technical assistance to the MOHFW for new programs and interventions and for improvements in existing ones. The health care delivery system under the DGHS extends from the national down to the community level and is implemented under regular revenue setups and development programs.

The DGHS in Bangladesh has a six-tier health care infrastructure that includes national, divisional, district, upazila (sub-district), union, and ward levels. The national level has institutions for public health functions, postgraduate medical education/training, and specialized treatment of patients. Each division is governed by a divisional director for health, with assistance from deputy directors and assistant directors. Divisional headquarters have one infectious disease hospital and one or more medical colleges, each with a hospital. Some divisional headquarters also have general hospitals and institutes of health technology, which provide tertiary-level care. The civil surgeon is the district health manager responsible for delivering secondary and primary care services. Each district has a district hospital, most of which are managed by superintendents, while others are headed by civil surgeons. Some district headquarters have medical colleges with attached hospitals, medical assistant training schools, and nursing training institutes. The Upazila Health & Family Planning Officer (UH&FPO) manages all public health programs, including primary health care services in the upazila and the

upazila hospital, which has 30 to 100 beds. Most upazilas have hospitals, except where the district headquarters are located; in those upazilas, the district hospital provides hospital services. At the union level, there are three types of health facilities: rural dispensaries (RDs), union sub-centers (USCs), and UHFWCs. Each union health facility employs a medical doctor and has sub-assistant community medical officers (SACMOs) who provide health services. Only outpatient services are available at union-level facilities.

The MOHFW in Bangladesh has established 13,948 community clinics (CCs) at the ward level, each serving around 6,000-12,000 people (Health Bulletin 2020). The CCs provide a basic health care package to the community, including maternal and child health care, reproductive health and family planning services, immunization, nutrition education, micronutrient supplementation, health education and counseling, communicable disease control, treatment for minor ailments and first aid, and referral to higher-level health centers. Some CCs also have a community skilled birth attendants (SBAs) and offer services for normal delivery. The CCs are governed by a management committee comprised of 13 to 17-members (of which, at least one-third are female members) selected from the communities served by the CC, including local government representatives. Additionally, community support groups work as unpaid community health volunteers and assist the management committee and community clinics.

The MOHFW in Bangladesh has recruited and trained 13,948 full-time Community Health Care Providers (CHCPs) to operate the CCs and provide quality health care. The CHCPs use internet-connected laptop computers to update local health data in an online database that is used for evidence-based decision making and planning. In addition to the CHCPs, the existing domiciliary staff members of the DGHS and Directorate General of Family Planning (DGFP) provide service to the CCs, with the DGHS staff and DGFP staff alternate their availability in three-day schedules each week. In total, there are 26,538 sanctioned posts of domiciliary workers under DGHS: 20,908 health assistants (HAs), 4,220 assistant health inspectors (AHIs), and 1,410 health inspectors (HIs). As of now, 74.72% posts were filled up (Health Bulletin 2020).

| Table 1.1 | Table 1.1 Secondary and primary health care facilities in public sector | | | | | | | |
|-----------|---|-----------------------------------|-------------------|--|--|--|--|--|
| Level | Type of facility | Type of service | No. of facilities | | | | | |
| District | District/general hospital Other secondary hospital | Hospital Hospital | 59 6 | | | | | |
| | | Total of district-level hospitals | 65 | | | | | |
| | Mother and child welfare center | Hospital | 62 | | | | | |
| Upazila | Upazila health complex (100-bed) | Hospital | 3 | | | | | |
| | Upazila health complex (50-bed) | Hospital | 345 | | | | | |
| | Upazila health complex (31-bed) | Hospital | 65 | | | | | |
| | Upazila health complex (10-bed) | Hospital | 11 | | | | | |
| | | Total of upazila health complexes | 424 | | | | | |
| | Hospitals outside health complexes | | | | | | | |
| | 50-bed hospital | Hospital | 2 | | | | | |
| | 31-bed hospital | Hospital | 7 | | | | | |
| | 30-bed hospital | Hospital | 3 | | | | | |
| | 25-bed hospital | Hospital | 1 | | | | | |
| | 20-bed hospital | Hospital | 38 | | | | | |
| | 10-bed hospital | Hospital | 13 | | | | | |
| | Mother and child welfare center | Hospital | 98 | | | | | |
| Union | Union-level facilities under DGHS | | | | | | | |
| | Union subcenter | Outpatient only | 1,312 | | | | | |
| | Union health and family welfare center | Outpatient only | 87 | | | | | |
| | Mother and child welfare center | Hospital | 54 | | | | | |
| | Union health and family welfare | e Hospital | 3.924 | | | | | |
| | center-upgraded Union health an | d Outpatient only | - / - | | | | | |
| | family welfare center | | | | | | | |
| Ward | Community clinic | Outpatient only | 13,948 | | | | | |
| Source: H | lealth Bulletin 2020; Renovation and Recons | struction plan, DGFP 2023 | | | | | | |

1.3.2 Management Structure and Health Facilities Under DGFP

The DGFP in Bangladesh implements a Maternal and Child Health/Family Planning (MCH-FP) service delivery system that extends from the district to the community level, with an extensive network of health facilities, satellite clinics, and domiciliary workers. In each district, there is an MCH-FP clinic at the District Sadar Hospital and a separate Mother and Child Welfare Center (MCWC) that delivers MCH-FP services. The MCWCs provide antenatal care (ANC), normal delivery, postnatal care (PNC), and child health care including an expanded program on immunization (EPI) and all contraceptives, including sterilization through medical officer (clinic), family welfare visitors (FWVs), and dai (nurses). The FWVs at the MCH-FP clinics in the District Sadar Hospital provide ANC, normal delivery (limited facilities for "at-risk" cases); PNC; all contraceptives, including sterilization; and follow-up services. Similar services are also available in model clinics attached to the public medical college hospitals. The Deputy Director-Family Planning (DD-FP) is responsible for overall management of the MCH-FP program at the district level, while the Divisional Director (FP) is the program head at the divisional level.

The MCWCs provide services at the upazila level and below that are similar to those offered at the district level and have an equivalent staffing profile. Almost all Upazila Health Complexes (UHCs) have MCH-FP units that provide ANC; normal delivery; PNC; child health care, including an EPI; health education; and all contraceptives, including sterilization. The UHC MCH-FP units are led by a medical officer (MCH-FP) along with the FWVs. The Upazila Family Planning Officer (UFPO) is the manager responsible for overall management of the FP program at the upazila level, while the medical officer (MCH-FP) is the upazila manager responsible for MCH service delivery. Almost all union-level facilities can provide MCH-FP services. There are 3,924 Union Health and Family Welfare Centers (UHFWCs) under the DGFP. The SACMOs and FWVs provide ANC (with an emphasis on screening for "at-risk" pregnancies and referral), normal delivery at upgraded facilities, PNC, child health care including health education, and contraceptives that include clinical methods, and treatment of general patients. The FWVs are also assigned in some Union Sub-Centers (USCs) and Rural Dispensaries (RDs) under the Directorate General of Health Services (DGHS) to deliver MCH-FP services. The FWVs and SACMOs are responsible for delivering MCH-FP services and health education at more than 30,000 satellite clinics, which are organized monthly at the community level by FP fieldworkers.

The DGFP operates a network of community-based fieldworkers for domiciliary FP services. The fieldworkers, known as family welfare assistants (FWAs), distribute contraceptives and provide counseling, health education, and referrals for MCH-FP services at the household level. The FWAs also offer services at CCs and EPI outreach sites. The FWAs' activities are supervised by the union-level family planning inspector (FPI), who works under DGFP.

1.3.3 Urban Primary Health Care Services Delivery Project (UPHCSDP)

Bangladesh's urban population is growing at an annual rate of 3.27%. As of 2016, 37.2% of the population lived in urban areas, and this number is expected to increase to 44% by 2030. This rapid expansion has placed significant pressure on health services and facilities in urban areas. Although the country's rural areas are supported by a strong public sector primary health care network, there is a significant lack of similar arrangements in urban areas. The MOHFW is the main player in the health sector, with the mandate for providing primary health care in urban areas falling under the administrative jurisdiction of the Ministry of Local Government, Rural Development, and Cooperatives. To address this issue, the Local Government Division of the Government of Bangladesh has taken the initiative to provide primary health care services to urban people through partnerships between urban local bodies and non-governmental organizations (NGOs), with the financial support of the Asian Development Bank (ADB). Since 1998, the Local Government Division (LGD), with the support of ADB, has been assisting Urban Local Bodies (ULBs) in providing PHC services in urban areas under UPHCP-I and II and UPHCSDP. The UPHCSDP project will continue to provide services until 2023 and has been extended to June 2024 under UPHCSDP-II covering 11 city corporations and 18 municipalities (17.70 million population) throughout Bangladesh. The key features of this project are: (1) an excellent model of Public Private Partnership (PPP); (2) safety net provision for the economically disadvantaged (at least 30% of services are free of cost); (3) involvement of local government and community; (4) client-friendly health service delivery model; (5) easy accessibility and low cost; (6) adolescent corner for adolescent counseling; (7) satellite clinic services, near to the doorsteps of those in the community; (8) developed capacity of ULBs to respond to emergency health needs; (9) physical infrastructure for future sustainability; and (10) 24-hour ambulance service.

1.3.4 NGO Health Programs

The NGO sector has played a significant role in improving the Bangladesh health sector. The involvement of NGOs in the country's health sector dates back to the pre-independence period. NGOs have been providing quality health services at either reasonable cost or sometimes free-of-charge. Health services provided by NGOs include not only preventive and curative services but also advocacy, research, awareness raising for healthseeking behavior, nutrition, environmental health, behavior change communication (BCC), and others. Notable NGOs that have contributed to the health sector of Bangladesh are BRAC, CARE, the Diabetic Association of Bangladesh, Gonoshasthaya Kendra (GK), Kumudini Welfare Trust, Save the Children, and World Vision. BRAC is currently the hub of the world's largest non-government network of community health workers. Across Bangladesh, this network of 50,000 women are taking essential healthcare to the doorsteps of families and reaching the last mile, standing by for 80 million people each year. The community health workers, known as Shasthya Shebikas, provide health promotion sessions and educate families on nutrition, safe delivery, family planning, immunizations, hygiene, and water and sanitation. They also use this time to sell health products, such as basic medicine, sanitary napkins, and soap. These NGOs have been instrumental in improving the health sector of Bangladesh by providing quality health services to millions of people in Bangladesh. Their contributions have been significant in reducing the burden of disease, improving maternal and child health, and promoting healthy behaviors among the people of Bangladesh.

1.3.5 Private Health Sector

The private health sector in Bangladesh has grown rapidly in recent years and has made significant contributions to the overall health of the country. The country's private health sector comprises large and small commercial companies, professionals, and informal providers. It includes health services provided at hospitals, nursing and maternity homes, clinics, diagnostic facilities, and pharmacies. As of June 2020, there were 16,444 registered private hospitals, clinics, diagnostic centers, and blood banks in Bangladesh. The total number of beds in the registered private hospitals and clinics was 91,537 (Health Bulletin 2020). Local pharmaceutical companies manufacture and distribute most of the targeted products, and the majority of Bangladeshis usually obtain their drugs from the private sector. Even among rural residents and the urban economically disadvantaged , the most common means of obtaining drugs is through direct, out-of-pocket purchases from private sources.

1.4 HEALTH FINANCING

According to Bangladesh National Health Accounts (BNHA), the total health expenditure in 2020 was Taka777 billion, accounting for 2.8% of Gross domestic product (GDP). In the same year, government spending on health was Taka179.74 billion, representing 23% of total health expenditures (0.66% of GDP). The general trend in total health expenditures as a proportion of GDP during 1997–2020 has been rising and averaged around 3%. However, government spending as a share of GDP has fallen and remained around less than one percent (<1%) throughout the period—falling even more sharply in 2020, to 0.66%. This is contrary to the international trend for public expenditures on health to increase as a share of GDP as per capita income rises. It is noteworthy, however, that household spending contributed to increased total health expenditures.

Public health spending as a proportion of GDP was 0.66% in 2020. This was lower than all other South Asian countries, with the exception of Afghanistan which had the lowest figures in the region (5%). Public hospital spending as a proportion of total government health expenditures rose from 42% in 1997 to 49% in 2020. In contrast, public hospital spending as a proportion of total hospital spending declined from 65% in 1997 to 48% in 2020, indicating a growing share of private sector spending (including NGOs).

The proportion of government spending on curative care increased while the share of preventive care decreased between 1997 and 2020. In 1997, the respective share was 36% and 17% of total public health expenditures. In 2020, the share of curative care increased to 47%, while the share of preventive care dropped to 15%. However, it is noteworthy that the share of preventive care as a proportion of public health expenditures was higher between 2003 (39%) and 2015 (33%). In contrast, the share of curative care as a proportion of public health expenditures was lower by comparison, increasing only post-2015 (28%). This trend could be attributed to the government's upscaling of disease control and prevention programs during this period, with a massive upscaling of hospitals and other facilities designated for curative healthcare taking place since 2015.

According to BNHA, the government's share of total health expenditures was on a declining trend from 1997–2020, remaining at less than 1% of GDP throughout the period. Further analysis of health spending revealed that the proportion of government health expenditures on hospitals decreased during the same period. Although preventive care received a higher proportion of public health expenditures between 2003 and 2015, curative healthcare received a higher share post-2015 (Public Expenditure Review 1997-2020).

To increase financial protection for the entire population, while decreasing out-of-pocket payments at the point of service, the MOHFW has three strategic objectives under the Health Care Financing Strategy 2012-2032 (HEU 2012):

- Generate more resources for effective health services.
- Improve equity and increase health care access, especially for the economically disadvantaged and vulnerable.
- Enhance efficiency in resource allocation and utilization.

1.5 HEALTH STEWARDSHIP

The government is taking measures to improve the quality and equity of health services, particularly for the poor and disadvantaged. However, further steps are required to make health services more responsive to the population's needs and demands. The governance of the system remains insufficient in terms of providing equitable services, and regulatory mechanisms, especially within the for-profit private sector, must be strengthened. Decentralization of authority and responsibility, combined with proper accountability at all levels, must be institutionalized, given the availability of a functional local government system.

METHODOLOGY

2.1 OVERVIEW

B angladesh has an extensive network of public, private, and nongovernmental organization (NGO) facilities for providing basic health services. The Government of Bangladesh (GOB) is committed to strengthening health systems and improving the quality of care. In the past two decades, investments in health facilities have expanded services and improved access to quality care. Subsequently, Bangladesh has achieved immense success in increasing health service coverage and utilization. Although service utilization is assessed and monitored intensively with various data sources, the improvements in health systems and quality of care have not been monitored systematically. Under the Health, Population and Nutrition Sector Program (HPNSP) January 2017- June 2024, the Ministry of Health and Family Welfare (MOHFW) is committed to periodic assessments of health systems and quality of care provided by various health facilities. As part of this assessment effort, the National Institute of Population Research and Training (NIPORT) was entrusted with conducting the 2022 Bangladesh Health Facility Survey (BHFS) under the Training, Research, and Development (TRD) operational plan of the HPNSP.

The 2022 BHFS is an assessment of health care facilities in the formal sector of Bangladesh. The survey provides information on the availability of basic and essential health care services¹ and the readiness of health facilities to provide quality services to clients. The 2022 BHFS is the fifth survey of its kind and follows the 2009, 2011, 2014 and 2017 BHFS surveys. As with the 2014 and 2017 surveys, the 2022 BHFS collected information from all types of health facilities managed by the government, NGO clinics and hospitals, and private for-profit hospitals with 10 or more beds in all 64 districts of the country. The 2022 BHFS tools adopted the 2017 BHFS which is based on standardized questionnaires from the Service Provision Assessment (SPA) component of USAID's Demographic and Health Surveys (DHS) Program to collect information on the availability of services and the preparation of facilities to provide quality, effective, and efficient services to clients. The 2022 BHFS focused primarily on the service readiness indicators that were jointly developed and proposed by WHO, USAID, the World Bank, the International Health Facility Assessment Network (IHFAN), and other stakeholders. The 2014, 2017, and 2022 BHFS data are not strictly comparable to the 2009 and 2011 survey results because the 2014, 2017, and 2022 BHFS used a different set of questionnaires and defined indicators slightly differently.

The 2022 BHFS provides information on child health, maternal and newborn care, family planning (FP), and services for selected non-communicable diseases (NCDs) (diabetes and cardiovascular diseases) and tuberculosis. The newborn signal function module has been added to the tools for the first time in the 2022 BHFS. For each of these services, the 2022 BHFS assessed whether components considered essential for quality service delivery were present and functioning. In general, the components that were assessed are those that are commonly considered important to various programs supported by the government and development partners. Unlike the 2017 BHFS, the 2022 BHFS also assessed whether more sophisticated components were present, such as higher-level diagnostic and treatment modalities or support systems for health services that are usually introduced after basic-level services have been put in place.

¹ Basic and essential health care services of interest in the 2022 BHFS included child health, family planning, maternal and newborn care, non-communicable diseases, and tuberculosis.

2.2 INSTITUTIONAL FRAMEWORK AND OBJECTIVES OF THE 2022 BHFS

2.2.1 Institutional Framework

The 2022 BHFS was conducted under the authority of NIPORT. icddr,b provided overall technical assistance under the USAID-funded Research for Decision Makers (RDM) project and Data for Impact (D4I) provided support for finalizing the final report. Associates for Community and Population Research (ACPR), a private research agency, collected the data. Financial support for the survey was provided by the Government of Bangladesh and the United States Agency for International Development (USAID). A Stakeholders Advisory Committee (SAC) and a Technical Working Committee (TWC) oversaw all policy and technical issues related to the survey.

2.2.2 Objectives of the 2022 BHFS

The main objectives of the 2022 BHFS were to:

- Assess the availability of health services, including maternal and child health, family planning, diabetes, cardiovascular disease, tuberculosis, and nutrition services.
- Ascertain general preparedness of the health facilities and availability of basic amenities, equipment, laboratory services, essential medicines, standard precautions for infection control, and human resources at the facilities.
- Assess service-specific readiness of health facilities to provide maternal, newborn, and child health care; family planning services; treatment of diabetes, cardiovascular disease, and tuberculosis, measured in terms of the average score of WHO-recommended minimum items required to provide quality services.
- Compare findings among facility types and managing authorities.

2.3 DATA COLLECTION METHODS

The 2022 BHFS used two types of data collection tools:

- Facility Inventory Questionnaire
- Health Care Provider Interview Questionnaire

Both the Facility Inventory and Health Care Provider Interview questionnaires were loaded onto Android-based tablets.

The **Facility Inventory Questionnaire** obtained information on the availability of each priority service and the preparedness of the facilities to provide the service. The questionnaire also collected information on the availability of specific items (including their location and functional status), components of support systems (e.g., logistics, maintenance, and management), and facility infrastructure, including the service delivery environment. The data collectors interviewed the person who is most knowledgeable about the facility and its services. If another person or provider needed to provide specific information, the data collectors consulted that person or provider to obtain the information. However, the data collectors considered only observed items as available in the facility.

The Facility Inventory Questionnaire was organized into three modules:

• **Module 1** collected information on service availability and included two sections.

- Module 2 collected information on general facility readiness. This module included seven sections that covered topics such as facility infrastructure (i.e., sources of water and electricity), staffing, health management information systems, health statistics, processing of instruments for re-use, health care waste management, availability of basic supplies and equipment, laboratory diagnostic capacity, and medicines and commodities.
- Module 3 collected information on service-specific readiness. This module has 13 sections including specific service areas such as child vaccination, growth monitoring, child curative care, FP, adolescent health, nutrition, antenatal care (ANC), delivery and newborn care, tuberculosis, NCDs, caesarean delivery, blood grouping and compatibility, blood transfusion services, and general facility cleanliness.

The **Health Care Provider Interview Questionnaire** collected information from a sample of health service providers. The data included qualifications, training, experience, continuing education/ training, supervision received, and perceptions of the service delivery environment.

2.4 SURVEY IMPLEMENTATION

2.4.1 Questionnaire Adaptation

The 2022 BHFS questionnaires are based on the 2017 BHFS survey instruments of the SPA core questionnaires developed by The DHS Program. The questionnaires were adapted for 2022 BHFS to focus on Bangladesh health services in consultation with technical specialists and experts from the MOHFW, DGHS, DGFP, and icddr,b. The questionnaire adaptation for the 2022 BHFS took place during July and August 2022 in Dhaka. A series of workshops were conducted with representatives from the MOHFW, NIPORT, DGHS, DGFP, icddr,b, ACPR, Program Management and Monitoring Unit of Ministry of Health and Family Welfare (PMMU-MOHFW), USAID, D4I, University of Dhaka, Jagannath University and other key stakeholders to review and update the questionnaires based on the country context. After being prepared in English, both the Facility Inventory and Health Care Provider Interview questionnaires were translated into Bangla.

Android-based tablets were employed for data collection, using an android application developed with native Java programming. These tablets stored data locally using SQLite database. A central server running SQL server 2019 was used to store and manage the collected data from all tablets. The central service hosted a web application and web services developed with ASP.NET, enabling reliable data synchronization over the internet.

2.4.2 Pre-test

A two-day pretesting was conducted after adaptation and translation of the hardcopy questionnaires and one-day field-testing was completed using the program for Android-based tablets. The pre-test was conducted in master training during August 2022 in non-sampled areas of Dhaka. Eleven medical doctors were recruited (four from icddr,b and seven from ACPR). During the pre-test data collection, health facilities within Dhaka district were surveyed over a 3-day period to test and refine the survey instruments and the computer programs. After the pre-test, the questionnaires and computer programs were finalized for the main survey assessment. At the end of the pre-test, 10 of 11 medical doctors who received the 2 weeks of training served as "master trainers" of the training of data collection enumerators. They also served as independent field monitors including interview observation and quality control (QC) during the main survey.

2.4.3 Main Training

The main training for the 2022 BHFS was conducted from September 6-25, 2022, in Dhaka. Eighty enumerators consisting of 40 interviewers [paramedics (medical assistants)] and 40 team leaders [medical doctors] were

recruited for data collection. They received training as interviewers in the application of survey instruments and Android-based data collection apps.

The training included classroom lectures and discussions, practical demonstrations, mock interviews, role-plays, and field practice. The trainees were also given daily homework to conduct mock interviews among themselves with the survey tools. The first week of training was dedicated exclusively to training interviewers on the use of paper questionnaires. This phase included two days of field practice using hardcopy of tools on 17 and 18 September 2022 to ensure that the trainees understood both the content of the questionnaires and approaches for organizing themselves while working in a health facility.

During the third week of training, trainees were introduced to tablet and then transitioned to the use of tabletbased program and instruments for both the Facility Inventory and Health Care Provider Interview questionnaires. Initially, this was done with completed paper questionnaires from the facilities visited during the pre-test. During the fourth training week, the trainees practiced all questionnaire types and tablet-based approaches in teams and pairs. A one-day field practice for interviewers was conducted using tablet-based instruments for familiarization. The training was conducted by master trainers and trainers from NIPORT, icddr,b, and ACPR.

2.4.4 Data Collection

After the training, 40 data collection teams with two interviewers each were formed. One interviewer (medical doctor) from each team was assigned to the role of team leader. Data collection was conducted from 27 September to 8 December 2022. On average, data collection took one day for one interviewer to complete the UH&FWCs, USCs, and CCs. However, it took 2 days for a team (two interviewers) to cover the survey for DHs and UHCs; and took 1 day for a team (two interviewers) to complete MCWC, NGO, and private health facilities.

Fieldwork supervision was led by icddr,b. The medical doctor master trainers and trainers of the master trainer formed the thirteen field supervision teams. The field supervision teams conducted frequent visits to their assigned data collection teams to review work, observe the interview, and conduct QC of data collection. In addition, supervisory teams and professionals from NIPORT and icddr,b periodically conducted field visits to monitor the data collection.

Observation is an essential tool to ensure quality data collection. The approach, methods, and techniques were assessed through observation. A standard observational checklist was established for unbiased and specific observation of the data collectors' approach. Simultaneously, a QC questionnaire tool, shortened, based on the BHFS 2022 inventory questionnaire was established to recheck and compare data by QC monitoring physicians in facilities which were assessed immediately through TAB-based QC tools. The QC physicians provide feedback to the interviewers just after QC observation and interviews are completed. The feedback aimed and was largely focused to improve the day-to-day data collection quality.

2.4.5 Data Management

The data collection team used two tablets for data collection in DHs, UHCs, MCWC, Private hospitals/clinics, and NGOs. Both team members are allowed to collect the data for any of the sections of the Facility Inventory Questionnaire and the Health Care Provider Interview Questionnaire. All tablets were equipped with internet access. When an interviewer syncs their tablet, the paired tablet shows all the collected data by the teammate. Initially, the background colour of all sections in the tablet shows white. When a section's data collection is complete, it shows green. But it is shown in red when the section is not complete. The team leader then reviewed the file and conducted consistency and structural checks on the data to identify any errors or missing information. When a team was satisfied that data collection and entry were complete for the facility, the team sent the data to

the icddr,b server via the internet. However, it shows the message in red during the final submission of the data when an entire section or some of the particular questions of a section are left blank while the facility reportedly offers that specific service. In this case, the team revisited the sections and finally submitted the data to the servers.

Data received from the field were reviewed and checked at icddr,b with the technical assistance of icddr,b experts. If any inconsistencies or errors were identified by the icddr,b experts, the data collection team members reviewed and rechecked the data. If necessary, the team visited the specific service facility again to obtain and resend the correct data to the central office.

Data editing and management activities commenced in October 2022 at the beginning of field work and ended in January 2023 after completing the data collection in the field.

2.4.6 Data Analysis and Report Preparation

The BHFS 2022 tabulation plan is based on the 2017 BHFS tabulation. The preparation of the tables for the report was conducted from February through April 2023. Feedback from NIPORT, the BHFS TWC, and the BHFS SAC members informed revision of the analysis plan.

Two conventions were observed during the analysis of the 2022 BHFS data:

- First, unless otherwise indicated, the 2022 BHFS considered only those items observed by the interviewers to be available.
- Second, in a majority of facilities, multiple health care providers contribute to the services received by clients. The health care provider who ultimately assessed the client, made the final diagnosis, and prescribed any treatment was therefore identified as the primary provider for the particular service.
- The final report was prepared with input from professionals from NIPORT, Maternal and Child Health Training Institute (MCHTI), USAID, D4I, and University of Dhaka, Jagannath University, and icddr,b with a continuous technical oversight by icddr,b.

2.5 SAMPLING

2.5.1 Sampling Methodology

The sample for the 2022 BHFS was a stratified random sample of 1,640 health facilities designed to provide representative results for Bangladesh, for the different facility types and different management authorities, and for each of the eight divisions of the country (**Map 2**).

The sample for the 2022 BHFS covered all types of registered health facilities in all eight divisions of the country: Barisal, Chittagong, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur, and Sylhet. The survey was designed to report results separately for the eight divisions and the six types of public health facilities included: community clinics (CCs), union subcenters/rural dispensaries (USC/RDs), union health and family welfare centers (UHFWCs), upazila health complexes (UHCs), mother and child welfare centers (MCWCs), and district hospitals (DHs). Results are also reported separately for NGO clinics and hospitals, and private hospitals with 10 or more beds.

2.5.2 Sampling Frame

A list of 21,943 registered health facilities, provided by the MOHFW, was used as a sampling frame for sample selection. The allocation of the 2022 BHFS sample took the type of facility and divisional distribution of the health facilities into account. At the same time, other factors such as indicator precision, at the national level (<5% relative standard error) and also at domain level- facility type division (12.5% relative standard error). The distribution of registered health facilities by division and by facility type is presented in **Table 2.1**.

2.5.3 2022 BHFS Sample Size and Sample Allocation

Like previous health facility surveys, the 2022 BHFS is designed to provide survey results separately for the eight administrative divisions, six types of public facilities, NGO static clinics/hospitals, and private hospitals with 10 or more beds. The sample size determination was achieved by controlling the survey precision at the division level and by facility type at the national level. The formula used for the sample size calculation is

$$n = \frac{\left(1 - p\right)}{\varepsilon^2 p}$$

where \mathcal{E} is the desired survey precision, which is the relative standard error for estimating a proportion *p*. By controlling the relative standard error (RSE) for an indicator at the 30% level (*p* = 0.30) within 12.5% of relative standard error ($\mathcal{E} \leq 0.125$) at facility type and division level, and 5% nonresponse rate, a sample of 1,640 health facilities was allocated based on a power allocation between sampling domains (health facility types and divisions). **Table 2.2** presents the allocation of the 1,640 health facilities by division and facility type. All of the 100 MCWCs (All MCWCs at district level and other MCWCs which were found functioning at the time of the listing of facilitates) and 62 DHs were included in the 2022 BHFS survey.

2.5.4 Sample of Health Facilities and Outcomes

The BHFS sample was a combination of census (for DHs and MCWCs) and random samples (for other facility types). **Table 2.3** presents an analysis of the 1,640 sampled health facilities according to the outcomes of the visits to those facilities. In this table and in subsequent tables in the report, the total for facilities is shown including and excluding CCs. However, the 2022 BHFS also reports the value of each characteristics in 'total public facilities excluding CCs'. The reason for showing the total public excluding CCs is that CCs are the lowest level of facilities that are supported by community health care providers, and often are least likely to offer all of the health services and/or to have the items necessary for providing a service if they offer it. As shown, data were successfully collected from 95% of the 1,640 sampled facilities (a total of 1,557 facilities).

About 5% of sampled facilities could not be interviewed for various reasons, but mainly because some facilities were closed or not operational at the time of the survey. Consequently, BHFS data are available for 1,557 facilities. **Table 2.4** shows the distribution of surveyed facilities by background characteristics.

2.5.5 Sample of Health Service Providers

The sample of health service providers was selected from providers who were present in the facility on the day of the assessment and who provided services assessed by the 2022 BHFS. For purposes of the BHFS data collection, a health service provider was defined as a person who provides consultation services, counseling, health education, or laboratory services to clients. Health workers were not eligible for interviewing if they only take measurements or complete registers and do not provide any type of professional client services. In each

facility, the aim was to interview an average of eight providers who provided the range of services being assessed. In facilities with less than eight health care providers, all providers present on the day of the visit were interviewed. In facilities with more than eight providers, efforts were made to interview eight providers, including all of those who provided information for any section of the Facility Inventory Questionnaire and others involved in delivering services for which information was collected in the BHFS. In a few cases, the staff members present on the day of the assessment may not have been representative of the staff that usually provides the services being assessed. The health care provider data were weighted based on interviewed staff versus posted staff during the analysis to account for a particular qualification in a facility type differential from over-sampling or under-sampling of providers.

Table 2.5 shows the distribution of health care providers who were interviewed with the Health Care Provider Interview Questionnaire, by background characteristics and provider qualification. A total of 6,209 providers were interviewed, with the highest percentage of interviews conducted in private clinics/hospitals (33%) and UHCs (25%).

METHODOLOGY

LIST OF TABLES

- Table 2.1 Bangladesh health facility distribution by division and health facility type
- Table 2.2 Health facility sample allocation by division and health facility type
- Table 2.3 Results of facility contacts by background characteristics
- Table 2.4 Distribution of surveyed facilities by background characteristics
- Table 2.5 Distribution of interviewed providers

| Table 2.1: Bangladesh health facility distribution by division and health facility type | | | | | | | | | |
|---|--------|--------|-------|-----|------|----|----------------------------|-------|--------|
| Division | СС | USC/RD | UHFWC | UHC | MCWC | DH | NGO clinic /hospital | PH | Total |
| Barishal | 1.131 | 48 | 226 | 34 | 10 | 6 | 12 | 96 | 1.563 |
| Chattogram | 2,469 | 238 | 654 | 87 | 18 | 11 | 103 | 518 | 4.098 |
| Dhaka | 2,650 | 248 | 603 | 77 | 16 | 14 | 131 | 735 | 4,474 |
| Khulna | 1,706 | 113 | 455 | 51 | 18 | 10 | 46 | 562 | 2,961 |
| Mymensingh | 1,245 | 93 | 229 | 31 | 6 | 3 | 16 | 91 | 1,714 |
| Rajshahi | 1,976 | 194 | 389 | 59 | 16 | 7 | 49 | 322 | 3,012 |
| Rangpur | 1,892 | 189 | 410 | 49 | 12 | 7 | 29 | 136 | 2,724 |
| Sylhet | 933 | 73 | 207 | 31 | 6 | 4 | 22 | 121 | 1,397 |
| Bangladesh | 14,002 | 1,196 | 3,173 | 419 | 102 | 62 | 408 | 2,581 | 21,943 |

DH = district hospital, MCWC = mother and child welfare center, UHC = upazila health complex, UHFWC = union health and family welfare center, USC/RD = union subcenter/rural dispensary, CC = community clinic, NGO = nongovernmental organization, PH = private hospital

| Table 2.2: Health facility sample allocation by division and health facility type | | | | | | | | | | | | |
|---|-----|--------|-------|-----|------|-----|-----------|-----|-------|--|--|--|
| | | | | | | NGO | | | | | | |
| | | | | | | | clinic | | | | | |
| Division | CC | USC/RD | UHFWC | UHC | MCWC | DH | /hospital | PH | Total | | | |
| Barishal | 47 | 15 | 29 | 15 | 10 | 6 | 10 | 20 | 152 | | | |
| Chattogram | 75 | 24 | 41 | 24 | 19 | 11 | 23 | 27 | 244 | | | |
| Dhaka | 80 | 22 | 49 | 26 | 16 | 14 | 25 | 32 | 264 | | | |
| Khulna | 65 | 20 | 38 | 21 | 17 | 10 | 15 | 22 | 208 | | | |
| Mymensingh | 49 | 15 | 31 | 16 | 6 | 3 | 11 | 20 | 151 | | | |
| Rajshahi | 66 | 21 | 40 | 21 | 15 | 7 | 17 | 25 | 212 | | | |
| Rangpur | 62 | 12 | 38 | 20 | 12 | 7 | 13 | 26 | 190 | | | |
| Sylhet | 44 | 12 | 27 | 14 | 5 | 4 | 13 | 17 | 136 | | | |
| Bangladesh | 488 | 141 | 293 | 157 | 100 | 62 | 127 | 189 | 1,557 | | | |

DH = district hospital, MCWC = mother and child welfare center, UHC = upazila health complex, UHFWC = union health and family welfare center, USC/RD = union subcenter/rural dispensary, CC = community clinic, NGO = nongovernmental organization, PH = private hospital

Table 2.3: Results of facility contacts by background characteristics

Percent distribution of sampled facilities according to result of visit of the survey team to the facility, by background characteristics, Bangladesh HFS 2022

| | | | | Closed/ | | | Number of |
|-------------------------------|-----------|---------------|---------|-------------|-------|---------|------------|
| Background | | Respondent | | not yet | | Total | facilities |
| characteristic | Completed | not available | Refused | operational | Other | percent | surveyed |
| Facility type | | | | | | | |
| District and upazila | | | | | | | |
| public facilities | 99.1 | 0.0 | 0.0 | 0.9 | 0.0 | 100.0 | 322 |
| DH | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 62 |
| MCWC | 97.1 | 0.0 | 0.0 | 2.9 | 0.0 | 100.0 | 103 |
| UHC | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 157 |
| Union-level public | | | | | | | |
| facilities | 94.8 | 0.0 | 0.0 | 5.2 | 0.0 | 100.0 | 458 |
| UHFWC | 97.3 | 0.0 | 0.0 | 2.7 | 0.0 | 100.0 | 301 |
| USC/RD | 89.8 | 0.0 | 0.0 | 10.2 | 0.0 | 100.0 | 157 |
| Community clinic | | | | | | | |
| (CC) | 99.6 | 0.0 | 0.0 | 0.4 | 0.0 | 100.0 | 490 |
| NGO clinic/hospital | 79.9 | 0.0 | 0.0 | 20.1 | 0.0 | 100.0 | 159 |
| Private hospital | 89.6 | 0.0 | 0.0 | 10.4 | 0.0 | 100.0 | 211 |
| Division | | | | | | | |
| Barishal | 98.7 | 0.0 | 0.0 | 1.3 | 0.0 | 100.0 | 154 |
| Chattogram | 94.6 | 0.0 | 0.0 | 5.4 | 0.0 | 100.0 | 258 |
| Dhaka | 96.0 | 0.0 | 0.0 | 4.0 | 0.0 | 100.0 | 275 |
| Khulna | 92.9 | 0.0 | 0.0 | 7.1 | 0.0 | 100.0 | 224 |
| Rajshahi | 96.4 | 0.0 | 0.0 | 3.6 | 0.0 | 100.0 | 220 |
| Rangpur | 91.8 | 0.0 | 0.0 | 8.2 | 0.0 | 100.0 | 207 |
| Sylhet | 94.4 | 0.0 | 0.0 | 5.6 | 0.0 | 100.0 | 144 |
| Mymensingh | 95.6 | 0.0 | 0.0 | 4.4 | 0.0 | 100.0 | 158 |
| Total | 94.9 | 0.0 | 0.0 | 5.1 | 0.0 | 100.0 | 1,640 |
| Total excluding CCs | 93.0 | 0.0 | 0.0 | 7.0 | 0.0 | 100.0 | 1,150 |
| Total public | 97.7 | 0.0 | 0.0 | 2.3 | 0.0 | 100.0 | 1,270 |
| Total public excluding CCs | 96.6 | 0.0 | 0.0 | 3.4 | 0.0 | 100.0 | 780 |

Note: The percentages in some rows may not add up to 100% due to rounding. DH = district hospital, MCWC = mother and child welfare center, UHC = upazila health complex, UHFWC = union health and family welfare center, USC/RD = union subcenter/rural dispensary, CC = community clinic

Table 2.4: Distribution of surveyed facilities, by background characteristics

Percent distribution and number of surveyed facilities, by background characteristics, Bangladesh HFS 2022 $\,$

| Background | Weighted percent distribution of surveyed | Number of facilities surveyed | | | |
|--|--|-------------------------------|------------|--|--|
| characteristic | facilities | Weighted | Unweighted | | |
| Facility type District and upazila public facilities | 2.6 | 41 | 319 | | |
| DH | 0.3 | 4 | 62 | | |
| MCWC | 0.4 | 7 | 100 | | |
| UHC | 1.9 | 30 | 157 | | |
| Union-level public facilities | 19.9 | 310 | 434 | | |
| UHFWC | 14.5 | 225 | 293 | | |
| USC/RD | 5.4 | 85 | 141 | | |
| Community clinic (CC) | 63.8 | 994 | 488 | | |
| NGO clinic/hospital ¹ | 1.9 | 29 | 127 | | |
| Private hospital | 11.8 | 183 | 189 | | |
| Location | | | | | |
| Urban | 15.1 | 235 | 514 | | |
| Rural | 84.9 | 1,322 | 1,043 | | |
| Division | | | | | |
| Barishal | 7.1 | 111 | 152 | | |
| Chattogram | 18.7 | 291 | 244 | | |
| Dhaka | 20.4 | 317 | 264 | | |
| Khulna | 13.5 | 210 | 208 | | |
| Rajshahi | 13.7 | 214 | 212 | | |
| Rangpur | 12.4 | 193 | 190 | | |
| Sylnet | 0.4 | 99 | 130 | | |
| Wymensingn | 7.0 | 122 | 151 | | |
| Total | 100.0 | 1,557 | 1,557 | | |
| Total excluding CCs | - | 563 | 1,069 | | |
| Total public | - | 1,345 | 1,241 | | |
| Total public excluding CCs | - | 351 | 753 | | |

¹The NGO category includes facilities run by local government. "-" means not applicable

Table 2.5: Distribution of interviewed providers

Percent distribution and number of interviewed providers, by background characteristics and provider qualification, Bangladesh HFS 2022

| | Weighted percent distribution of | | |
|--|-------------------------------------|------------------|------------------|
| Background | interviewed _ | Number of interv | liewed providers |
| | providers | Weighted | Onweighted |
| Facility type District and upazila public | | | |
| facilities | 33.6 | 2.080 | 3.410 |
| DH | 7.9 | 489 | 896 |
| MCWC | 0.5 | 29 | 382 |
| UHC | 25.2 | 1,562 | 2,132 |
| Union-level public facilities | 7.4 | 461 | 794 |
| UHFWC | 5.3 | 332 | 515 |
| USC/RD | 2.1 | 129 | 279 |
| Community clinic (CC) | 22.4 | 1,391 | 756 |
| NGO clinic/hospital | 3.2 | 201 | 468 |
| Private hospital | 33.4 | 2,076 | 781 |
| Location | | | |
| Urban | 57.3 | 3,561 | 3,543 |
| Rural | 42.7 | 2,648 | 2,666 |
| Division | | | |
| Barishal | 6.4 | 398 | 620 |
| Chattogram | 16.7 | 1,035 | 948 |
| Dhaka | 29.6 | 1,840 | 1,143 |
| Knuina | 10.2 | 632 | 796 |
| Rajsnani | 14.3 | 885 | 862 |
| Sylbot | 10.5 | 222 | //4 |
| Mymensingh | J.2 7 1 | 322 | 490 570 |
| Wymensingn | 7.1 | | 570 |
| Provider type | | | |
| Specialist ¹ | 1.7 | 107 | 211 |
| General practitioner ² | 10.2 | 635 | 1,192 |
| Paramedics ^o | 9.2 | 568 | 1,303 |
| Field supervisors ⁵ | 23.5 | 1,460 | 1,071 |
| Medical/pharmaceutical | 0.4 | 20 | 25 |
| technician ⁶ | 22.6 | 1 300 | 608 |
| Other health providers ⁷ | 32.4 | 2 014 | 1 199 |
| | 02.1 | 2,0 | 1,100 |
| Total | 100 | 6,209 | 6,209 |
| Total excluding CCs | - | 4,818 | 5,453 |
| Total public | - | 3,932 | 4,960 |
| Total public excluding CCs | - | 2,541 | 4,204 |

¹ Specialist (consultant) medicine [including cardiology], specialist (consultant) general surgery, specialist (consultant) obstetrics/gynecology, specialist (consultant) pediatrics, specialist (consultant) psychiatry, specialist (consultant) anesthesia, or any other specialist not listed above ² Medical officer (MBBS) (any non-specialist doctor, including assistant surgeon, emergency medical officer (EMO), indoor medical officer (IMO), maternal and child health/family planning medical officer (MCH/FP), residential medical officer (RMO), regardless of designation or title) or medical officer—anesthetist or dental surgeon

³ SACMO [(sub-assistant community medical officer/medical assistant (MA)], family welfare visitor

(FWV), or paramedics in private/NGO
⁴ Nurse/midwife, matron, nursing supervisor, senior staff nurse, assistant nurse/staff nurse in

private, midwife, or DAI nurse ⁵ Health inspector or assistant health inspector

⁶ Medical technologist-laboratory or medical technologist- Expanded Program on Immunization

(EPI) ⁷ Family welfare assistant (FWA), health assistant, community health care provider (CHCP), TB leprosy control assistant (TLCA), counselor, nutritionist or health educator, other providers "-" means not applicable

FACILITY-LEVEL INFRASTRUCTURE, RESOURCES, MANAGEMENT, AND SUPPORT

Key Findings

- Almost all (≥90%) of public health facilities provide antenatal care, child curative care, child vaccination, postnatal care, adolescent health care, nutrition, and postpartum family planning services. Child growth monitoring (88%) and family planning services (83%) are available in most public facilities (Table 3.1).
- More than half (55%) of public facilities excluding CCs provide normal delivery services. About one in five public facilities offer laboratory diagnosis (23%) and non-communicable disease diagnosis or management (23%), and tuberculosis (TB) diagnosis or treatment services (24%). Very few public facilities offer blood grouping and typing (3%) and blood transfusion services (2%) (Table 3.1).
- Across all public facilities (excluding CCs), connection to the national electricity grid/pollibiddut is almost universal (97%). The availability of national electricity grid/polli-biddut in public facilities has increased from 57% to 85% between 2017 and 2022 (Table 3.3 and Figure 3.2).
- Among all public facilities, DHs (100%) and UHC (99%) have universal access to a functioning computer/laptop/tablet with internet access. (**Table 3.3** and **Figure 3.6**).
- Overall, about one in ten public facilities and 18% of public facilities excluding CCs have a functioning communication equipment (land-line or mobile phone). Very few CCs (6%), UHFWCs (8%), and USC/RDs (9%) have a functioning communication equipment. But there is some improvement in the availability across public facilities between 2017 and 2022 (Table 3.3 and Figure 3.5).
- Six basic amenities—regular electricity, an improved source of water, privacy during patient consultations, a latrine for clients, communication equipment, and a computer with Internet access—are considered essential for rendering services in health facilities. Overall, the average score for basic amenities for all public facilities is 3.0 (out of 6). District hospitals and private hospitals have the highest average score of basic amenities (5.2 out of 6), whereas CCs and USC/RDs have the lowest average score (<3 out of 6) (Table 3.3 and Figure 3.8).</p>
- The most widely available test is blood glucose, provided by 57% of public facilities (Table 3.7.1). The average capacity to conduct five basic diagnostic tests (hemoglobin, blood glucose, urine protein, urine glucose, urine pregnancy test) among public facilities is 1.0 out of 5.0 (1.8 excluding CCs). Across all facility types, availability has gradually increased over the years (Table 3.7.2 and Figure 3.10).
- NGO clinics/hospitals have the highest capacity of diagnostic testing on average (4.4 out 5.0) followed by UHCs (4.3 out 5.0) (Table 3.7.2 and Figure 3.10).
- Functional X-ray machines have increased in DHs and UHCs between 2017 and 2022. Also, functional ultrasonography machine slightly increased in DHs and substantially (three-folds) increased in UHCs during the same time period (Table 3.7.2 and Figure 3.11, 3.12).
- About one-fifth of UHCs (17%) have senior/ junior consultants (surgery) and only 5% have radiologists compared to DHs (79% and 44% respectively) (**Table 3.11.1**).
- Among public facilities, the most widely practiced method of managing sharps waste (36%) and non-sharp (19%) is offsite removal followed by open burning. Urban facilities are more likely to use offsite removal, compared to rural facilities for both types of waste (Table 3.12.1, 3.12.2 and Figure 3.14, 3.15).

3.1 BACKGROUND

n order to improve the health status of the population, a health system requires fundamental inputs and vital support systems that promote efficient and effective provision of health care services. While healthcare services can be delivered in diverse conditions, some common inputs are necessary under all conditions to ensure the quality, acceptability, and utilization of these services. These essential inputs include human resources and equipment as well as pharmaceutical and medical supplies.

This chapter reports results from the 2022 BHFS on the availability of basic health services and essential resources and on management and support systems at the facility level. The chapter is divided into the following parts:

- Availability of services: Section 3.2, including Tables 3.1 through 3.8.5 and Figures 3.1 through 3.12, describes the availability of client services at health care facilities in Bangladesh and reports on a range of indicators recommended by WHO to assess the readiness of facilities to provide good-quality client services, including the availability of basic amenities and equipment, infection control processes, diagnostic capacity, essential medicines, medicines of DDS kit and medicine supply in community clinics.
- Basic management and support: Section 3.3, including Tables 3.9 and 3.10 and Figure 3.13, considers the extent to which essential management and administrative systems are in place to support the provision of quality services, including quality assurance monitoring and supportive management practices.
- **Staffing:** Section 3.4, including **Tables 3.11.1 and 3.11.2** provide information on staffing patterns at the different facility levels.
- Medical waste management: Section 3.5, including Tables 3.12.1 and 3.12.2 and Figures 3.14 and 3.15, provides information on management of medical waste at the different facility levels.

3.2 AVAILABILITY OF SERVICES

3.2.1 Overall Availability of Specific Client Services

Policymakers and program managers can use information on the overall availability of health services in Bangladesh to identify gaps in the provision of key services. **Table 3.1** shows the percentages of all facilities that offer various client services.

- Almost all public facilities (≥90%) in Bangladesh provide antenatal care, child curative care, child vaccination (through the government's Expanded Program on Immunization [EPI], postnatal care, adolescent health care, nutrition, and postpartum family planning services (Table 3.1).
- Child growth monitoring (88%) and family planning services (83%) are available in more than fourfifths of public facilities (**Table 3.1**).
- More than one in five public facilities offer services for laboratory diagnosis (23%), non-communicable disease diagnosis or management (23%), and tuberculosis (TB) diagnosis or treatment (24%) (Table 3.1).
- Very few public facilities provide blood grouping and typing (3%) and blood transfusion services (2%) (Table 3.1).

- About one-fifth of the public facilities provide normal delivery services as compared to about four-fifths (82%) of NGOs and private facilities (**Table 3.1**).
- Most NGOs and private facilities offer laboratory services (85%) followed by cesarean delivery (83%) (Table 3.1).

3.2.2 Availability of Basic Client Services

The availability of a basic package of health services, the frequency with which these services are offered, the presence of qualified staff for their delivery, and the overall ease of access to the health care system all contribute to client utilization of services at a health facility.

The BHFS defines basic client services as the following:

- Outpatient curative care for sick children
- Child growth monitoring services
- Facility-based child vaccination services
- Provision of any modern method of family planning (FP)
- Antenatal care (ANC)
- Normal delivery

Table 3.2 and Figure 3.1 present information on the availability of FP and basic maternal and child health services.

- Almost all of the district and upazila-level facilities (97%) and UHFWCs (96%) offer modern family planning services. Modern family planning service is available in one-fourth of the private hospitals (Table 3.2).
- All DHs (100%), MCWCs (99%), UHCs (100%), UHFWCs (100%), CCs (100%), and NGOs (100%) provide antenatal care services. However, antenatal care service is available in a relatively lower proportion of private facilities (72%) (Table 3.2).
- Availability of normal delivery service is universal in DHs (100%) and UHCs (100%). Although most of the MCWCs (87%) and private facilities (89%) offer normal delivery services, the service is available in only one-third of the NGO facilities (34%) (**Table 3.2**).
- Child curative care is available in 100% of the DHs and UHCs, and in more than 90% of MCWCs, UHFWCs, USC/RDs, and CCs. Most NGOs (88%) and more than half of private facilities (57%) offer curative care for sick children (Table 3.2).
- Child vaccination service is mainly offered through the EPI program conducted mostly in public health facilities where it is provided universally. However, this service is minimally available at private hospitals (3%) (**Table 3.2**).



Figure 3.1 Availability of basic client services in public health facilities



3.2.3 General Service Preparedness

The 2022 BHFS collected information to assess the general preparedness of health facilities to offer quality health services. According to WHO (2013), an assessment of the general preparedness of a facility to provide quality services should consider the following six components:

- Basic amenities for client services
- Basic equipment to support quality health services
- Standard precautions for infection control in service delivery areas
- Capacity for adherence to standards for quality sterilization
- Diagnostic capacity
- Availability of essential medicine

During data collection, the 2022 BHFS interviewers verified that all components were present in the facility and in working order. The following sections present information from the 2022 BHFS on each of the six components for assessing general service preparedness.

Basic Amenities for Client Services

The components reviewed here are necessary but not sufficient to provide quality services. However, the availability of the following basic amenities is important to a client's satisfaction with health services rendered at a facility:

- Regular electricity
- Improved water source
- Visual and auditory privacy during consultations
- Client latrine
- Communication equipment (land-line/mobile phone)
- Computer with Internet access

In addition to these basic amenities, the 2022 BHFS considered the availability of emergency transport, another key component of client services. Emergency transport is expected to be available primarily in higher-level facilities; its availability also depends on the services provided by the facility. **Table 3.3** provides information on the availability of basic amenities for client services.

Electricity:

The 2022 BHFS obtained information on the connectivity of facilities with the national electricity grid/pollibiddut and the availability of regular electricity. Regular electricity is considered to be available at a facility if one of the following conditions is met: the facility is connected to a central power grid and the power supply was not interrupted for more than 2 hours at a time during normal working hours in the 7 days before the survey, the facility has a functioning generator on the day of the survey, or the facility has back-up solar power.

- Across public facilities excluding CCs, connection to the national electricity grid/polli-biddut is almost universal (97%) (Table 3.3).
- Almost all urban facilities (95%) and most rural facilities (85%) are connected to the national electricity grid / polli-biddut (**Table 3.3**).
- In CCs, connection to the national electricity grid/polli-biddut has increased from 47% in 2017 to 81% in 2022 and in UHFWCs from 83% in 2017 to 97% in 2022 (Figure 3.2).
- Overall, 85% of all public facilities are connected to national electricity grid/polli-biddut, as compared to 57% in 2017 (**Figure 3.2**).



Figure 3.2 Availability of electricity from national grid/polli-biddut, by facility type

• Around half (48%) of all public facilities have regular electricity, as compared to 40% of public facilities in 2017 (**Table 3.3** and **Figure 3.3**).

- Almost all of district and upazila level public facilities (90%), 53% of union-level public facilities, 45% of community clinics, 77% of NGO facilities, and 97% of private hospitals have regular electricity (Table 3.3 and Figure 3.3).
- Electricity is available regularly in almost all urban facilities (92%), while nearly half (48%) of rural facilities have consistent regular electricity (**Table 3.3**).
- Since 2017, the availability of regular electricity has increased across all public health facilities (Figure 3.3).



Figure 3.3 Availability of regular electricity*, by facility type

*regular electricity is considered to be available if facility is considered to a central power grid and there has not been an interruption in power supply lasting for more than 2 hours at a time during normal working hours in the 7 days before the survey, or the facility has a functioning generator, or else facility has back up solar power

Improved water source

- Most public facilities (81%) have an improved water source (**Table 3.3**).
- By facility type, the proportion with an improved water source varies from 77% in CCs to 100% in private hospitals (**Table 3.3**).
- Rural facilities (81%) exhibit lower improved water source, compared to urban facilities (97%). Almost all facilities in Rajshahi (96%) have an improved water source which is higher than other divisions (Table 3.3).

Privacy during consultations

Visual and auditory privacy are important in consultations with health care providers because this allows clients to express their problems without hesitation and in detail.

- Around 40% of public facilities have the capacity to ensure privacy for clients during consultations (**Table 3.3**).
- Most private hospitals (82%) are able to provide visual and auditory privacy (**Table 3.3**).

Client latrine

- Three-quarters of the public facilities (75%) have a functioning client latrine (**Table 3.3**).
- In DHs, MCWCs, UHCs, and private hospitals, the availability of a functional latrine for general outpatient client use is universal (≥90%) (Figure 3.4).
- The client latrine is available in 88% UHFWCs, 76% in USC/RDs, and 71% in CCs (Figure 3.4).
- Across the total public facilities excluding CCs, the availability of client latrine has increased from 82% in 2017 to 86% in 2022 (**Figure 3.4**).
- Around one-quarter of public facilities excluding CCs have a separate latrine for women (29%), whereas the availability of a separate toilet for women is very low in CCs (8%) (**Table 3.3**).



Figure 3.4 Availability of client latrine, by facility type

Communication equipment (land-line/mobile phone)

- Almost all DHs (98%), UHCs (90%), most private facilities (86%), and majority of NGO facilities (75%) have communication equipment– a functioning official land phone/mobile phone or private mobile phone reimbursed by the facility. However, communication equipment is less available in CCs (6%), UHFWCs (8%), and USC/RDs (9%) (Table 3.3 and Figure 3.5).
- There has been improvement in the availability of communication equipment across the public facilities between 2017 and 2022 (Figure 3.5).

• The availability of communication equipment in NGO facilities has increased from 71% in 2017 to 75% in 2022, whereas it has declined in private hospitals from 97% to 86% (**Figure 3.5**).



Figure 3.5 Availability of communication equipment (land-line/mobile phone), by facility type

Computer with Internet

- DHs (100%) and UHCs (99%) have universal access to a functioning computer/laptop/tablet with internet access (**Table 3.3** and **Figure 3.6**).
- The availability of a functioning computer/laptop/tablet with internet access is 77% in MCWCs and 32% in UHFWCs (**Figure 3.6**).
- Although 86% of the CCs have a computer/laptop/tablet, only 44% of CCs have a functioning device with internet access (**Table 3.3**).
- The availability of functioning computers with internet access has increased in MCWCs and UHFWCs between 2017 and 2022 (**Figure 3.6**).



Figure 3.6 Availability of a functioning computer with Internet, by facility type

Emergency transport

- Most of district and upazila level public facilities (84%) have functional ambulance or vehicle for emergencies (Table 3.3).
- Almost all DHs (97%) and most of the UHCs (89%) have an ambulance or other vehicle for emergency transportation of clients (**Table 3.3** and **Figure 3.7**).
- Majority of the MCWCs (56%) have a functional ambulance or vehicle (**Table 3.3** and **Figure 3.7**).
- Only one-third (32%) of private hospitals have a functional ambulance or vehicle which has decreased from 2017 (65%) (**Table 3.3** and **Figure 3.7**).



Figure 3.7 Availability of emergency transport, by facility type

■2014 BHFS □2017 BHFS ■2022 BHFS

In summary, the ideal facility should have a safe and welcoming environment with regular electricity, an improved water source, privacy during consultations, a client latrine, a landline/mobile phone, a computer with internet access and emergency transport. **Figure 3.8** presents the average score of amenities (out of 6 amenities) in health facilities in Bangladesh, by facility type.

- The average score for six amenities of total public facilities is 3.0 (out of 6 amenities) which has reduced from 2017 BHFS (3.4) (**Table 3.3** and **Figure 3.8**).
- DHs and private hospitals have the highest average score of amenities (5.2), whereas USC/RDs have the lowest average score (2.7) (**Table 3.3** and **Figure 3.8**).



Figure 3.8 Average score of basic amenities (out of 6 amenities), by facility type

Basic Equipment to Support Quality Health Services

Average

Delivery of quality basic health services requires certain equipment. WHO and USAID have proposed a list of seven basic pieces of equipment that should be available at a health facility to guarantee its readiness to deliver basic health services (WHO 2012). The basic equipment considered necessary to support quality health services includes adult scale, child or infant scale, thermometer, stethoscope, blood pressure apparatus and light source. **Table 3.4** provides information on the availability of this basic equipment from the 2022 BHFS.

- Most public facilities have stethoscope (89%), adult weighing scale (83%), blood pressure apparatus (81%), and thermometer (74%) (Table 3.4).
- Nearly two-thirds of public facilities (63%) have an infant or child weighing scale and about threequarters (76%) have a light source (**Table 3.4**).

Basic Items for Infection Control in Service Delivery Areas

Around the world, infections acquired in a health facility (known as nosocomial infections) often complicate the delivery of health care services. Strict adherence to infection control guidelines and constant vigilance are necessary to prevent such infections. It is essential that a health facility have the supplies and equipment for infection control appropriate to the services offered. **Table 3.5** provides information on 18 items considered important for compliance with standard precautions for infection control in service delivery areas.

- Only 9% of public facilities (36% excluding CCs) have basic sterilization equipment for processing instruments for reuse. Such equipment includes: Functioning electric dry heat sterilizer, functioning electric autoclave, non-electric autoclave with a functioning heat source (**Table 3.5**).
- Sterilization equipment is universally available in DHs (98%), in most of the UHCs (85%), and majority of NGO facilities (70%). CCs generally do not have provision for sterilization equipment, so it is expected that none of these facilities reported having such equipment (**Table 3.5**).

- More than half public facilities (55%) can safely dispose of sharps waste by incinerating or burning it, dumping it in a protected area, or storing it in a protected container for later removal. Majority of public facilities (60%) have containers for storage of sharps waste prior to disposal (sharps boxes). Syringes and needles for safely managing injections are available in nearly 54% of public facilities (**Table 3.5**).
- Disinfectants are available in half of the public facilities (54%). However, disinfectants are available in 73% of DHs, 68% of MCWCs, and 64% of UHCs (**Table 3.5**).
- Overall, 76% of public facilities (78% excluding CCs) have soap and running water or else alcoholbased hand disinfectant. Almost all the NGO clinics/hospitals (97%) have soap and running water or alcohol-based hand disinfectant (**Table 3.5**).
- Medical masks are available in most of public facilities (81%). Availability of medical masks is highest in CCs (85%) and lowest in USC/RD (66%) (Table 3.5).
- The average score of availability of all infection control items in public facilities is 7.4 (8.0 public facilities excluding CCs) out of 16, which has increased from 6.7 in 2017. The average availability of all infection control items is highest in NGO clinics/ hospital (11.2) and lowest in CCs (7.1) (**Table 3.5** and **Figure 3.9**).

Figure 3.9 Average score of availability of all infection control items (out of 16), by facility type



Average

Capacity for Adherence to Standards for Quality Sterilization or High-level Disinfection

For most equipment used in client examinations, either sterilization or high-level disinfection (HLD) procedures are sufficient to prevent the spread of infection. However, to effectively kill the spores that cause illnesses such as tetanus, either dry heat sterilization or an autoclave system is required. This type of treatment is necessary for processing surgical equipment that will be reused, such as blade handles and scissors used to cut umbilical cords. The standard equipment necessary to process these types of instruments for reuse includes:

- Functioning electric dry heat sterilizer
- Functioning electric autoclave
- Non-electric autoclave with a functioning heat source
- Electric boiler or steamer
- Non-electric boiler or steamer with a functioning heat source

Depending on the size of the facility, equipment may be processed through different methods or at more than one site within the facility. The information presented in this chapter refers to the primary site in the facility where equipment is processed.

Table 3.6 reports on the capacity of health facilities to process instruments for reuse, including the availability of sterilization equipment and an automatic timer and knowledge of the correct processing time.

- About 59% of public facilities excluding CCs have the functioning equipment or items necessary to perform an accepted method of equipment processing (**Table 3.6**).
- Almost half of public facilities excluding CCs (47%) have both functioning equipment and correct knowledge of the processing time (**Table 3.6**).
- One-fourth (24%) of all public facilities excluding CCs have the necessary equipment, knowledge of processing time, and an automatic timer (**Table 3.6**).
- Written guidelines for sterilization or HLD are available in 12% of total public facilities excluding CCs (**Table 3.6**).

Diagnostic Capacity

Provision of diagnostic services, including laboratory tests and diagnostic imaging, is essential for clinical decision making and for enhancing delivery of quality health care. In fact, case management for conditions such as TB depends entirely on laboratory and/or imaging results. **Tables 3.7.1** and **3.7.2** and **Figures 3.10** present information on the availability and capacity of laboratory diagnostic testing of health facilities in Bangladesh. The items included in these tables and figures are based on the methodology proposed by WHO and USAID for measuring diagnostic capacity as a component of assessing general service preparedness (WHO 2012).

- Overall, the availability of basic laboratory diagnostic tests in public health facilities except blood glucose test is still limited. Majority of the public facilities (57%) offer blood glucose test (**Table 3.7.1**).
- Although blood glucose test is universally offered by UHCs (98%) (Table 3.7.1), less than eighty percent of UHCs (79%) have functioning necessary equipment to conduct blood glucose test (Table 3.7.2).
- Urine protein test is available in most of DHs (89%), UHCs (85%), and NGOs (86%), however, only 4% of CCs have this test available (Table 3.7.1).

Figure 3.10 presents average score on the diagnostic testing capacity (hemoglobin, blood glucose, urine protein, urine glucose, urine pregnancy test), by facility type.

• The average diagnostic testing capacity among public facilities excluding CCs is 1.8 (out of 5.0). The availability has gradually increased over the years except DHs, CCs, and private hospitals (**Table 3.7.2** and **Figure 3.10**).
• The average capacity to conduct basic diagnostic tests (out of five) is highest in NGO clinics/ hospitals (4.4), followed by UHCs (4.3) (**Table 3.7.2** and **Figure 3.10**).



Figure 3.10 Average capacity to conduct basic diagnostic tests (out of five), by facility type

Figure 3.11 and **Table 3.7.1**, shows the availability and **Table 3.7.2** provision of a functional X-ray machine in district and upazila facilities, NGO facilities, and private hospitals.

- Almost all DHs (97%) and majority of UHCs (59%) have an X-ray machine (Table 3.7.1). But 87% DHs and only 47% of UHCs have a functioning X-ray machine (Table 3.7.2 and Figure 3.11).
- Functioning X-ray machines have increased in DHs and UHCs between 2017 and 2022 (Figure 3.11).
- More than half of private hospitals (54%) have a functioning X-ray machine, but only a very negligible proportion of NGO facilities (<1%) have a functioning X-ray machine (Table 3.7.2 and Figure 3.11).

Average

Figure 3.11 Functioning x-ray machine in health facilities, by facility type Percent



■2014 BHFS ■2017 BHFS ■2022 BHFS

The BHFS 2022 also obtained information on the availability and provision of functional ultrasound machines.

- Almost all DHs (90%) and more than half (58%) of UHCs have an ultrasonography machine (Table 3.7.1). But only four-fifths of DHs (79%) and nearly half (49%) of UHCs have a functioning ultrasonography machine (Table 3.7.2 and Figure 3.12).
- Functioning ultrasonography machine slightly increased in DHs and substantially (three-folds) increased in UHCs between 2017 and 2022 (**Table 3.7.2** and **Figure 3.12**).
- Three-quarters of private hospitals (75%) and 42% of NGO facilities have functioning ultrasound machines (Table 3.7.2 and Figure 3.12).



Figure 3.12 Functioning ultrasound machines in health facilities, by facility type

■2014 BHFS ■2017 BHFS ■2022 BHFS

Availability of Essential Medicines

Percent

Consistent availability of essential medicines is critical to the delivery of quality health services. The 2022 BHFS assessed the availability of 14 essential medicines in concurrence with the service readiness indicators proposed by WHO and USAID (WHO 2012). **Table 3.8.1** and **3.8.2** present information on the availability of these medicines at health facilities.

- Across all types of facilities, ciprofloxacin tablets/capsules are the most widely available essential medicine followed by omeprazole/cimetidine tablets/capsules. Almost all of public facilities had ciprofloxacin available on the day of the survey. Public facilities (97%) and NGO facilities (91%) are more likely than private hospitals (59%) to have ciprofloxacin tablets/capsules (Table 3.8.1).
- Amitriptyline tablets, atenolol tablets, captopril tablets, ceftriaxone injectable, glibenclamide tablets, and simvastatin tablets are the least available, in 10% or less of public facilities. The limited availability of these medicines reflects the fact that they are generally provided only at higher-level health facilities and are not expected to be available at lower-level facilities (**Table 3.8.1**).
- Examining differences by division, ciprofloxacin tablets/capsules, omeprazole/cimetidine tablets/capsules, paracetamol oral suspension, and salbutamol inhaler are the most widely available medicines (81% or more in all divisions) (**Table 3.8.2**).

Availability of medicines supplies in community clinics

The information obtained in the BHFS can be used to assess the availability of nineteen essential medicines supplied in CCs. CCs receive drugs at different intervals from the central warehouse, with the following items including albendazole tablet, amoxicillin capsule, syrup & pediatrics drop, antacid tablet, benzoic and salicylic acid ointment, calcium lactate tablet, chlorpheniramine tablet & syrup, co-trimoxazole tablet (120 & 960 mg), doxycycline capsule, erythromycin tablet & syrup, ferrous fumarate & folic acid tablet, gentian violet, hyoscine-

n-butyl bromide tablet, metronidazole tablet, neomycin and bacitracin skin ointment, paracetamol tablet & syrup, penicillin tablet, salbutamol syrup, vitamin B complex tablet, zinc dispersible tablet.

- The availability of doxycycline capsules is universal (100%) followed by paracetamol tablets and syrup (98%) and metronidazole tablets (97%) (**Table 3.8.3**).
- Erythromycin tablets and syrup are almost unavailable throughout all the CCs (**Table 3.8.3**).

Availability of medicines in drug and dietary supply (DDS) kit

As another measure of the availability of drugs, the information obtained in the BHFS can be used to assess the availability of twenty medicines included in the drug and dietary supply (DDS) kit. A DDS kit is a box of preselected essential drugs and dietary items to be dispensed by service providers at MCWCs and UHFWCs for patient management. They receive DDS kits at different intervals from the central warehouse, with the following items including iron and folic acid tablet, folic acid tablet, calcium carbonate tablet, vitamin B complex tablet, albendazole tablet, paracetamol tablet, antacid tablet, drotaverine hydrochloride tablet, pantoprazole tablet, metronidazole tablet, chlorpheniramine maleate tablet, salbutamol tablet, diazepam tablet, cotrimoxazole (120 mg) dispersible, cotrimoxazole DS (960 mg), amoxycillin (500 mg), doxycycline (100 mg) capsule, suspension amoxycillin, cotrimoxazole suspension, paracetamol suspension, and benzoic acid and salicylic acid (Packaging list of DDS kit, DGFP 2022-2023).

- Almost all UHFWCs and MCWCs (more than 90%) have iron and folic acid tablet, vitamin B complex tablet, albendazole tablet, paracetamol tablet, pantoprazole tablet, and metronidazole tablet (Table 3.8.4).
- The least available is calcium carbonate tablet (available in 15% of UHFWCs and 18% of MCWCs) (**Table 3.8.4**).

Availability of medicines and equipment of delivery kit

The information obtained in the BHFS can be used to assess the availability of eleven medicines included in the delivery kit. A delivery kit is a box of pre-selected essential drugs and dietary items to be dispensed by service providers at MCWCs and UHFWCs for patient management. They receive delivery kits at different intervals from the central warehouse, with the following items including paracetamol, amoxicillin capsule, infusion 5% dextrose in aqua (500 ml), infusion 5% dextrose in normal saline (500 ml), chlorhexidine cream (60 ml bottle), sterile surgical gloves, cotton ball thread, sterile gauge pack, surgical adhesive tape, 1% lidocaine injection (50 ml), disposable syringe (5 ml).

- The availability of paracetamol, sterile surgical gloves and disposable syringe (5 ml) is universal throughout the MCWCs and UHFWCs, followed by amoxicillin capsule (available in 91% of UHFWCs and 89% of MCWCs) (**Table 3.8.5**).
- However, the availability of surgical adhesive tape is the lowest in UHFWCs (5%). For MCWCs, availability of 1% lidocaine injection (50 ml) is the lowest (50%) (**Table 3.8.5**).

3.3 BASIC MANAGEMENT SYSTEMS TO SUPPORT AND MAINTAIN QUALITY SERVICES, AND APPROPRIATE CLIENT UTILIZATION

Establishing Basic management and administrative systems are essential for the consistent delivery of health care services at an acceptable level of quality. The 2022 BHFS elicited information on:

- Supportive management practices for providers
- Quality assurance

3.3.1 Supportive Management Practices for Providers

Health facilities are considered to have supportive management practices if they had external supervision during the 6 months before the survey and if staff members received routine training and personal supervision. **Table 3.9** presents information on supportive management practices.

External Supervision

Supervision by external managers has many benefits. It can ensure that system-wide standards and protocols are followed at the facility level and promote an organizational culture that expects standards and protocols to be followed. External supervision provides an opportunity to expose staff members to a wider scope of ideas and relevant experiences. It can also motivate service providers, especially if the supervisor is supportive.

- Most public facilities (89%) have reported receiving external supervision in the 6 months preceding the BHFS (Table 3.9).
- Recent external supervision is common across all types of facilities, ranging from 84% among unionlevel public facilities to 100% in DHs (**Table 3.9**).

Staff Training

Staff training is essential for updating health workers with knowledge, skills, and technical competence to improve the quality of health care services. The 2022 BHFS assessed whether healthcare providers had received any formal or structured in-service training related to the services they offer in the 24 months preceding the survey. If more than half of the providers at a facility had received such training, the facility was deemed to have routine staff training.

- Overall, 72% of public facilities have routine staff training (Table 3.9).
- Routine staff training is less common in private hospitals (16%), compared to other type of facilities (**Table 3.9**).
- Routine training is much more common in rural facilities (73%) than in urban facilities (26%) (Table 3.9).
- The proportion of facilities with routine staff training varies considerably by division, from 49% in Khulna to 77% in Mymensingh (**Table 3.9**).

Personal Supervision of Health Service Providers

While facility-level supervision is critical to support quality service provision throughout the health care system, personal supervision is essential to assess the work of individual staff members. Personal supervision identifies each employee's strength and weakness and provides appropriate support.

- The 2022 BHFS defined a health facility as having personal supervision of health care providers if at least half of the providers reported being personally supervised at least once during the 6 months preceding the survey.
- Almost all public facilities (93%) have at least half of the providers reported receiving personal supervision during the last six months before the survey. About 87% of private facilities and 99% of NGOs have the same personal supervision (Table 3.9).

Staff Training and Personal Supervision

The combination of staff training and personal supervision is crucial to achieving competence and sustaining quality health service delivery.

- Majority of public facilities (68%) have both routine staff-training and personal supervision of health workers (Table 3.9).
- CCs have the highest percentage (73%) with respect to routine training and personal supervision (Table 3.9).
- By division, the percentage of facilities with both training and supervision varies from 47% in Khulna to 72% in Sylhet (**Table 3.9**).

Supportive Management Practices

A facility is considered to have supportive management practices if the facility had an external supervisory visit during the 6 months before the assessment and staff had received both routine training and supervision.

- Majority of public facilities (65%) have supportive management practices (Table 3.9).
- Around six in ten NGOs (59%) and only 15% private facilities have supportive management practices (**Table 3.9**).

3.3.2 Quality Assurance

Quality assurance (QA) refers to a system for monitoring quality of care, identifying problems, and instituting changes to resolve those problems. QA systems require an established standard against which quality is measured. There must be systematic methods to assess results and develop interventions. The following are examples of QA activities and approaches:

- I. A supervisory checklist for health systems, which documents the presence of equipment and supplies, the completeness of health management information system accounts, and other process indicators.
- II. A supervisory checklist for health service provision, which verifies specific content in client assessments, treatments, or consultations. This list is often used to document the provision of care.

- III. A facility-wide review of mortality, which is a structured system to review the records of each client who dies. There is usually a committee established for this purpose.
- IV. Audits of medical records or registers, which check medical records for specific items or information and assess adherence to protocols.
- Nearly one in five (18%) public facilities report that they have regular QA activities and have evidence of such activities, an increase from 2017 (13%) (Table 3.10 and Figure 3.13).
- Supportive documentation of QA activities is observed highest in DHs (66%), followed by UHCs (61%), NGOs (48%) and MCWCs (22%) (Table 3.10 and Figure 3.13).
- Only 16% both in union-level facilities and private facilities have documented QA activities (Table 3.10).



Figure 3.13 Quality assurance, by facility type

Percent

3.4 AVAILABILITY OF HUMAN RESOURCES FOR HEALTH

The WHO considers health workforce as one of the key building blocks in any healthcare system. As part of its continuing commitment to improve the provision of health services in the public sector, the Government of Bangladesh is committed to expanding the human resources for health.

As part of the assessment of general preparedness for service delivery, the BHFS collected information on staffing patterns. **Table 3.11.1** shows the percentages of sanctioned healthcare provider posts filled in the surveyed facilities, by provider category and type of facility.

Senior/junior consultant (anesthesia) are available in almost all DHs (97%) and two-thirds of UHCs (65%) (Table 3.11.1).

- Senior/ junior consultant (medicine) are available in 87% of DHs and 32% of UHCs (Table 3.11.1).
- Almost eight in ten (79%) DHs have senior/junior consultant (Surgery) assigned in contrast to only about one-fifth of UHCs (17%) (**Table 3.11.1**).
- Radiologist are assigned in 44 % of DHs as compared to 5% of UHCs (Table 3.11.1).
- Medical Technologist (lab) are available universally (97%) in DHs and four-fifths of UHCs (83%) (Table 3.11.1).
- On average, 21 medical officers are working in a DH as compared with 10 in UHC. Number of medical officer ranges from 4 to 52 among the DHs and from 1 to 22 among the UHCs (**Table 3.11.2**).
- In a DH, 126 nurses are available on an average, as compared with 25 in a UHC. The number of nurse ranges from 5 to 398 among the DHs and from 1 to 37 among the UHCs (**Table 3.11.2**).

3.5 MANAGEMENT OF MEDICAL WASTE IN HEALTH CARE FACILITIES

WHO considers poor medical waste management to be one of the key elements posing potential risks of infection, toxic effects, and injuries among health care workers, waste handlers, patients, and the community at large. As part of its continuing effort to control infections in public sector health care facilities, the Government of Bangladesh is committed to ensuring proper management of medical waste management. It is therefore essential that all medical waste materials be disposed of safely.

As part of the assessment of general preparedness for service delivery, the BHFS collected information on management of medical waste. **Tables 3.12.1-3.12.2** and **Figure 3.14-3.15** present information on final disposal of medical waste at the surveyed facilities.

Sharps waste: According to WHO (2001), sharps waste includes needles, intravenous tubing with a needle attached, glass Pasteur pipettes, disposable glass pipettes, scalpels, razor blades, lancets, and broken glass that may be contaminated with blood or other potentially infectious material.

- Only, less than 1% of public facilities use an incinerator to burn their sharps waste (**Table 3.12.1**).
- 11% of public facilities practice open burning (19% excluding CCs), 8% dump waste without burning it (10% excluding CCs), and 36% remove waste offsite (21% excluding CCs) (Table 3.12.1).
- The most widely practiced method of managing sharps waste is removal offsite followed by open burning (Table 3.12.1 and Figure 3.14).



Figure 3.14 Sharps waste management, by facility type

Non-sharps medical waste: According to WHO (2001), non-sharps medical waste includes infectious waste (pathogens), pathological waste (human tissues, body parts, or fluids), pharmaceutical waste, genotoxic waste, chemical waste, and radioactive waste.

- Overall, 13% of public facilities engage in open burning to manage non-sharps waste (Table 3.12.2).
- Less than 1% (1% excluding CCs) of public facilities burn their medical waste using an incinerator.
- One-third of DHs (31%), MCWCs (37%), and one-fifth of CCs (21%) practice the remove offsite method, whereas the majority of NGOs (60%) and private facilities (71%) use the remove offsite method (Table 3.12.2 and Figure 3.15).

Percent



Figure 3.15 Non-sharps waste management, by facility type

Percent

FACILITY-LEVEL INFRASTRUCTURE, RESOURCES, MANAGEMENT, AND SUPPORT

LIST OF TABLES

- Table 3.1 Availability of specific services
- Table 3.2 Availability of basic client services
- Table 3.3 Availability of basic amenities for client services
- Table 3.4 Availability of basic equipment
- Table 3.5 Standard precautions for infection control
- Table 3.6 Capacity for processing equipment for reuse
- Table 3.7.1 Availability of laboratory diagnostic testing by facility type
- Table 3.7.2 Laboratory diagnostic capacity by facility type
- Table 3.7.3 Laboratory diagnostic capacity by location and division
- Table 3.8.1 Availability of essential medicines by facility type
- Table 3.8.2 Availability of essential medicines by location and division
- Table 3.8.3 Availability of medicines in community clinics
- Table 3.8.4 Availability of DDS kit medicines
- Table 3.8.5 Availability of medicines and equipment of delivery kit
- Table 3.9 Supportive management practices at the facility level
- Table 3.10 Quality assurance
- Table 3.11.1 Staffing pattern in surveyed facilities
- Table 3.11.2 Staffing pattern in surveyed facilities (medical officer and nurse)
- Table 3.12.1 Sharps waste management
- Table 3.12.2 Non-sharps waste management
- Table 3.13 Key indicators for general service preparedness in MCWCs

Table 3.1: Availability of specific services

Among all facilities, the percentages and numbers that offer the indicated specific services, Bangladesh HFS 2022

| | | Percentage of total public | | | Number of facilitie | es offering service |
|--|---|--|--|-----------------------------------|---------------------|---------------------|
| Service provided | Percentage of total public facilities offering service | facilities excluding CCs offering service | Percentage of NGO and private facilities offering service | Percentage of total facilities | Weighted | Unweighted |
| | | | | g | | |
| Curative care for sick children | 94.8 | 95.7 | 61.0 | 90.2 | 1,404 | 1,406 |
| Child growth monitoring | 88.2 | 76.4 | 30.6 | 80.4 | 1,251 | 1,196 |
| Child vaccination (EPI) ¹ | 93.1 | 80.5 | 12.4 | 82.1 | 1,278 | 1,178 |
| Any family planning ² | 82.9 | 85.1 | 31.8 | 75.9 | 1,182 | 1,197 |
| Antenatal care | 99.5 | 98.5 | 76.2 | 96.3 | 1,499 | 1,500 |
| Normal delivery | 17.8 | 55.0 | 81.5 | 26.5 | 412 | 740 |
| Cesarean delivery ³ | 1.7 | 6.6 | 82.7 | 12.7 | 199 | 378 |
| TB diagnosis or treatment ⁴ | 23.8 | 22.2 | 20.6 | 23.4 | 364 | 455 |
| Non-communicable disease ⁵ | 22.9 | 29.8 | 65.1 | 28.6 | 446 | 654 |
| Laboratory diagnostic ⁶ | 23.3 | 35.8 | 85.1 | 31.7 | 493 | 735 |
| Blood grouping and typing ⁷ | 2.9 | 9.4 | 81.4 | 13.6 | 212 | 487 |
| Blood transfusion ⁸ | 2.2 | 8.4 | 67.6 | 11.1 | 173 | 371 |
| Postnatal care9 | 97.3 | 97.1 | 78.9 | 94.8 | 1.476 | 1.482 |
| Postpartum family planning ¹⁰ | 89.7 | 90.5 | 52.9 | 84.7 | 1,318 | 1,329 |
| Adolescent health ¹¹ | 90.8 | 89.8 | 37.8 | 83.5 | 1.301 | 1.281 |
| Nutrition ¹² | 95.2 | 91.5 | 44.3 | 88.3 | 1,375 | 1,335 |
| Total | - | - | - | - | 1,557 | 1,557 |

¹ Routine series of DPT/pentavalent, polio, and measles vaccinations offered from the facility, excluding any outreach services

² Facility provides, prescribes, or counsels clients on any of the following: contraceptive pills (combined or progestin only), injectables (progestin only), implants, intrauterine contraceptive devices (IUCDs), male condoms, female sterilization (tubal ligation), male sterilization (vasectomy), or lactational amenorrhea method (LAM).

³ Facility reports that it provides cesarean delivery services at the facility.

⁴ Facility reports that providers assigned to the facility diagnose TB, prescribe treatment for TB, or provide TB treatment follow-up services for clients put on treatment elsewhere. ⁵ Facility reports that it provides diagnosis or management of non-communicable diseases, especially diabetes and cardiovascular diseases.

⁶ Facility reports that it provides laboratory diagnostic services at the facility.
 ⁷ Facility reports that it provides blood grouping and typing services at the facility.

⁸ Facility reports that it provides blood transfusion services at the facility.

⁹ Facility reports that it provides postnatal care in facility.

¹⁰ Facility reports that it provides postpartum family planning services at the facility.

¹¹ Facility reports that it provides adolescent health care at the facility. ¹² Facility reports that it provides nutrition services at the facility.

Table 3.2: Availability of basic client services

Among all facilities, the percentages offering the indicated basic client services and all basic client services, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Child curative care | Child growth monitoring services | Child vaccination services | Any modern method of family planning | Antenatal care services | Normal delivery | All basic client services ¹ with normal delivery | All basic client services without normal delivery | Number of facilities (weighted) | Number of facilities (unweighted) |
|------------------------------|---------------------------|---|----------------------------------|--|-------------------------------|--------------------|--|--|---------------------------------------|---|
| Facility type | | | | | | , | | | | (g) |
| District and upazila | | | | | | | | | | |
| public facilities | 99.1 | 92.9 | 90.9 | 97.1 | 99.8 | 97.4 | 85.1 | 85.8 | 41 | 319 |
| DH | 100.0 | 98.4 | 95.2 | 93.5 | 100.0 | 100.0 | 88.7 | 88.7 | 4 | 62 |
| MCWC | 95.1 | 68.1 | 57.0 | 95.1 | 99.0 | 87.1 | 40.1 | 42.0 | 7 | 100 |
| UHC | 100.0 | 98.1 | 98.5 | 98.2 | 100.0 | 99.5 | 95.5 | 96.0 | 30 | 157 |
| Union-level public | | | | | | | | | | |
| facilities | 95.3 | 74.2 | 79.1 | 83.5 | 98.3 | 49.4 | 32.6 | 52.3 | 310 | 434 |
| UHFWC | 94.2 | 78.2 | 79.2 | 96.0 | 100.0 | 59.2 | 39.2 | 59.3 | 225 | 293 |
| USC/RD | 98.1 | 63.6 | 79.1 | 50.3 | 93.7 | 23.4 | 15.2 | 33.7 | 85 | 141 |
| Community clinic (CC) | 94.5 | 92.4 | 97.5 | 82.1 | 99.8 | 4.7 | 3.6 | 70.4 | 994 | 488 |
| NGO clinic/hospital | 88.4 | 81.9 | 71.9 | 74.8 | 100.0 | 33.5 | 16.6 | 50.5 | 29 | 127 |
| Private hospital | 56.6 | 22.5 | 3.0 | 25.0 | 72.4 | 89.0 | 1.1 | 1.1 | 183 | 189 |
| Location | | | | | | | | | | |
| Urban | 65.7 | 36.9 | 22.4 | 38.6 | 79.3 | 81.7 | 11.2 | 16.2 | 235 | 514 |
| Rural | 94.6 | 88.1 | 92.7 | 82.6 | 99.3 | 16.6 | 11.6 | 66.2 | 1,322 | 1,043 |
| Division | | | | | | | | | | |
| Barishal | 92.7 | 87.6 | 84.5 | 82.0 | 98.8 | 19.2 | 11.1 | 63.6 | 111 | 152 |
| Chattogram | 97.4 | 87.5 | 85.1 | 80.9 | 97.4 | 30.6 | 16.0 | 71.5 | 291 | 244 |
| Dhaka | 83.8 | 75.7 | 80.0 | 70.7 | 95.4 | 30.1 | 11.2 | 49.1 | 317 | 264 |
| Khulna | 76.7 | 76.2 | 70.9 | 70.3 | 90.5 | 30.4 | 8.8 | 50.1 | 210 | 208 |
| Rajshahi | 95.1 | 73.9 | 82.5 | 75.4 | 97.8 | 26.4 | 10.2 | 55.4 | 214 | 212 |
| Rangpur | 95.7 | 84.7 | 87.3 | 79.3 | 97.7 | 17.1 | 8.0 | 64.5 | 193 | 190 |
| Sylhet | 88.4 | 82.7 | 84.0 | 74.4 | 98.0 | 31.6 | 20.4 | 59.2 | 99 | 136 |
| Mymensingh | 94.7 | 78.8 | 87.2 | 78.6 | 97.5 | 18.2 | 7.1 | 59.3 | 122 | 151 |
| Total | 90.2 | 80.4 | 82.1 | 75.9 | 96.3 | 26.5 | 11.5 | 58.7 | 1,557 | 1,557 |
| Total excluding CCs | 82.7 | 59.1 | 54.9 | 65.0 | 90.1 | 65.0 | 25.4 | 38.0 | 563 | 1,069 |
| Total public | 94.8 | 88.2 | 93.1 | 82.9 | 99.5 | 17.8 | 12.8 | 66.7 | 1,345 | 1,241 |
| Total public excluding CCs | 95.7 | 76.4 | 80.5 | 85.1 | 98.5 | 55.0 | 38.8 | 56.2 | 351 | 753 |

¹ Basic client services include outpatient curative care for sick children, child growth monitoring, facility-based child vaccination services, any modern methods of family planning, antenatal care, and normal delivery.

Table 3.3: Availability of basic amenities for client services

Among all facilities, the percentages with indicated amenities considered basic for quality services, by background characteristics, Bangladesh HFS 2022

| | | | | | | Am | enities | | | | | | |
|-------------------------------|---------------------------------|-------------------------------------|--|---|--------------------------------|---|----------|---|-------------------------------------|---|---|---------------------------------------|---|
| Background characteristic | National electricity grid | Regular electricity ¹ | Improved water source ² | Visual and auditory privacy ³ | Client latrine ⁴ | Communic ation equipment ⁵ | Computer | Computer with Internet ⁶ | Emergency transport ⁷ | Separate latrine for female clients | Average basic amenities available (out of 6) ⁸ | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type | | | | | | | | | | | | | |
| District and upazila public | | | | | | | | | | | | | |
| facilities | 97.5 | 89.5 | 96.2 | 45.6 | 93.9 | 86.9 | 96.3 | 95.2 | 84.3 | 59.1 | 5.1 | 41 | 319 |
| | 100.0 | 98.4 | 98.4 | 38.7 | 90.3 | 98.4 | 100.0 | 100.0 | 96.8 | 67.7 | 5.2 | 4 | 62 |
| | 95.1 | 80.2 | 99.0 05.2 | 0.00 77 0 | 95.Z 0/ 1 | 00.8 90.1 | 100.0 | 77.0 | 20.3 | 47.4 60.6 | 4.0 | 30 | 100 |
| UNC | 91.1 | 09.2 | 95.2 | 44.0 | 94.1 | 50.1 | 100.0 | 90.9 | 09.5 | 00.0 | 5.1 | 50 | 157 |
| Union-level public | | | | | | | | | | | | | |
| facilities | 96.5 | 52.6 | 89.6 | 53.9 | 84.7 | 8.2 | 49.6 | 28.7 | 1.1 | 24.8 | 3.2 | 310 | 434 |
| UHFWC | 96.5 | 57.6 | 92.5 | 59.0 | 87.9 | 8.1 | 50.7 | 32.1 | 1.2 | 27.6 | 3.4 | 225 | 293 |
| USC/RD | 96.4 | 39.2 | 82.1 | 40.2 | 76.0 | 8.6 | 46.7 | 19.8 | 0.8 | 17.2 | 2.7 | 85 | 141 |
| Community clinic | | | | | | | | | | | | | |
| (CC) | 81.2 | 45.2 | 77.2 | 36.0 | 70.8 | 5.6 | 85.9 | 44.4 | 0.4 | 8.1 | 2.8 | 994 | 488 |
| NGO clinic/ | | | | | | | | | | | | | |
| hospital | 97.9 | 76.8 | 98.9 | 53.4 | 74.7 | 75.2 | 53.7 | 49.8 | 24.7 | 64.8 | 4.3 | 29 | 127 |
| Private hospital | 97.0 | 96.6 | 99.8 | 82.2 | 92.6 | 86.2 | 73.7 | 65.2 | 32.4 | 62.6 | 5.2 | 183 | 189 |
| Location | | | | | | | | | | | | | |
| Urban | 95.2 | 91.5 | 97 4 | 73.8 | 89.9 | 83.0 | 73 5 | 65.5 | 36.8 | 61.2 | 5.0 | 235 | 514 |
| Rural | 85.4 | 47.9 | 80.7 | 40.6 | 74.5 | 7.7 | 77.5 | 41.6 | 1.7 | 12.9 | 2.9 | 1,322 | 1,043 |
| Division | | | | | | | | | | | | | |
| Barishal | 83.2 | 44.2 | 76 1 | 48 5 | 68 7 | 13.7 | 76 1 | 53.8 | 39 | 14.2 | 3.1 | 111 | 152 |
| Chattogram | 75.4 | 60.4 | 74.5 | 45.1 | 75.1 | 19.6 | 79.5 | 48.6 | 7.0 | 20.3 | 3.1 | 291 | 244 |
| Dhaka | 89.0 | 52.3 | 82.4 | 48.7 | 75.2 | 24.2 | 70.7 | 43.3 | 11.2 | 25.0 | 3.3 | 317 | 264 |
| Khulna | 95.3 | 67.1 | 92.0 | 50.6 | 85.0 | 19.6 | 71.1 | 36.3 | 7.5 | 26.0 | 3.5 | 210 | 208 |
| Rajshahi | 98.0 | 54.3 | 95.8 | 43.6 | 83.9 | 18.9 | 76.8 | 53.6 | 6.6 | 19.1 | 3.5 | 214 | 212 |
| Rangpur | 85.3 | 55.0 | 82.1 | 39.6 | 69.4 | 13.6 | 88.1 | 41.2 | 3.8 | 13.2 | 3.0 | 193 | 190 |
| Sylhet | 87.9 | 41.3 | 91.4 | 68.5 | 86.0 | 17.5 | 83.6 | 52.0 | 7.3 | 25.5 | 3.6 | 99 | 136 |
| Mymensingh | 79.5 | 42.6 | 70.9 | 21.7 | 70.0 | 19.0 | 74.3 | 35.5 | 3.2 | 11.7 | 2.6 | 122 | 151 |
| Total | 86.9 | 55.4 | 83.3 | 45.6 | 76.8 | 19.1 | 76.9 | 45.2 | 7.0 | 20.2 | 3.2 | 1,557 | 1,557 |
| Total excluding CCs | 96.8 | 70.8 | 93.9 | 62.5 | 87.4 | 42.8 | 61.1 | 46.5 | 18.6 | 41.6 | 4.0 | 563 | 1,069 |
| Total public | 85.3 | 48.2 | 80.7 | 40.4 | 74.7 | 8.7 | 77.8 | 42.4 | 3.1 | 13.5 | 3.0 | 1,345 | 1,241 |
| Total public excluding CCs | 96.6 | 56.9 | 90.4 | 52.9 | 85.8 | 17.5 | 55.1 | 36.6 | 10.9 | 28.8 | 3.4 | 351 | 753 |

Note: The indicators presented in this table comprise the basic amenities domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012). ¹ Facility is connected to a central power grid and there was no interruption in power supply lasting for more than 2 hours at a time during normal working hours in the 7 days

before the survey, or facility has a functioning generator on the day of the survey, or else facility has back-up solar power.

² Water is piped into facility or piped onto facility grounds, bottled water is available, or water is available from a public tap or standpipe, a tube well or borehole, a protected dug well, a protected spring, or rainwater and the outlet from this source is within 500 meters of the facility.

³ A private room or screened-off space available in the general outpatient service area that is a sufficient distance from other clients so that a normal conversation could be held without the client being seen or heard by others

⁴ Facility has a functioning flush or pour-flush toilet, a ventilated improved pit latrine, or a composting toilet.
 ⁵ Facility has a functioning land-line telephone, a functioning facility-owned cellular phone, or a private cellular phone that is supported by the facility.
 ⁶ Facility has a functioning computer with access to the Internet, or facility has access to the Internet via a cellular phone inside the facility.

⁷ Facility has a functioning ambulance or other vehicle for emergency transport that is stationed at the facility and operates from the facility

⁸ Average number of basic amenities include regular electricity, an improved water source, visual and auditory privacy, a client latrine, communication equipment, and a computer with Internet.

Table 3.4: Availability of basic equipment

Among all facilities, the percentages with equipment considered basic to quality client services available in the general outpatient service area, by background characteristics, Bangladesh HFS 2022

| | | | | Equipment | | | | _ | |
|--|------------------------------|---|-----------------------------|-----------------------------|---|------------------------------|------------------------------|---------------------------------------|---|
| Background characteristic | Adult scale | Child scale ¹ or infant scale ² | Thermometer | Stethoscope | Blood pressure apparatus ³ | Light source ⁴ | All basic equipment | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type | | | | | | | | | |
| District and upazila public facilities DH MCWC | 91.9 88.7 88.0 | 86.7 82.3 76.3 | 88.6 80.6 80.8 | 98.4 98.4 97.1 | 96.0 90.3 97.1 | 92.1 91.9 91.1 | 72.2 59.7 60.2 | 41 4 7 | 319 62 100 |
| UHC | 93.3 | 89.9 | 91.7 | 98.8 | 96.5 | 92.3 | 77.0 | 30 | 157 |
| Union-level public facilities UHFWC USC/RD | 81.0 87.7 63.2 | 63.5 70.1 46.3 | 71.1 74.9 61.1 | 97.0 98.0 94.5 | 93.2 94.7 89.2 | 78.8 79.3 77.5 | 36.2 40.8 24.0 | 310 225 85 | 434 293 141 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 83.5 97.7 85.1 | 61.6 80.9 77.8 | 73.8 93.9 84.8 | 86.0 96.6 91.5 | 76.7 92.0 90.5 | 74.1 94.8 92.0 | 27.7 71.3 70.2 | 994 29 183 | 488 127 189 |
| Location | | | | | | | | | |
| Urban Rural | 87.0 83.1 | 77.7 62.7 | 86.2 73.4 | 93.8 88.6 | 92.2 80.7 | 93.0 75.4 | 68.9 30.6 | 235 1,322 | 514 1,043 |
| Division | 72.0 | F0 F | 77 4 | 04.5 | 74 5 | 02.2 | 28.0 | 444 | 450 |
| Chattogram Dhaka Khulna | 73.9 81.3 89.0 80.4 | 60.1 59.1 71.6 | 81.2 73.3 79.0 | 91.6 91.2 90.4 | 83.8 84.5 78.2 | 83.3 81.6 82.2 76.7 | 28.0 38.4 43.5 37.0 | 291 317 210 | 244 264 208 |
| Rajshahi Rangpur Sylhet | 89.8 83.1 86.3 | 71.1 69.0 78.7 | 79.9 67.8 78.7 | 87.1 87.8 87.3 | 83.4 80.2 90.1 | 70.6 67.5 81.7 | 34.7 26.6 50.3 | 214 193 99 | 212 190 136 |
| Mymensingh | 78.1 | 59.3 | 59.7 | 89.8 | 86.7 | 83.0 | 26.9 | 122 | 151 |
| Total | 83.7 | 64.9 | 75.3 | 89.3 | 82.4 | 78.0 | 36.4 | 1,557 | 1,557 |
| Total excluding CCs | 84.0 | 70.8 | 78.0 | 95.3 | 92.5 | 84.9 | 51.7 | 563 | 1,069 |
| Total public | 83.2 | 62.8 | 73.6 | 88.9 | 81.1 | 75.8 | 31.0 | 1,345 | 1,241 |
| Total public excluding CCs | 82.3 | 66.3 | 73.2 | 97.2 | 93.5 | 80.4 | 40.4 | 351 | 753 |

Note: The indicators presented in this table comprise the basic equipment domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012).

¹ A scale with a gradation of 250 grams, or a digital standing scale with a gradation of 250 grams or lower where an adult can hold a child to be weighed, available

somewhere in the general outpatient area ² A scale with a gradation of 100 grams, or a digital standing scale with a gradation of 100 grams where an adult can hold an infant to be weighed, available somewhere in the general outpatient area ³ A digital blood pressure machine or a manual sphygmomanometer with a stethoscope available somewhere in the general outpatient area

⁴ A spotlight source that can be used for client examination or a functioning flashlight available somewhere in the general outpatient area

Table 3.5: Standard precautions for infection control

Percentages of facilities with sterilization equipment somewhere in the facility and other items for standard precautions available in the general outpatient area of the facility on the day of the survey, by facility type, Bangladesh HFS 2022

| | | | | | | Facility typ | be | | | | | | | |
|--|--|--------------|------|------|---|--------------|-------------|--------------------------|----------------------------|---------------------|-------|---------------------------|-----------------|-------------------------------------|
| Items | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Community clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Sterilization equipment ¹ | 84.4 | 98.4 | 74.0 | 84.8 | 29.7 | 36.5 | 11.6 | 0.0 | 69.7 | 86.3 | 19.6 | 54.1 | 9.4 | 36.1 |
| Equipment for high-level disinfection ² | 67.0 | 82.3 | 57.4 | 67.0 | 37.7 | 43.6 | 22.1 | 6.4 | 58.7 | 53.5 | 20.8 | 46.0 | 15.5 | 41.1 |
| Safe final disposal of | | | | | | | | | | | | | | |
| sharps waste ³ Safe final disposal of | 60.3 | 54.8 | 61.4 | 60.9 | 49.6 | 54.1 | 37.7 | 56.3 | 76.5 | 73.0 | 57.4 | 59.4 | 54.9 | 50.9 |
| infectious waste4 | 56.0 | 58.1 | 60.1 | 54.7 | 41.7 | 44.6 | 34.0 | 40.3 | 78.7 | 76.2 | 45.9 | 55.9 | 41.1 | 43.4 |
| Appropriate storage of | 46.4 | 40.0 | 64.0 | 40.0 | E4 0 | 62.0 | 20.6 | 60.4 | co 7 | 24.0 | 50.0 | 47.0 | 60.4 | 50.0 |
| snarps waste | 46.1 | 40.3 | 61.2 | 43.3 | 54.8 | 63.9 | 30.6 | 62.4 | 68.7 | 31.2 | 56.9 | 47.2 | 60.1 | 53.8 |
| infoctious wasto ⁶ | 69.0 | 62.0 | 62.9 | 71.2 | 54 9 | 59.2 | 45 5 | 52.5 | 79 / | 63 / | 55.2 | 50.0 | 526 | 56 F |
| Disinfoctant ⁷ | 65.6 | 72.6 | 62.0 | 64.0 | 54.0 62.2 | 50.5 69.5 | 40.5 | 50.6 | 20.4 | 67.2 | 56.0 | 59.9 | 54.0 | 50.5 62.6 |
| Syringes and needles ⁸ | 58.1 | 72.0 58.1 | 76.3 | 53.6 | 71 / | 80.4 | 49.5 | 48.1 | 70.9 | 58.2 | 54.8 | 66.6 | 53.8 | 60.8 |
| Soon | 75.5 | 75.9 | 70.5 | 76.9 | 55.0 | 50.4 | 47.5 | 55.2 | 99.6 | 75.6 | 59.9 | 64.0 | 55.0 | 57.4 |
| Rupping water ⁹ | 88.5 | 88.7 | 81.0 | 00.0 | 56.8 | 61.2 | 42.0 | 50.8 | 87.9 | 70.0 | 57.1 | 68.2 | 53.3 | 60.6 |
| Soan and running water | 74.9 | 75.8 | 70.1 | 75.9 | 49.8 | 54.3 | 37.9 | 45.8 | 87.2 | 74.6 | 51.5 | 61.6 | 47.6 | 52.8 |
| Alcohol-based hand | 05.0 | 00.0 | 70.1 | 10.0 | 45.0 | 05.0 | 07.0 | 40.0 | 07.2 | 79.0 | 01.0 | 70.0 | 47.0 | 02.0 |
| Soap and running water or else alcohol-based hand | 85.2 | 90.3 | 77.0 | 86.4 | 65.5 | 65.6 | 65.4 | 66.6 | 89.0 | 79.9 | 68.9 | 72.9 | 67.0 | 67.9 |
| disinfectant | 94.0 | 96.8 | 90.0 | 94.6 | 75.8 | 77.1 | 72.4 | 75.1 | 96.7 | 86.1 | 77.5 | 81.6 | 75.9 | 78.0 |
| Latex gloves/other | | | | | | | | | | | | | | |
| gloves ¹⁰ | 70.2 | 62.9 | 84.0 | 68.0 | 71.7 | 80.7 | 47.9 | 59.8 | 84.1 | 74.0 | 64.6 | 73.0 | 62.9 | 71.6 |
| Medical masks | 81.5 | 82.3 | 73.2 | 83.4 | 69.9 | 71.2 | 66.3 | 84.8 | 83.8 | 75.0 | 80.6 | 73.1 | 81.3 | 71.3 |
| Gowns | 23.0 | 14.5 | 38.1 | 20.6 | 31.7 | 36.4 | 19.0 | 42.5 | 44.1 | 36.5 | 39.2 | 33.3 | 39.4 | 30.7 |
| Eye protection | 18.8 | 16.1 | 24.1 | 17.8 | 18.5 | 21.7 | 9.9 | 28.9 | 15.8 | 20.1 | 25.3 | 18.9 | 26.2 | 18.5 |
| precautions ¹¹ | 9.9 | 3.2 | 12.9 | 10.1 | 8.6 | 9.4 | 6.3 | 6.2 | 32.5 | 11.5 | 7.9 | 10.9 | 6.9 | 8.7 |
| Average number of the indicated items available | 9.6 | 9.6 | 9.8 | 9.5 | 7.8 | 8.6 | 5.8 | 7.1 | 11.2 | 9.6 | 7.7 | 8.7 | 7.4 | 8.0 |
| Number of facilities (weighted) | 41 | 4 | 7 | 30 | 310 | 225 | 85 | 994 | 29 | 183 | 1,557 | 563 | 1,345 | 351 |
| Number of facilities (unweighted) | 319 | 62 | 100 | 157 | 434 | 293 | 141 | 488 | 127 | 189 | 1,557 | 1,069 | 1,241 | 753 |

Note: The indicators presented in this table comprise the standard precautions domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012).

¹ Facility reports that some instruments are processed in the facility and the facility has a functioning electric dry heat sterilizer, a functioning electric autoclave, or a non-electric autoclave with a functioning heat source available somewhere in the facility.

² Facility reports that some instruments are processed in the facility and the facility has an electric pot or other pot with a heat source for high-level disinfection by boiling or high-level disinfection by steaming, or else facility had chlorine, formaldehyde, CIDEX, or glutaraldehyde for chemical high-level disinfection available somewhere in the facility on the day of the survey.

³ The process of sharps waste disposal is incineration and the facility had a functioning incinerator, or else the facility disposes of sharps waste by means of open burning in a Protected area, dumping without burning in a protected area, or removal offsite with storage in a protected area prior to removal offsite. ⁴ The process of infectious waste disposal is incineration and the facility had a functioning incinerator, or else the facility disposes of infectious waste by means of open burning

in a protected area, dumping without burning in a protected area, or removal offsite with storage in a protected area prior to removal offsite.

⁵ Sharps container observed in general outpatient service area

⁶ Waste receptacles observed in general outpatient service area

⁷ Chlorine-based or other country-specific disinfectants used for environmental disinfection available in the general outpatient area

⁸ Single-use standard disposable syringes with needles or else auto-disable syringes with needles available in the general outpatient area

⁹ Piped water, water in bucket with specially fitted tap, or water in pour pitcher available in the general outpatient area

¹⁰ Non-latex equivalent gloves are acceptable

¹¹ Any guideline for infection control in health facilities available in the general outpatient area

Table 3.6: Capacity for processing equipment for reuse

Percentage of facilities with the equipment and other items to support the final processing of instruments for reuse, by background characteristics, Bangladesh HFS 2022

| | | Percentage | of facilities having: | | _ | |
|------------------------------|------------------------|--|---|---|---------------------------------------|---|
| Background characteristic | Equipment ¹ | Equipment and knowledge of process time ² | Equipment, knowledge of process time, and automatic timer ³ | Written guidelines for sterilization or HLD ⁴ | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type | | | | | | |
| District and upazila | | | | | | |
| public facilities | 93.9 | 89.1 | 69.7 | 22.0 | 41 | 319 |
| DH | 100.0 | 100.0 | 88.7 | 14.5 | 4 | 62 |
| MCWC | 82.8 | 74.9 | 55.0 | 27.0 | 7 | 100 |
| UHC | 95.8 | 90.9 | 70.5 | 21.9 | 30 | 157 |
| Union-level public | | | | | | |
| facilities | 54.7 | 40.8 | 18.0 | 11.0 | 310 | 434 |
| UHFWC | 64.0 | 48.3 | 22.4 | 14.5 | 225 | 293 |
| USC/RD | 30.1 | 20.8 | 6.3 | 1.9 | 85 | 141 |
| Community clinic (CC) | 11.9 | 4.4 | 0.0 | 1.3 | 994 | 488 |
| NGO clinic/hospital | 78.2 | 74.5 | 59.1 | 42.6 | 29 | 127 |
| Private hospital | 92.0 | 85.7 | 53.7 | 4.8 | 183 | 189 |
| Location | | | | | | |
| Urban | 89.5 | 83.7 | 56.3 | 10.3 | 235 | 514 |
| Rural | 23.3 | 14.3 | 5.1 | 4.0 | 1,322 | 1,043 |
| Division | | | | | | |
| Barishal | 25.0 | 19.5 | 9.6 | 1.6 | 111 | 152 |
| Chattogram | 39.1 | 28.8 | 16.9 | 7.0 | 291 | 244 |
| Dhaka | 38.2 | 25.1 | 16.6 | 5.9 | 317 | 264 |
| Khulna | 39.4 | 30.9 | 9.0 | 3.4 | 210 | 208 |
| Rajshahi | 28.9 | 25.7 | 12.6 | 4.9 | 214 | 212 |
| Rangpur | 27.8 | 19.3 | 8.6 | 2.5 | 193 | 190 |
| Sylhet | 29.4 | 24.3 | 15.7 | 7.4 | 99 | 136 |
| Mymensingh | 23.2 | 16.4 | 8.0 | 5.2 | 122 | 151 |
| Total | 33.3 | 24.8 | 12.9 | 4.9 | 1,557 | 1,557 |
| Total excluding CCs | 70.9 | 60.7 | 35.5 | 11.5 | 563 | 1,069 |
| Total public | 24.3 | 15.4 | 6.3 | 4.2 | 1,345 | 1,241 |
| Total public excluding CCs | 59.3 | 46.5 | 24.1 | 12.3 | 351 | 753 |

¹ Facility reports that some equipment is processed in the facility and facility has a functioning electric dry heat sterilizer, a functioning electric autoclave, a non-electric autoclave with a functioning heat source, an electric boiler or steamer, or a non-electric boiler or steamer with a functioning heat source available anywhere in the facility or high-level disinfectant used for sterilization or high-level disinfection of equipment for reuse. ² Processing area has functioning equipment and power source for processing method and the responsible worker reports the correct

processing time (or equipment automatically sets the time) and processing temperature (if applicable) for at least one method. Definitions for capacity for each method assessed were functioning equipment and the following processing conditions:

• Dry heat sterilization: temperature at 160°C-169°C and processed for at least 120 minutes, or temperature at least 170°C and processed for at least 60 minutes

• Autoclave: wrapped items processed for at least 30 minutes, unwrapped items processed for at least 20 minutes

Boiling or steaming: items processed for at least 20 minutes
Chemical high-level disinfection: items processed in chlorine-based or glutaraldehyde or CIDEX or formaldehyde solution and

soaked for at least 20 minutes ³ An automatic timer here refers to a passive timer that can be set to indicate when a specified time has passed. It may be part of the sterilization process or the HLD equipment ⁴ Handwritten instructions that are pasted on walls and clearly outline the procedures to follow for processing of equipment are

acceptable

Table 3.7.1: Availability of laboratory diagnostic testing by facility type

Among all facilities, the percentages with basic and advanced laboratory diagnostic tests in the facility available, by background characteristics, Bangladesh HFS 2022

| | | | | | Facilit | y type | | | | | | | | |
|---|--|-------|------|------|---|--------|--------|----------------------------------|----------------------------|---------------------|-------|---------------------------|-----------------|-------------------------------------|
| Laboratory tests | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commu- nity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Basic tosts | | | | | | | | | | | | | | |
| Hemoglobin | 89.7 | 98.4 | 58.8 | 95 9 | 24.0 | 29.0 | 10.8 | 42 | 88.2 | 78 6 | 20.7 | 49 9 | 114 | 31.8 |
| Blood alucose | 91.5 | 100.0 | 60.7 | 97.7 | 40.9 | 45.6 | 28.3 | 61.2 | 96.8 | 90.1 | 62.0 | 63.5 | 57.4 | 46.8 |
| Urine protein | 81.3 | 88.7 | 61.4 | 85.0 | 36.2 | 45.8 | 10.8 | 3.7 | 86.0 | 73.2 | 21.9 | 54 1 | 13.6 | 41.5 |
| Urine alucose | 89.1 | 90.3 | 68.4 | 94.0 | 35.2 | 44.6 | 10.0 | 3.7 | 84.9 | 72.9 | 21.9 | 53.9 | 13.6 | 41.5 |
| TB microscopy | 37.5 | 48.4 | 0.0 | 45.0 | 0.6 | 0.0 | 2.0 | 0.0 | 0.0 | 12.7 | 2.6 | 72 | 1.3 | 4.9 |
| Syphilis rapid | 0110 | 10.1 | 0.0 | 10.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | | 2.0 | 7.2 | 1.0 | 1.0 |
| diagnostic test | 50.2 | 75.8 | 14.9 | 55.0 | 9.3 | 9.4 | 8.8 | 9.8 | 55.3 | 43.7 | 15.6 | 25.8 | 10.9 | 14.1 |
| General | | | | | | •••• | | | | | | | | |
| microscopy | 65.9 | 88.7 | 5.9 | 77.2 | 0.0 | 0.0 | 0.0 | 0.0 | 38.3 | 60.8 | 9.6 | 26.6 | 2.0 | 7.8 |
| Urine pregnancy | | | | | | | | | | | | | | |
| test | 80.4 | 90.3 | 42.4 | 88.2 | 24.4 | 11.0 | 3.0 | 3.0 | 90.0 | 75.4 | 19.6 | 48.0 | 10.3 | 31.0 |
| Advanced-level diagnostic tests | | | | | | | | | | | | | | |
| Full blood count | | | | | | | | | | | | | | |
| with differentials | 31.0 | 61.3 | 4.0 | 33.1 | 0.2 | 0.2 | 0.0 | 0.0 | 39.8 | 59.5 | 8.6 | 23.2 | 1.0 | 3.8 |
| Blood typing and | | | | | | | | | | | | | | |
| cross matching | 77.4 | 100.0 | 6.9 | 91.2 | 0.3 | 0.0 | 1.1 | 0.6 | 93.4 | 79.5 | 13.6 | 36.5 | 2.9 | 9.4 |
| Gram stain | 7.9 | 8.1 | 0.0 | 9.8 | 0.0 | 0.0 | 0.0 | 0.0 | 5.8 | 15.4 | 2.1 | 5.9 | 0.2 | 0.9 |
| Stool microscopy | 49.6 | 75.8 | 0.9 | 57.5 | 0.0 | 0.0 | 0.0 | 0.0 | 23.4 | 48.0 | 7.4 | 20.4 | 1.5 | 5.8 |
| TB culture | 3.3 | 6.5 | 0.0 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.3 | 0.8 | 2.3 | 0.1 | 0.4 |
| TB rapid diagnostic | , · · · | | | | | | | | | | | | | |
| test | 50.7 | 37.1 | 0.0 | 65.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.9 | 1.8 | 5.0 | 1.6 | 6.0 |
| Equipment for diagnostic imaging X-ray machine | | | | | | | | | | | | | | |
| (linked with TB) | 42.5 | 58.1 | 0.0 | 50.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 16.7 | 3.1 | 8.6 | 1.3 | 5.0 |
| X-ray machine | 52.5 | 96.8 | 0.0 | 58.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 54.9 | 7.9 | 21.7 | 1.6 | 6.2 |
| Ultrasonogram | 52.2 | 90.3 | 4.8 | 58.1 | 0.0 | 0.0 | 0.0 | 0.0 | 42.9 | 75.9 | 11.1 | 30.7 | 1.6 | 6.1 |
| CT scan | 1.5 | 14.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.8 | 0.4 | 1.0 | 0.0 | 0.2 |
| Number of facilities (weighted) | 41 | 4 | 7 | 30 | 310 | 225 | 85 | 994 | 29 | 83 | 1,557 | 563 | 1,345 | 351 |
| Number of facilities (unweighted) | 319 | 62 | 100 | 157 | 434 | 293 | 141 | 488 | 127 | 189 | 1,557 | 1,069 | 1,241 | 753 |

CT = Computed tomography

Table 3.7.2: Laboratory diagnostic capacity by facility type

Among all facilities, the percentages with the capacity to conduct basic and advanced laboratory diagnostic tests in the facility, by background characteristics, Bangladesh HFS 2022

| | | | | | Facili | ty type | | | | | | | | |
|--|--|------------------------------|--------------------------|-----------------------------|---|--------------------------|--------------------------|---------------------------------|----------------------------|-----------------------------|---------------------------|----------------------------|--------------------------|-------------------------------------|
| Laboratory tests | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commu nity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Basic tests | | | | | | | | | | | | | | |
| Hemoglobin | 86.4 | 93.5 | 58.8 | 92.0 | 23.9 | 28.8 | 10.8 | 4.2 | 87.0 | 77.6 | 20.5 | 49.2 | 11.2 | 31.2 |
| Blood glucose | 72.2 | 50.0 | 57.8 | 79.0 | 39.3 | 44.2 | 26.1 | 58.2 | 93.0 | 74.2 | 57.3 | 55.8 | 54.3 | 43.1 |
| Urine protein | 76.6 | 87.1 | 58.4 | 79.5 | 34.6 | 43.5 | 10.8 | 2.7 | 84.4 | 73.2 | 20.8 | 52.8 | 12.3 | 39.5 |
| Urine glucose | 86.1 | 88.7 | 65.4 | 90.7 | 33.5 | 42.4 | 10.0 | 2.9 | 83.3 | 72.9 | 20.9 | 52.7 | 12.5 | 39.7 |
| TB microscopy Syphilis rapid | 15.5 | 19.4 | 0.0 | 18.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.8 | 1.1 | 3.0 | 0.5 | 1.8 |
| diagnostic test General | 50.2 | 75.8 | 14.9 | 55.0 | 9.3 | 9.4 | 8.8 | 9.8 | 55.3 | 43.7 | 15.6 | 25.8 | 10.9 | 14.1 |
| microscopy Urine pregnancy | 61.9 | 85.5 | 5.9 | 72.0 | 0.0 | 0.0 | 0.0 | 0.0 | 37.0 | 60.8 | 9.5 | 26.2 | 1.9 | 7.3 |
| test Average basic | 80.4 | 90.3 | 42.4 | 88.2 | 24.4 | 29.4 | 11.0 | 3.0 | 90.0 | 75.4 | 19.6 | 48.5 | 10.3 | 31.0 |
| testing capacity ¹ | 4.0 | 4.1 | 2.8 | 4.3 | 1.6 | 1.9 | 0.7 | 0.7 | 4.4 | 3.7 | 1.4 | 2.6 | 1.0 | 1.8 |
| Advanced-level diagnostic tests Full blood count with differentials | 31.0 | 61.3 | 4.0 | 33.1 | 0.2 | 0.2 | 0.0 | 0.0 | 39.8 | 59.5 | 8.6 | 23.7 | 1.0 | 3.8 |
| Blood typing and cross matching | 2.5 | 4.8 | 0.0 | 2.7 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 7.4 | 1.0 | 2.9 | 0.1 | 0.3 |
| Syphilis serology Gram stain Stool microscopy TB culture | 6.6 36.9 1.8 | 8.1 61.3 1.6 | 0.0 0.9 | 8.1 42.0 2 3 | 0.0 0.0 | 0.0 0.0 0.0 | 0.0 0.0 0.0 | 0.0 0.0 0.0 | 4.7 14.6 0.0 | 12.4 36.4 0.7 | 1.7 5.5 0 1 | 4.8 15.3 0.4 | 0.2 1.1 0.1 | 0.8 4.3 0.2 |
| TB rapid diagnostic test | 35.5 | 29.0 | 0.0 | 45.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.9 | 1.4 | 3.9 | 1.1 | 4.2 |
| Equipment for diagnostic imaging X-ray machine | | | | | | | | | | | | | | |
| (linked with TB) X-ray machine Ultrasonogram CT scan | 41.0 42.7 44.1 1.4 | 75.8 87.1 79.0 12.9 | 0.0 0.0 4.8 0.0 | 45.8 46.5 48.5 0.0 | 0.0 0.0 0.0 0.0 | 0.0 0.0 0.0 0.0 | 0.0 0.0 0.0 0.0 | 0.0 0.0 0.0 0.0 | 0.0 0.3 41.6 0.0 | 18.4 53.8 74.9 2.8 | 3.2 7.5 10.8 0.4 | 9.0 20.6 29.7 1.0 | 1.3 1.3 1.4 0.0 | 4.8 5.0 5.2 0.2 |
| Number of facilities (weighted) | 41 | 4 | 7 | 30 | 310 | 225 | 85 | 994 | 29 | 83 | 1,557 | 563 | 1,345 | 351 |
| Number of facilities (unweighted) | 319 | 62 | 100 | 157 | 434 | 293 | 141 | 488 | 127 | 189 | 1,557 | 1.069 | 1.241 | 753 |

Note: The basic test indicators presented in this table comprise the diagnostic capacity domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012).

CT = Computed tomography ¹ The capacity to conduct average number of basic diagnostic tests (hemoglobin, blood glucose, urine protein, urine glucose, and urine pregnancy test)

Table 3.7.3: Laboratory diagnostic capacity by location and division

Among all facilities, the percentages with the capacity to conduct basic and advanced laboratory diagnostic tests in the facility, by background characteristics, Bangladesh HFS 2022

| | Loc | ation | | | | Di | vision | | | | |
|--|-------|-------|----------|--------|-------|--------|----------|---------|--------|------------|-------|
| | | | | Chat- | | | | | | | |
| Laboratory tests | Urban | Rural | Barishal | togram | Dhaka | Khulna | Rajshahi | Rangpur | Sylhet | Mymensingh | Total |
| Basic tests | | | | | | | | | | | |
| Hemoglobin | 77.3 | 10.3 | 11.4 | 26.8 | 26.5 | 17.3 | 22.9 | 10.2 | 23.7 | 12.9 | 20.5 |
| Blood glucose | 73.8 | 54.4 | 37.9 | 57.5 | 63.0 | 62.5 | 70.5 | 46.2 | 53.9 | 48.0 | 57.3 |
| Urine protein | 72.3 | 11.6 | 12.5 | 25.6 | 27.3 | 17.3 | 20.6 | 13.4 | 26.1 | 13.6 | 20.8 |
| Urine glucose | 73.3 | 11.6 | 13.8 | 26.9 | 26.6 | 16.8 | 21.0 | 14.0 | 27.0 | 11.3 | 20.9 |
| TB microscopy | 6.0 | 0.2 | 1.7 | 1.4 | 1.9 | 0.3 | 0.9 | 0.6 | 1.2 | 0.2 | 1.1 |
| Syphilis rapid | | | | | | | | | | | |
| diagnostic test | 45.4 | 10.3 | 20.3 | 11.5 | 26.0 | 14.7 | 14.7 | 10.5 | 16.2 | 4.8 | 15.6 |
| General microscopy | 56.5 | 1.1 | 5.9 | 10.9 | 12.7 | 9.4 | 11.8 | 5.4 | 7.7 | 5.1 | 9.5 |
| Urine pregnancy test | 75.9 | 9.4 | 10.5 | 28.2 | 25.1 | 17.5 | 17.2 | 14.0 | 14.0 | 12.4 | 19.5 |
| capacity ¹ | 3.7 | 1.0 | 0.9 | 1.6 | 1.7 | 1.3 | 1.5 | 1.0 | 1.4 | 1.0 | 1.4 |
| Advanced-level | | | | | | | | | | | |
| diagnostic tests | | | | | | | | | | | |
| Full blood count with | | | | | | | | | | | |
| differentials | 53.4 | 0.6 | 5.6 | 12.5 | 12.9 | 4.2 | 9.9 | 3.6 | 7.7 | 4.7 | 8.6 |
| Blood typing and | | | | | | | | | | | |
| cross matching | 6.6 | 0.0 | 0.8 | 0.6 | 2.3 | 1.0 | 0.5 | 0.8 | 0.7 | 0.5 | 1.0 |
| Syphilis serology | | | | | | | | | | | |
| Gram stain | 10.7 | 0.1 | 0.7 | 1.7 | 5.0 | 0.3 | 0.4 | 0.8 | 1.5 | 0.6 | 1.7 |
| Stool microscopy | 34.0 | 0.5 | 4.9 | 7.2 | 9.0 | 2.0 | 6.2 | 2.8 | 5.6 | 2.4 | 5.5 |
| TB culture | 0.8 | 0.0 | 0.1 | 0.6 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| TB rapid diagnostic | | | | | | | | | | | |
| test | 6.6 | 0.5 | 1.6 | 2.6 | 1.0 | 1.6 | 0.8 | 0.9 | 1.5 | 1.0 | 1.4 |
| Equipment for diagnostic imaging X-ray machine | | | | | | | | | | | |
| (linked with TB) | 19.2 | 0.4 | 1.8 | 4.9 | 4.5 | 1.9 | 3.2 | 1.6 | 3.7 | 2.1 | 3.2 |
| X-ray machine | 46.3 | 0.6 | 5.6 | 11.0 | 11.2 | 4.5 | 8.3 | 3.2 | 4.8 | 3.5 | 7.5 |
| Ultrasonogram | 66.1 | 0.9 | 6.2 | 13.7 | 15.6 | 11.1 | 11.7 | 4.5 | 8.2 | 5.0 | 10.8 |
| CT scan | 2.4 | 0.0 | 0.1 | 0.0 | 1.1 | 0.9 | 0.1 | 0.0 | 0.0 | 0.0 | 0.4 |
| Number of facilities (weighted) | 235 | 1,322 | 111 | 291 | 318 | 210 | 214 | 193 | 99 | 122 | 1,557 |
| Number of facilities (unweighted) | 319 | 62 | 100 | 157 | 434 | 293 | 141 | 488 | 127 | 189 | 1,557 |

Note: The basic test indicators presented in this table comprise the diagnostic capacity domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012). CT = Computed tomography ¹ The capacity to conduct average number of basic diagnostic tests (hemoglobin, blood glucose, urine protein, urine glucose, and urine pregnancy test)

Table 3.8.1: Availability of essential medicines by facility type

Percentages of facilities having the 14 essential medicines available, by background characteristics, Bangladesh HFS 2022

| | | | | | Facilit | y type | | | | | | | | |
|---|--|------|------|-------|---|--------|--------|------------------------------|----------------------------|---------------------|-------|---------------------------|-----------------|-------------------------------------|
| Essential medicines | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commun ity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Amitriptyline tablets/capsules1 | 7.4 | 9.7 | 0.0 | 8.9 | 0.9 | 0.2 | 2.5 | 0.2 | 10.8 | 28.7 | 4.1 | 10.9 | 0.6 | 1.6 |
| Amoxicillin tablets/capsules ² | 86.8 | 67.7 | 89.0 | 89.0 | 89.9 | 91.3 | 86.3 | 91.6 | 55.4 | 27.5 | 82.9 | 67.6 | 91.0 | 90.0 |
| Atenolol tablets/capsules ³ | 58.1 | 37.1 | 0.9 | 75.2 | 8.9 | 0.3 | 31.7 | 0.2 | 12.5 | 25.8 | 6.7 | 18.2 | 4.0 | 14.7 |
| Captopril tablets/capsules ⁴ | 82.4 | 96.8 | 5.9 | 98.9 | 16.9 | 2.6 | 54.7 | 0.4 | 43.9 | 55.4 | 13.2 | 35.6 | 6.8 | 24.6 |
| Ceftriaxone injectable ⁵ | 89.2 | 95.2 | 51.1 | 97.6 | 17.0 | 20.7 | 7.2 | 4.4 | 34.6 | 60.6 | 16.3 | 37.4 | 9.9 | 25.5 |
| Ciprofloxacin tablets/capsules ⁶ | 99.0 | 98.4 | 95.0 | 100.0 | 97.5 | 97.2 | 98.4 | 97.1 | 91.2 | 58.8 | 92.6 | 84.7 | 97.2 | 97.7 |
| Cotrimoxazole oral | | | | | | | | | | | | | | |
| suspension ⁷ | 86.0 | 75.8 | 93.1 | 85.8 | 84.4 | 92.6 | 62.7 | 94.0 | 53.8 | 44.9 | 85.3 | 70.1 | 91.5 | 84.6 |
| Diazepam tablets/capsules8 | 90.5 | 64.5 | 90.3 | 94.4 | 69.7 | 82.8 | 35.2 | 3.7 | 35.6 | 46.7 | 24.8 | 62.0 | 21.6 | 72.2 |
| Diclofenac tablets/capsules9 | 96.0 | 98.4 | 80.1 | 99.5 | 65.1 | 61.1 | 75.6 | 69.6 | 84.6 | 59.9 | 68.6 | 66.6 | 69.4 | 68.7 |
| Glibenclamide tablets/ | | | | | | | | | | | | | | |
| capsules ¹⁰ | 78.3 | 91.9 | 3.0 | 94.6 | 10.8 | 1.0 | 36.8 | 0.2 | 16.6 | 40.5 | 9.5 | 25.7 | 5.1 | 18.8 |
| Omeprazole/cimetidine | | | | | | | | | | | | | | |
| tablets/capsules11 | 97.7 | 96.8 | 93.1 | 99.0 | 91.8 | 93.4 | 87.6 | 92.6 | 89.0 | 60.1 | 88.7 | 81.8 | 92.6 | 92.5 |
| Paracetamol oral suspension ¹² | 86.4 | 88.7 | 85.1 | 86.3 | 83.6 | 91.0 | 64.2 | 96.0 | 64.4 | 44.9 | 86.7 | 70.2 | 93.9 | 84.0 |
| Salbutamol inhaler ¹³ | 97.2 | 98.4 | 86.9 | 99.5 | 86.6 | 88.3 | 82.1 | 93.8 | 57.1 | 58.7 | 87.6 | 76.8 | 92.2 | 87.8 |
| Simvastatin/atorvastatin | | | | | | | | | | | | | | |
| tablets/capsules14 | 73.6 | 87.1 | 2.0 | 89.1 | 6.9 | 1.5 | 21.1 | 0.0 | 2.6 | 36.5 | 7.7 | 21.2 | 3.9 | 14.7 |
| Number of facilities (weighted) | 41 | 4 | 7 | 30 | 310 | 225 | 85 | 994 | 29 | 183 | 1,557 | 563 | 1,345 | 351 |
| Number of facilities | 210 | 62 | 100 | 157 | 434 | 202 | 1.4.1 | 499 | 127 | 190 | 1 557 | 1.060 | 1 2/1 | 752 |
| (unweighted) | 319 | 02 | 100 | 157 | 434 | 293 | 141 | 400 | 121 | 103 | 1,007 | 1,009 | 1,241 | 100 |

Note: The indicators presented in this table comprise the essential medicines domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012).

¹ For management of depression in adults

² First-line antibiotic for adults

³ Beta blocker for management of angina/hypertension

⁴ Vasodilator for management of hypertension

⁵ Second-line injectable antibiotic

⁶ Second-line oral antibiotic

⁷ Oral antibiotic for children

⁸ Muscle relaxant for management of anxiety or seizures

⁹ Oral analgesic

¹⁰ For management of type 2 diabetes

¹¹ Proton pump inhibitor for treatment of peptic ulcer disease, dyspepsia, and gastroesophageal reflux disease

¹² Fever reducer and analgesic for children

¹³ For management and relief of bronchospasms in conditions such as asthma and chronic obstructive pulmonary disease

¹⁴ For control of elevated cholesterol

Table 3.8.2: Availability of essential medicines by location and division

Percentages of facilities having the 14 essential medicines available, by background characteristics, Bangladesh HFS 2022

| | Loc | ation | | | | Div | ision | | | | |
|---|-------|-------|----------|------------|-------|--------|----------|---------|--------|------------|-------|
| Essential medicines | Urban | Rural | Barishal | Chattogram | Dhaka | Khulna | Rajshahi | Rangpur | Sylhet | Mymensingh | Total |
| Amitriptyline tablets/ | | | | | | | | | | | |
| capsules ¹ | 24.0 | 0.5 | 3.0 | 6.1 | 6.0 | 4.3 | 2.4 | 1.9 | 2.7 | 2.3 | 4.1 |
| Amoxicillin tablets/capsules ² | 37.5 | 91.0 | 84.4 | 82.3 | 81.5 | 74.8 | 84.2 | 86.1 | 85.3 | 91.2 | 82.9 |
| Atenolol tablets/capsules ³ | 27.6 | 3.0 | 5.1 | 8.6 | 9.3 | 5.5 | 6.3 | 4.5 | 4.6 | 5.3 | 6.7 |
| Captopril tablets/capsules ⁴ | 55.8 | 5.6 | 8.5 | 16.7 | 17.5 | 13.5 | 13.2 | 8.2 | 8.6 | 8.8 | 13.2 |
| Ceftriaxone injectable ⁵ | 59.0 | 8.7 | 11.5 | 19.1 | 23.5 | 16.4 | 11.1 | 10.3 | 16.1 | 14.3 | 16.3 |
| Ciprofloxacin tablets/ | | | | | | | | | | | |
| capsules ⁶ | 65.9 | 97.4 | 94.5 | 93.3 | 93.2 | 88.2 | 91.9 | 93.7 | 92.4 | 94.8 | 92.6 |
| Cotrimoxazole oral | | | | | | | | | | | |
| suspension ⁷ | 49.2 | 91.8 | 89.2 | 86.7 | 84.2 | 81.6 | 81.1 | 86.1 | 89.6 | 90.7 | 85.3 |
| Diazepam tablets/capsules8 | 49.5 | 20.4 | 25.0 | 30.2 | 28.5 | 21.8 | 20.2 | 24.5 | 23.5 | 17.3 | 24.8 |
| Diclofenac tablets/capsules9 | 66.1 | 69.0 | 59.9 | 64.5 | 69.4 | 68.7 | 69.8 | 74.4 | 61.8 | 77.8 | 68.6 |
| Glibenclamide tablets/ | | | | | | | | | | | |
| capsules ¹⁰ | 40.8 | 3.9 | 6.2 | 14.5 | 13.3 | 7.8 | 7.0 | 3.8 | 7.9 | 8.0 | 9.5 |
| Omeprazole/cimetidine | | | | | | | | | | | |
| tablets/capsules11 | 66.7 | 92.6 | 90.5 | 84.7 | 87.2 | 85.0 | 90.3 | 96.1 | 91.3 | 90.5 | 88.7 |
| Paracetamol oral | | | | | | | | | | | |
| suspension ¹² | 51.0 | 93.0 | 91.0 | 90.8 | 88.8 | 81.0 | 80.7 | 84.4 | 89.6 | 88.8 | 86.7 |
| Salbutamol inhaler ¹³ | 63.0 | 92.0 | 89.3 | 88.4 | 86.3 | 84.2 | 87.0 | 91.2 | 82.4 | 92.8 | 87.6 |
| Simvastatin/atorvastatin | | | | | | | | | | | |
| tablets/capsules14 | 36.0 | 2.6 | 5.6 | 11.8 | 11.8 | 3.6 | 5.0 | 4.9 | 7.0 | 5.7 | 7.7 |
| | | | | | | | | | | | |
| Number of facilities | | | | | | | | | | | |
| (weighted) | 235 | 1,322 | 111 | 291 | 318 | 210 | 214 | 193 | 99 | 122 | 1,557 |
| Number of facilities | | | | | | | | | | | |
| (upweighted) | 210 | 60 | 100 | 157 | 124 | 202 | 1.1.1 | 100 | 107 | 190 | 1 557 |
| (unweighted) | 319 | 62 | 100 | 157 | 434 | 293 | 141 | 400 | 127 | 169 | 1,557 |

Note: The indicators presented in this table comprise the essential medicines domain for assessing general service readiness within the health facility assessment methodology proposed by WHO and USAID (WHO 2012).

¹ For management of depression in adults

² First-line antibiotic for adults

³ Beta blocker for management of angina/hypertension

⁴ Vasodilator for management of hypertension

⁵ Second-line injectable antibiotic

⁶ Second-line oral antibiotic

⁷ Oral antibiotic for children

⁸ Muscle relaxant for management of anxiety or seizures

9 Oral analgesic

¹⁰ For management of type 2 diabetes

¹¹ Proton pump inhibitor for treatment of peptic ulcer disease, dyspepsia, and gastroesophageal reflux disease

¹² Fever reducer and analgesic for children

¹³ For management and relief of bronchospasms in conditions such as asthma and chronic obstructive pulmonary disease

¹⁴ For control of elevated cholesterol

Table 3.8.3: Availability of medicines in community clinics

Percentages of community clinics having the indicated medicines available on the day of survey, Bangladesh HFS 2022

| Medicines | Percentage of community clinics |
|---|------------------------------------|
| Albendazole tablets ¹ | 96.6 |
| Amoxicillin capsules, syrup and paediatric drops ² | 95.8 |
| Antacid tablets ³ | 92.4 |
| Benzoic and salicylic acid ointment ⁴ | 79.5 |
| Calcium lactate tablets ⁵ | 86.6 |
| Chlorpheniramine tablet and syrup ⁶ | 89.3 |
| Co-trimoxazole tablets (120mg and 960 mg) ⁷ | 87.6 |
| Doxycycline capsule ⁸ | 100.0 |
| Erythromycin tablets and syrup ⁹ | 0.2 |
| Ferrous fumarate & folic acid tablets ¹⁰ | 81.4 |
| Gentian violet ¹¹ | 20.9 |
| Hyoscine-n-butyl bromide tablets ¹² | 67.6 |
| Metronidazole tablets ¹³ | 97.2 |
| Neomycin and bacitracin skin ointment ¹⁴ | 71.3 |
| Paracetamol tablets and Syrup ¹⁵ | 98.2 |
| Penicillin tablets ¹⁶ | 81.8 |
| Salbutamol syrup ¹⁷ | 87.0 |
| Vitamin-B complex tablets ¹⁸ | 97.0 |
| Zinc dispersible tablets ¹⁹ | 79.8 |
| Number of facilities (weighted) | 994 |
| Number of facilities (unweighted) | 488 |

Note: These are the medicines routinely supplied in community clinics for patient management. ¹ Anti-helminthics for the management of worm infestation

²Broad-spectrum oral antibiotics

³ Stomach acid neutralizers to relieve indigestion and heartburn

⁴ Ointment for the management of skin infections

⁵ Drugs for the management of calcium deficiency ⁶ First-generation sedating antihistamine

⁷ Combined anti-biotics for the management of bacterial infections

8 Oral antibiotics of tetra-cyclin group

⁹ Oral antibiotics of Macrolides group

¹⁰ Combined supplements to mitigate iron and folic acid deficiency in

¹¹ Antiseptic solution to treat skin infections

¹² Antiseptic solution to treat skin infections
 ¹² Antiseptic solution to treat skin infections
 ¹³ Antiseptic solution to treat skin infections
 ¹³ Oral anti-protozoal drugs
 ¹⁴ Combined antibiotic ointment for the management of skin infections
 ¹⁵ Environment antibiotic ointment for the management of skin infections

¹⁵ Fever reducer (antipyretic) and analgesic drugs

¹⁶ Broad-spectrum oral antibiotics

¹⁷ Drugs for management and relief of bronchospasms in conditions such

as asthma and chronic obstructive pulmonary disease ¹⁸ Combined vitamin B preparations to mitigate vitamin B deficiency

¹⁹ For the management of diarrhea in children

Table 3.8.4: Availability of drug and dietary supply (DDS) kit medicines

Percentages of facilities having the indicated medicines available on the day of survey, Bangladesh HFS 2022

| | Facilit | у Туре |
|---|---------|--------|
| Medicines | MCWC | UHFWC |
| Albendazole tablets ¹ | 96.0 | 98.2 |
| Amoxicillin suspension ² | 89.0 | 91.3 |
| Amoxicillin capsules ² (500 mg) | 89.0 | 91.3 |
| Antacid tablets ³ | 82.8 | 89.1 |
| Benzoic acid and salicylic acid ⁴ | 80.8 | 80.8 |
| Calcium carbonate tablets ⁵ | 17.9 | 15.2 |
| Chlorpheniramine maleate tablets ⁶ | 79.1 | 79.6 |
| Cotrimoxazole ⁷ (120 mg) dispersible | 80.4 | 83.9 |
| Cotrimoxazole DS (960 mg) | 81.1 | 83.3 |
| Cotrimoxazole suspension | 69.2 | 66.0 |
| Diazepam tablets ⁸ | 90.3 | 82.8 |
| Doxycycline (100 mg) capsules ⁹ | 71.7 | 70.1 |
| Drotaverine hydrochloride tablets ¹⁰ | 45.7 | 39.5 |
| Folic acid tablets ¹¹ | 96.0 | 94.5 |
| Iron and folic acid tablets ¹² | 96.0 | 98.4 |
| Metronidazole tablets ¹³ | 90.0 | 90.1 |
| Pantoprazole tablets ¹⁴ | 90.0 | 90.1 |
| Paracetamol ¹⁵ tablets | 91.9 | 96.2 |
| Paracetamol suspension | 84.1 | 91.0 |
| Salbutamol tablets ¹⁶ | 84.9 | 85.8 |
| Vitamin B complex tablets ¹⁷ | 94.8 | 94.3 |
| | | |
| Number of facilities (weighted) | 7 | 225 |
| Number of facilities (unweighted) | 100 | 293 |

Note: A DDS kit consists of pre-selected essential drugs and dietary items to be dispensed by service providers at MCWCs and UHFWCs for patient management. They receive DDS kits at different intervals from the central warehouse.

¹ Anti-helminthics for the management of worm infestation

² Broad-spectrum oral antibiotics

- ³ Stomach acid neutralizers to relieve indigestion and heartburn
- ⁴ Ointment for the management of skin infections
- ⁵ Drugs for the management of calcium deficiency
- ⁶ First-generation sedating antihistamine

```
    <sup>9</sup> Combined anti-biotics for the management of bacterial infections
    <sup>8</sup> Sedative drugs for management of anxiety or seizures
    <sup>9</sup> Oral antibiotics of tetra-cyclin group
    <sup>10</sup> Antispasmodics for the treatment of abdominal pain induced by
```

gastrointestinal/genitourinary spasms¹¹ Supplements to mitigate iron and folic acid deficiency in pregnancy

¹² Supplements to mitigate folic acid deficiency in pregnancy

¹³ Oral anti-protozoal drugs

¹⁴ Proton pump inhibitor for treatment of peptic ulcer disease, dyspepsia, and gastroesophageal reflux disease ¹⁵ Fever reducer (antipyretic) and analgesic drugs

¹⁶ Drugs for management and relief of bronchospasms in conditions such as asthma and chronic obstructive pulmonary disease

¹⁷ Combined vitamin B preparations to mitigate vitamin B deficiency

Table 3.8.5: Availability of medicines and equipment of delivery kit

Percentages of facilities having the indicated medicines and equipment available on the day of survey, Bangladesh HFS 2022

| | Facility Type | | | | |
|---|---------------|-------|--|--|--|
| Items | MCWC | UHFWC | | | |
| Paracetamol ¹ | 91.9 | 96.2 | | | |
| Amoxicillin capsules ² | 89.0 | 91.3 | | | |
| Infusion 5% dextrose in aqua ³ (500 ml) | 78.3 | 43.3 | | | |
| Infusion 5% dextrose in normal saline ³ (500 ml) | 68.3 | 38.0 | | | |
| Chlorhexidine ⁴ cream (60 ml bottle) | 84.2 | 70.5 | | | |
| Sterile surgical gloves | 95.0 | 95.5 | | | |
| Cotton ball thread | 96.0 | 89.9 | | | |
| Sterile gauge pack | 92.9 | 72.3 | | | |
| Surgical adhesive tape | 63.4 | 4.8 | | | |
| 1% lidocaine injection ⁵ (50 ml) | 50.1 | 19.8 | | | |
| Disposable syringe (5ml) | 94.9 | 95.8 | | | |
| Number of facilities (weighted) | 7 | 225 | | | |
| Number of facilities (unweighted) | 100 | 293 | | | |

Note: A delivery kit comprises essential medicines, infusions, and logistical items necessary for conducting a safe normal delivery. These kits are supplied from the central warehouse and are dispensed by service providers at MCWCs and UHFWCs for patient management. ¹ Fever reducer (antipyretic) and analgesic drugs ² Broad-spectrum oral antibiotics

³ Intravenous solutions for the management of hypovolemia
 ⁴ Antiseptic agents for infection prevention
 ⁵ Local anesthetic agents

Table 3.9: Supportive management practices at the facility level

Among all facilities, the percentages that had an external supervisory visit during the 6 months before the survey, and the percentages of facilities where at least half of the interviewed providers reported receiving routine work-related training and personal supervision recently, by background characteristics, Bangladesh HFS 2022

| | | | | | | | | Number of facilities where | Number of facilities where at |
|--|---|---------------------------------------|---|-----------------------------|-----------------------------------|---|--|--|---|
| | Percentage of | | | Pe | rcentage of faci | at least two | least two eligible | | |
| Background characteristic | facilities with supervisory visit during the 6 months before the survey ¹ | Number of facilities (weighted) | Number of facilities (unweighted) | Staff training ² | Personal supervision ³ | Training and personal supervision | Percentage of facilities with supportive management practices ⁴ | eligible providers were interviewed with health worker interview questionnaire (weighted) | providers were interviewed with health worker interview questionnaire (unweighted) |
| Facility type | | | | | | | | | |
| District and upazila | | | | | | | | | |
| public facilities | 97.9 | 41 | 319 | 59.4 | 97.8 | 56.9 | 56.3 | 41 | 308 |
| DH | 100.0 | 4 | 62 | 50.0 | 96.8 | 46.8 | 46.8 | 4 | 62 |
| MCWC | 94.0 | 7 | 100 | 54.4 | 96.6 | 53.3 | 52.2 | 7 | 89 |
| UHC | 98.6 | 30 | 157 | 61.9 | 98.3 | 59.2 | 58.6 | 30 | 157 |
| Union-level public | | | | | | | | | |
| facilities | 83.9 | 310 | 434 | 66.4 | 92.3 | 58.7 | 54.8 | 171 | 244 |
| UHFWC | 83.9 | 225 | 293 | 68.8 | 91.9 | 61.4 | 57.8 | 118 | 156 |
| USC/RD | 83.8 | 85 | 141 | 61.0 | 93.3 | 52.7 | 48.1 | 53 | 88 |
| Community clinic (CC) NGO clinic/hospital Private bospital | 89.5 90.2 84.4 | 994 29 183 | 488 127 189 | 75.8 62.2 | 93.5 99.2 86.8 | 72.7 62.2 14 7 | 69.9 58.9 14 7 | 434 26 167 | 210 113 117 |
| i mate nospital | 04.4 | 100 | 100 | 10.0 | 00.0 | 14.7 | 14.7 | 107 | |
| Location | | | | | | | | | |
| Urban | 87.2 | 235 | 514 | 26.0 | 90.0 | 25.0 | 24.8 | 213 | 487 |
| Rural | 88.2 | 1,322 | 1,043 | 72.6 | 93.1 | 68.2 | 65.1 | 626 | 564 |
| Division | | | | | | | | | |
| Barishal | 80.0 | 111 | 152 | 70.9 | 93.4 | 66.2 | 62.7 | 49 | 103 |
| Chattogram | 86.2 | 291 | 244 | 63.9 | 91.9 | 61.3 | 59.6 | 143 | 158 |
| Dhaka | 86.1 | 318 | 264 | 55.7 | 92.6 | 52.6 | 50.9 | 186 | 185 |
| Khulna | 87.7 | 210 | 208 | 48.9 | 88.6 | 47.0 | 44.4 | 103 | 120 |
| Rajshahi | 89.8 | 214 | 212 | 57.6 | 91.6 | 53.2 | 50.0 | 138 | 160 |
| Rangpur | 92.5 | 193 | 190 | 62.1 | 93.9 | 58.6 | 55.7 | 121 | 142 |
| Sylhet | 93.7 | 99 | 136 | 74.1 | 96.1 | 71.5 | 71.5 | 39 | 81 |
| Mymensingh | 90.5 | 122 | 151 | 77.2 | 93.6 | 68.8 | 65.7 | 60 | 102 |
| Total | 88.0 | 1,557 | 1,557 | 60.8 | 92.3 | 57.2 | 54.8 | 839 | 1,051 |
| Total excluding CCs | 85.4 | 563 | 1,069 | 44.6 | 91.0 | 40.6 | 38.6 | 405 | 841 |
| Total public | 88.5 | 351 | 1,241 | 72.3 | 93.4 | 68.0 | 65.0 | 646 | 762 |
| Total public excluding CCs | 85.5 | 1,345 | 753 | 65.0 | 93.4 | 58.4 | 55.1 | 212 | 552 |

¹ Facility reports that it received at least one external supervisory visit from the district, regional, or national office during the 6 months before the survey.
² At least half of all interviewed providers reported that they had received any in-service training as part of their work in the facility during the 24 months before the survey. This training refers to structured sessions; it does not include individual instruction that a provider might have received during routine supervision.
³ At least half of all interviewed providers reported that they had been personally supervised at least once during the 6 months before the survey. Personal supervision refers to any there has before the survey. Personal supervision refers to any there has a survey of the service of the service the survey of the service the survey.

form of technical support or supervision from a facility-based supervisor or from a visiting supervisor. It may include, but is not limited to, review of records and observation of work, with or without any feedback to the health worker. ⁴ Facility had an external supervisory visit during the 6 months before the survey, and staff members have received routine training and supervision.

Table 3.10: Quality assurance and client opinion

Among all facilities, the percentages having quality assurance activities and documentation of quality assurance activities, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Percentage of facilities with regular quality assurance activities and observed documentation of quality assurance activities ¹ | Number of facilities (weighted) | Number of facilities (unweighted) |
|--|--|---|--|
| Facility type | | | |
| District and upazila public facilities DH MCWC UHC | 54.9 66.1 21.8 61.3 | 41 4 7 30 | 319 62 100 157 |
| Union-level public facilities UHFWC USC/RD | 15.6 17.9 9.5 | 310 225 85 | 434 293 141 |
| Community clinic (CC) | 16.5 | 994 | 488 |
| NGO clinic/hospital | 48.4 | 29 | 127 |
| Private hospital | 15.9 | 183 | 189 |
| Location Urban Rural | 25.0 16.6 | 235 1,322 | 514 1,043 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 7.4 18.5 14.9 17.9 18.5 14.4 22.0 34.6 | 112 291 317 210 214 193 99 122 | 152 244 264 208 212 190 136 151 |
| Total | 17.8 | 1,557 | 1,557 |
| Total excluding CCs Total public | 20.3 <i>17.5</i> | 563 1,345 | 1,069 1,241 |
| Total public excluding CCs | 20.2 | 351 | 753 |

¹ Facility reports that it routinely carries out quality assurance activities and had documentation of a recent quality assurance activity. This could be a report or minutes of a quality assurance meeting, a supervisory checklist, a mortality review, or an audit of records or registers.

Table 3.11.1: Staffing pattern in surveyed facilities

Among all facilities, the percentages of facilities having assigned health care providers, by facility type, Bangladesh HFS 2022

| _ | Facility type | | | | | |
|---|---------------|------|------|-------|--------|--------------------------|
| Health care providers | DH | MCWC | UHC | UHFWC | USC/RD | Community clinic (CC) |
| Senior/junior consultant (medicine) | 87.1 | - | 32.2 | - | - | - |
| Senior/junior consultant (surgery) | 79.0 | - | 16.5 | - | - | - |
| Senior/junior consultant (obs & gyn) | 91.9 | - | 49.0 | - | - | - |
| Senior/junior consultant (pediatric) | 88.7 | - | 30.5 | - | - | - |
| Senior/junior consultant (anesthesia) | 96.8 | - | 64.8 | - | - | - |
| Radiologist | 43.6 | - | 4.9 | - | - | - |
| Medical technologist (Radiology) | 82.3 | - | 52.4 | - | - | - |
| Pathologist | 53.2 | - | 18.6 | - | - | - |
| Medical technologist (Laboratory) | 96.8 | - | 83.0 | - | - | - |
| Midwife | - | - | 83.8 | - | - | - |
| Community health care provider (CHCP) | - | - | - | - | - | 96.4 |
| Sub assistant community medical officer (SACMO) | - | - | 89.6 | 62.5 | 79.2 | - |
| Family welfare visitor (FWV) | - | 90.0 | 75.5 | 80.4 | - | - |
| Number of facilities (weighted) | 4 | 7 | 30 | 225 | 85 | 994 |
| Number of facilities (unweighted) | 62 | 100 | 157 | 293 | 141 | 488 |

Note: Numbers were provided by the facility in-charge. "-" Means the post is not applicable

Table 3.11.2: Staffing pattern in surveyed facilities (Medical officer and Nurse):

Among all DHs and UHCs, the descriptive statistics (mean and range) of the number of medical officers and nurses, Bangladesh HFS 2022

| | Facility Type | | | | | |
|--|---------------|-------------|-------------------|------------------|--|--|
| | DH UHC DH UHC | | | | | |
| Health care providers | (Mean) | (Mean) | (Range) | (Range) | | |
| Medical Officer ¹ Nurse ² | 21.3 126.1 | 9.8 25.0 | (4-52) (5-398) | (1-22) (1-37) | | |

Note: Numbers were provided by the facility in-charge. ¹ Residential medical officer (RMO), medical officer (MO)/physician, anesthetist, emergency medical officer (EMO), indoor medical officer (IMO), indoor medical officer (IMO) cardiology, indoor medical officer (IMO) surgery, indoor medical officer (IMO) orthopedic surgery, indoor medical officer (IMO) medicine, indoor medical officer (IMO) and afficer indoor medical officer (IMO) medicine, indoor medical officer (IMO) production indoor medical officer (IMO) medicine, indoor medical officer (IMO) pediatrics, indoor medical officer (IMO) OBS/GYN, medical officer (MO-blood transfusion), dental surgeon, assistant surgeon/equivalent, assistant registrar (medicine), assistant registrar (surgery), assistant registrar (OBS/GYN), assistant registrar (pediatric), medical officer (MCH-FP)

² Matron, nursing supervisor, senior staff nurse, nurse midwife, staff nurse, assistant nursing attendant

Table 3.12.1: Sharps waste management

Among all facilities, the percentages that reported having different methods of sharps waste management, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Incinerator | Open burning | Dump without burning | Remove offsite | Number of facilities (weighted) | Number of facilities (unweighted) |
|-------------------------------|-------------|--------------|-------------------------|-------------------|---------------------------------------|---|
| Facility type | | | | | | |
| District and upazila public | | | | | | |
| facilities | 2.3 | 24.7 | 10.2 | 22.7 | 41 | 319 |
| DH | 4.8 | 9.7 | 11.3 | 29.0 | 4 | 62 |
| MCWC | 1.0 | 18.1 | 6.4 | 35.9 | 7 | 100 |
| UHC | 2.2 | 28.6 | 11.0 | 18.6 | 30 | 157 |
| Union-level public facilities | 0.5 | 17.7 | 9.9 | 21.2 | 310 | 434 |
| UHFWC | 0.7 | 20.8 | 10.7 | 21.5 | 225 | 293 |
| USC/RD | 0.0 | 9.7 | 7.8 | 20.2 | 85 | 141 |
| Community clinic (CC) | 0.0 | 7.8 | 7.0 | 41.5 | 994 | 488 |
| NGO clinic/hospital | 1.8 | 4.2 | 8.0 | 62.6 | 29 | 127 |
| Private hospital | 0.4 | 3.2 | 4.0 | 65.4 | 183 | 189 |
| Location | | | | | | |
| Urban | 0.9 | 5.4 | 4.5 | 63.8 | 235 | 514 |
| Rural | 0.1 | 10.3 | 7.8 | 36.0 | 1,322 | 1,043 |
| Division | | | | | | |
| Barishal | 0.0 | 7.5 | 5.7 | 41.2 | 111 | 152 |
| Chattogram | 0.0 | 6.4 | 6.1 | 46.1 | 291 | 244 |
| Dhaka | 0.3 | 14.5 | 6.0 | 40.7 | 317 | 264 |
| Khulna | 0.2 | 9.6 | 7.3 | 47.1 | 210 | 208 |
| Rajshahi | 0.5 | 8.1 | 9.2 | 30.1 | 214 | 212 |
| Rangpur | 0.3 | 8.3 | 8.6 | 42.0 | 193 | 190 |
| Sylnet | 0.0 | 15.4 | 12.1 | 25.1 | 99 | 136 |
| Mymensingh | 0.5 | 6.1 | 5.9 | 38.8 | 122 | 151 |
| Total | 0.2 | 9.6 | 7.3 | 40.2 | 1,557 | 1,557 |
| Total excluding CCs | 0.7 | 12.8 | 7.9 | 37.8 | 563 | 1,069 |
| Total public | 0.2 | 10.6 | 7.8 | 36.3 | 1,345 | 1,241 |
| Total public excluding CCs | 0.7 | 18.6 | 9.9 | 21.3 | 351 | 753 |

Table 3.12.2: Non-sharps waste management

Among all facilities, the percentages that reported having different methods of medical waste management, by background characteristics, Bangladesh HFS 2022

| Background | | | Dump without | | Number of facilities | Number of facilities |
|-------------------------------|-------------|--------------|--------------|----------------|-------------------------|-------------------------|
| characteristic | Incinerator | Open burning | burning | Remove offsite | (weighted) | (unweighted) |
| Facility type | | | | | | |
| District and upazila public | | | | | | |
| facilities | 1.5 | 24.7 | 10.3 | 19.1 | 41 | 319 |
| DH | 1.6 | 11.3 | 14.5 | 30.6 | 4 | 62 |
| MCWC | 1.0 | 17.1 | 4.9 | 37.1 | / | 100 |
| UHC | 1.6 | 28.5 | 10.9 | 13.1 | 30 | 157 |
| Union-level public facilities | 1.4 | 16.4 | 10.5 | 13.1 | 310 | 434 |
| UHFWC | 1.3 | 17.4 | 11.1 | 14.5 | 225 | 293 |
| USC/RD | 1.6 | 13.9 | 9.1 | 9.4 | 85 | 141 |
| Community clinic (CC) | 0.2 | 11.6 | 7.4 | 21.0 | 994 | 488 |
| NGO clinic/hospital | 5.6 | 5.2 | 8.4 | 59.5 | 29 | 127 |
| Private hospital | 0.4 | 1.6 | 3.6 | 70.7 | 183 | 189 |
| Location | | | | | | |
| Urban | 1.2 | 4.4 | 3.7 | 66.3 | 235 | 514 |
| Rural | 0.5 | 12.9 | 8.4 | 18.8 | 1,322 | 1,043 |
| Division | | | | | | |
| Barishal | 0.0 | 5.6 | 8.3 | 23.1 | 111 | 152 |
| Chattogram | 0.4 | 9.3 | 10.6 | 23.9 | 291 | 244 |
| Dhaka | 0.4 | 10.8 | 4.8 | 34.3 | 317 | 264 |
| Khulna | 0.6 | 10.8 | 9.5 | 27.6 | 210 | 208 |
| Rajsnani | 1.0 | 13.0 | 9.6 | 21.7 | 214 | 212 |
| Sylbot | 1.3 | 15.7 | 0.Z | 27.4 | 193 | 190 |
| Mymensingh | 0.5 | 20.0 | 24 | 18.2 | 122 | 150 |
| Wymenangn | 0.0 | 5.7 | 2.4 | 10.2 | 122 | 101 |
| Total | 0.6 | 11.6 | 7.7 | 25.9 | 1,557 | 1,557 |
| Total excluding CCs | 1.3 | 11.6 | 8.2 | 34.6 | 563 | 1,069 |
| Total public | 0.5 | 13.1 | 8.2 | 19.1 | 1,345 | 1,241 |
| Total public excluding CCs | 1.4 | 17.4 | 10.5 | 13.8 | 351 | 753 |

Table 3.13: Key indicators for general service preparedness in MCWCs

Among all facilities, the percentages with key indicators, Bangladesh HFS 2022

| | Facility type | | |
|---|---|---|--|
| Indicators | MCWC (district-level) | MCWC (upazila and union-level) | |
| Basic client services | · · · · · · · · · · · · · · · · · · · | | |
| Child curative care Child growth monitoring Child vaccination Any modern methods of family planning Antenatal care services Normal delivery | 98.3 72.3 66.1 96.8 100.0 98.5 | 89.8 61.1 42.2 92.2 97.2 68.3 | |
| Basic Amenities Regular electricity ¹ Improved water source ² Visual and auditory privacy ³ Client latrine ⁴ Communication equipment ⁵ Computer with Internet ⁶ | 95.2 98.3 58.7 95.3 87.0 98.5 | 68.8 100.0 53.6 95.1 33.7 41.9 | |
| Basic equipment Adult scale Child scale ⁷ or infant scale ⁸ Thermometer Stethoscope Blood pressure apparatus ⁹ Light source ¹⁰ | 93.6 77.6 86.9 98.4 98.4 90.5 | 78.8 74.1 70.7 94.8 94.8 92.1 | |
| Quality assurance | 32.0 | 5.0 | |
| Human resource ¹¹ | | | |
| Medical officer (clinic) Family welfare visitor (FWV) | 79.1 96.8 | 12.8 78.7 | |
| Number of facilities (weighted) | 5 | 3 | |
| Number of facilities (unweighted) | 62 | 38 | |

¹ Facility is connected to a central power grid and there was no interruption in power supply lasting for more than 2 hours at a time during normal working hours in the 7 days before the survey, or facility has a functioning generator on the day of the survey, or else facility has back-up solar power. ² Water is piped into facility or piped onto facility grounds, bottled water is available, or water is available from a public tap

or standpipe, a tube well or borehole, a protected dug well, a protected spring, or rainwater and the outlet from this source is within 500 meters of the facility.

³ A private room or screened-off space available in the general outpatient service area that is a sufficient distance from other clients so that a normal conversation could be held without the client being seen or heard by others

⁴ Facility has a functioning flush or pour-flush toilet, a ventilated improved pit latrine, or a composting toilet.

⁵ Facility has a functioning land-line telephone, a functioning facility-owned cellular phone, or a private cellular phone that is supported by the facility.

⁶ Facility has a functioning computer with access to the Internet, or facility has access to the Internet via a cellular phone inside the facility.

⁷ A scale with a gradation of 250 grams, or a digital standing scale with a gradation of 250 grams or lower where an adult can hold a child to be weighed, available somewhere in the general outpatient area

⁸ A scale with a gradation of 100 grams, or a digital standing scale with a gradation of 100 grams where an adult can hold an infant to be weighed, available somewhere in the general outpatient area

⁹ A digital blood pressure machine or a manual sphygmomanometer with a stethoscope available somewhere in the general outpatient area ¹⁰ ¹⁰ A spotlight source that can be used for client examination or a functioning flashlight available somewhere in the general

outpatient area

¹¹percentages of MCWCs having the assigned health care providers

Key Findings

- Almost all public facilities offer outpatient curative care for sick children (95%). Most public facilities offer child vaccination (93%) and growth monitoring services (88%). (Table 4.1).
- Outpatient curative care for sick children is universally available in DHs and UHCs (both 100%), whereas it is available in about nine in ten NGO facilities and in more than half (57%) of the private facilities (Table 4.1 and Figure 4.1).
- Availability of growth monitoring services is the highest in DHs and UHCs (both 98%) and lowest in private facilities (23%). (**Table 4.1** and **Figure 4.1**).
- Availability of child vaccination service is universal in DHs (95%), UHCs (99%), and CCs (97%) (Table 4.1 and Figure 4.1).
- Of the facilities that offer child curative care, half of public facilities (51%) have at least one staff member trained in IMCI, while less than one-fifth (18%) have IMCI guidelines available on site. Since 2017, the percentage of public health facilities having IMCI guidelines has decreased from 42% to 18% (Table 4.6 and Figures 4.2–4.3).
- Around half (46%) of the public health facilities offering outpatient child curative care have a child weighing scale, while most have a stethoscope (92%) available on-site. Thermometers (78%), length or height boards (74%), and growth charts (66%) are observed in the majority of public facilities (Table 4.2 and Figures 4.4–4.6).
- The availability of essential medicines in public health facilities that offer curative care services for children, such as oral rehydration solution (ORS), amoxicillin, and paracetamol, has increased in the last five years. However, the availability of zinc tablets or syrup has slightly decreased in public health facilities (Table 4.4 and Figures 4.7–4.10).
- Almost all public health facilities offering child curative care have ORS (84%), amoxicillin (89%), paracetamol (92%), and albendazole/mebendazole (92%) available on the day of the survey (Table 4.4 and Figures 4.10–4.12).
- The average readiness score (out of 10 items) for facilities providing outpatient child curative care is higher among DHs (6.9), UHCs (7.9), and CCs (7.2). However, the scores are considerably lower in MCWCs (5.6), USC/RDs (4.7), and private facilities (4.3) (Table 4.6 and Figure 4.14).

4.1 BACKGROUND

ach year, millions of under-five children are dying due to preventable causes. It is common for healthcare providers to address a sick child's most evident symptoms without providing a complete assessment of the child's overall health status. For this reason, the World Health Organization (WHO) and other collaborative agencies developed the integrated management of childhood illness (IMCI) strategy (WHO 1998). This strategy advocates using every visit to a health care provider as an opportunity not only to conduct a full assessment of the child's current health and possible underlying cause of illness but also to provide interventions, such as vaccinations, that can prevent illness or minimize its progression. Bangladesh has adopted the IMCI strategy, and the country's 4th Health, Population and Nutrition Sector Program (HPNSP) aims to strengthen the existing IMCI program (Government of the People's Republic of Bangladesh, 2017).

The IMCI strategy aims to reduce morbidity and mortality among children under age five through the following three activities:

- Improving health workers' skills through training and supportive supervision;
- Improving health systems, including equipment, supplies, organization of work, and referral systems; and
- Improving child care at the community and household levels in line with key family practices.

The 2022 BHFS examined the availability of necessary services for assessing sick children for main symptoms (fever, cough, or difficulty breathing; diarrhoea; ear pain or discharge), nutrition and immunization status, feeding problems, and other potential problems and danger signs that indicate the need for immediate referral or hospital admission. The survey also assessed service readiness in facilities where services were available. IMCI guidelines were used as the basis for this assessment.

This chapter explores the following issues related to the provision of quality child health services at health facilities:

- Availability of child health services: Section 4.2, including Tables 4.1–4.5 and Figures 4.1–4.13, assesses the availability of child health services and of trained staff, equipment, guidelines, medicines, infection control items, and laboratory diagnostic capacity necessary to provide those services.
- Service readiness: Section 4.3, including Table 4.6 and Figures 4.14, addresses the overall readiness of facilities to provide quality child health services.
- Basic management and administrative systems: Section 4.4, including Tables 4.7–4.8 and Figures 4.15–4.16, summarizes two aspects of management and administrative systems that support quality services: personal supervision and in-service training for providers of child health services.

4.2 AVAILABILITY OF CHILD HEALTH SERVICES

4.2.1 Provision of Outpatient Curative Care, Child Growth Monitoring, Child Vaccination, and Nutrition Services

The 2022 BHFS assessed the availability of child health services including the availability of curative care services for sick children, child growth monitoring, and child vaccination services.

- Outpatient curative care for sick children is universally available (95%) in public health facilities. Most of the public facilities provide growth monitoring (88%) services, while more than 9 of 10 public facilities (93%) provide child vaccination services (**Table 4.1**).
- Child curative care services are universally available in DHs (100%), UHCs (100%), UHFWCs (94%), USC/RDs (98%), MCWCs (95%), and CCs (95%); whereas only 57% of the private facilities provide these services (Table 4.1 and Figure 4.1).
- Growth monitoring services are available in most DHs (98%), UHCs (98%), and CCs (92%). However, only about one-fourth (23%) of the private facilities provide these services (**Table 4.1** and **Figure 4.1**).
- More than 9 in 10 DHs (95%), UHCs (99%), and CCs (97%) provide child vaccination services. However, only 3% of the private facilities provide child vaccination services, which indicates a sharp decline from 57% in 2014 and 23% in 2017 (Table 4.1 and Figure 4.1).
- Most public facilities (88%) provide the service of diagnosis and/or treatment for child nutrition (Table 4.1).



Figure 4.1 Availability of child health services in 2022, by facility type (N=1,557)

4.2.2 Availability of Guidelines, Trained Staff, and Basic Equipment for Child Health Care

Facilities providing curative care for sick children should have basic equipment, necessary medicines, and trained staff to provide quality services. The 2022 BHFS addressed the availability of these logistics.

- Twenty-one percent of the public facilities (excluding CCs) that offer outpatient curative care for sick children have guidelines for IMCI available on the day of the survey. The availability of IMCI guidelines is highest in UHCs (51%) and lowest in private facilities (5%) (**Table 4.2** and **Figure 4.2**).
- Nearly half of DHs (48%) and UHCs (51%) had IMCI guidelines available on the day of survey (Table 4.2 and Figure 4.2).

• At the divisional level, Khulna has the lowest availability of IMCI guidelines (10%), while Barishal has the highest percentage (26%) (**Table 4.2**).





Percent

- Half of public facilities (51%) that offer outpatient curative care for sick children have at least one staff member trained at any time in IMCI (**Table 4.3** and **Figure 4.3**).
- The majority of DHs (84%) and UHCs (78%) have at least one provider ever trained in IMCI, while around half of the CCs (52%) and NGOs (53%) offering child curative care have at least one provider ever-trained in IMCI (**Table 4.3** and **Figure 4.3**).
- The percentage of having at least one IMCI ever-trained staff member has decreased gradually in almost all type of facilities except UHFWCs, CCs and NGOs from 2014 and 2017 (**Figure 4.3**).
- The proportion of having at least one IMCI ever-trained staff member is higher in rural regions (51%) compared to urban areas (32%) (**Table 4.3**).
- Among the divisions, 35% of facilities that offer curative care for sick children in Sylhet have at least one ever-trained IMCI provider, compared to 65% in Barishal (**Table 4.3**).




Availability of Equipment for Outpatient Child Curative Care Services

The availability of all necessary equipment for outpatient child curative care service ensures high-quality child curative care. The equipment includes child weighing scale, length or height board, thermometer, stethoscope, infant weighing scale, growth monitoring chart, MUAC tape, timer, and pulse oximeter.

- More than half (53%) of the public facilities (excluding CCs) had a functioning child weighing scale available on the day of the visit (Table 4.2 and Figure 4.4).
- The availability of child weighing scales is relatively higher in DHs (82%), UHCs (79%), NGOs (66%), and private facilities (70%) (**Table 4.2** and **Figure 4.4**).
- Since 2017, the availability of child weighing scales have increased in DHs, UHCs, MCWCs, and UHFWCs (Figure 4.4).





- Almost 8 in 10 public facilities (78%) have thermometers available for outpatient curative care for sick children in 2022 (Table 4.2 and Figure 4.5).
- In 2022, the availability of thermometers is nearly universal in DHs (90%), UHCs (94%), NGOs (99%), and private facilities (95%) (**Table 4.2** and **Figure 4.5**).
- The availability of thermometers has gradually decreased in almost all facility types. However, it slightly increased in UHFWCs (76% in 2017 to 79% in 2022) and remained unchanged in NGO facilities (**Figure 4.5**).



Figure 4.5 Availability of functioning thermometer for outpatient child curative care, by facility type

- Two-thirds of the public facilities (66%) offering curative care for sick children had growth monitoring charts available on the day of survey (**Table 4.2, 4.6**, and **Figure 4.6**).
- DHs (65%), UHCs (75%), CCs (71%), and NGOs (63%) are most likely to have growth monitoring charts and private facilities are least likely (11%) on the day of the survey (Table 4.2, 4.6, and Figure 4.6).
- Nearly all facilities experienced a reduction in the availability of growth monitoring charts, except for UHFWCs (54% in 2017 to 56% in 2022) and CCs (71%), where the situation remained unchanged (Figure 4.6).



Figure 4.6 Availability of growth chart in health facilities for outpatient child curative care, by facility type

Percent

- Three in four public facilities (74%) offering curative care for sick children have a length or height board available in 2022. The availability of length or height boards is higher in DHs (90%), UHCs (94%), and NGOs (82%) and lowest in private facilities (40%) in 2022 (Table 4.2).
- Stethoscope is a basic and necessary equipment in providing child curative care services. The availability of a stethoscope is universal (92%) in public health facilities (**Table 4.2**).
- For providing outpatient curative care for sick children, more than half (54%) of public facilities have pulse oximeter. Moreover, over three-quarters of private facilities (77%) have pulse oximeters available on site in 2022 (**Table 4.2**).

The availability of equipment for outpatient curative care for sick children is evaluated by the average equipment score. **Figure 4.7** presents the average score constituted by the mean number of equipment items out of a total of four equipment (child weighing scale, length or height board, thermometer, and stethoscope), that are available on-site on the day of the visit.

- The average equipment score is 2.9 in public facilities (excluding CCs) offering child curative care services, indicating an average presence of three items in the child curative care service corner (Table 4.2 and Figure 4.7).
- The average readiness score of equipment is higher among DHs (3.6), UHCs (3.7), and NGOs (3.5) and lowest among CCs (2.9) and USC/RDs (2.5) (Figure 4.7).
- There has been a decline observed in the average readiness score of equipment among the public facilities, from 3.1 in 2017 to 2.9 2022 (**Figure 4.7**).



Figure 4.7 Average score of equipment (child weighing scale, length or height board, thermometer, and stethoscope) for child curative care, by facility type

Availability of Equipment for Inpatient Child Curative Care Services

Inpatient curative care is crucial for critically ill children who require continuous monitoring and intensive medical attention by healthcare providers. In Bangladesh, all public facilities at district and upazila level, as well as most private and NGO facilities, operate inpatient department to serve critically ill patients. On the other hand, union-level and public facilities and community clinics are designed only outpatient services. The BHFS 2022 assessed the availability of necessary logistics and equipment for treating sick children in the indoor department.

Table 4.2.1 provides information on the availability of equipment (child weighing scale, infant scale, thermometer, stethoscope, nebulizer, timer, torch, growth chart, height board, pulse oximeter, nasal catheter/cannula, and oxygen) at inpatient child curative care service site, among facilities that offer inpatient curative care for sick children.

- The availability of oxygen (96%) and nebulizer (93%) are universal in public facilities offering inpatient curative care for sick children in 2022. The availability of oxygen in NGO facilities (32%) is remarkably lower than any other facilities that offer inpatient child curative care (**Table 4.2.1**).
- Most public facilities (59%) and private hospitals (69%) have a nasal cannula or catheter available on the day of the survey; it is observed in only 45% of the NGO clinics (**Table 4.2.1**).
- Eight of ten public facilities have pulse oximeters for inpatient child curative care. Pulse oximeters are universally available in private hospitals (90%) (**Table 4.2.1**). Although most of the DHs (83%) and UHCs (81%) have pulse oximeters in child inpatient care departments, it is not available in 41% of the MCWCs (**Table 4.2.1**).

• Stethoscopes (98%), thermometers (96%), and timers for assessing respiratory rate (92%) are universally available in public facilities offering inpatient curative care for children (**Table 4.2.1**).

4.2.3 Availability of Medicines and Commodities for Child Health Care

A range of medicines and commodities are essential to provide curative care for sick children. Both the WHO and USAID have proposed a set of essential and priority medicines and commodities that facilities should have available to support the provision of health services for sick children (WHO 2013). Overall, essential medicines are in better supply than priority medicines in Bangladesh's health facilities that offer curative care for sick children.

- Among facilities offering outpatient curative care for sick children, the availability of ORS is universal among DHs (98%), UHCs (99%), and CCs (99%). However, it is comparatively low in private facilities (65%), MCWCs (25%), and UHFWCs (19%) (Table 4.4 and Figure 4.8).
- Between 2017 and 2022, the availability of ORS increased from 80% to 84% among public health facilities (Figure 4.8).

Figure 4.8 Availability of ORS in health facilities for outpatient child curative care, by facility type



- Two-thirds of the public facilities (65%) offering outpatient curative care for sick children have zinc tablets or syrup available on the day of the survey in 2022 (Table 4.4 and Figure 4.9).
- Most UHCs (82%), CCs (81%), and DHs (74%) have zinc tables or syrup available on the day of the survey; however, the proportion is lower in UHFWCs (7%) and MCWCs (6%) (Table 4.4 and Figure 4.9).
- The availability of zinc tablets or syrup has increased notably in DHs and UHCs, from 2017 to 2022 (**Figure 4.9**).



Figure 4.9 Availability of zinc tablets or syrup in health facilities for outpatient child curative care, by facility type

Percent

- Around 9 in 10 public facilities (89%) offering outpatient curative care services for sick children have amoxicillin syrup, suspension, or dispersible pediatric-dosed tablets available on the day of the survey in 2022 (**Table 4.4** and **Figure 4.10**).
- The availability of oral amoxicillin for children is highest among CCs (93%), which has increased substantially from 80% in 2017. However, the availability is relatively low in DHs (32%), NGO facilities (30%), and private facilities (24%) (**Table 4.4** and **Figure 4.10**).



Figure 4.10 Availability of amoxicillin in health facilities for outpatient child curative care, by facility type

- In 2022, almost all (92%) of the public health facilities have paracetamol syrup or suspension available for outpatient child curative care, an increase from 84% in 2017 (**Table 4.4** and **Figure 4.11**).
- The availability of paracetamol for child curative care is more than 80% percent across all types of public health facilities, except USC/RDs (64%).
- Nearly two-thirds of the NGO facilities (63%) and more than half of the private facilities (53%) have paracetamol syrup or suspension available on site (**Table 4.4** and **Figure 4.11**).



Figure 4.11 Availability of paracetamol syrup or suspension in health facilities for outpatient child curative care, by facility type

- Almost all public health facilities offering outpatient curative care for sick children (92%) have mebendazole or albendazole tablets available on the day of the survey in 2022 (Table 4.4 and Figure 4.12).
- On the other hand, only three in ten DHs (31%) and nearly half of the private (49%) and NGO (51%) facilities have mebendazole/albendazole available, decreasing markedly compared to 2017 (**Table 4.4** and **Figure 4.12**).
- The availability of ceftriaxone powder for injection (75%) and gentamicin injection (69%) is comparatively high in district and upazila-level facilities (**Table 4.4**)



Figure 4.12 Availability of mebendazole/albendazole in health facilities for outpatient child curative care, by facility type

4.2.4 Infection Control in Sick Child Services

Infection control is one of the most essential health service components. **Table 4.5** summarizes information related to the availability of necessary infection control items (soap, running water, alcohol-based disinfectant, latex gloves, sharps containers, and waste receptacles).

- Over half of public facilities offering child curative care services have soap (54%) and running water (52%); on the other hand, about 8 of 10 private and NGO facilities are equipped with soap and running water (**Table 4.5**).
- The majority of public facilities have alcohol-based hand disinfectant (67%) and latex gloves (60%) available on site (**Table 4.5**).
- Public facilities have an average of 3.4 items for infection control available at outpatient child curative care service site in 2022, which has increased gradually since 2014 (Table 4.5 and Figure 4.13).
- DHs, MCWCs, NGOs, and private facilities scored higher in the average score for infection control. Nevertheless, there has been a decrease in scores for DHs, UHCs, NGOs, and private facilities from 2017 to 2022 (Table 4.5 and Figure 4.13).



Figure 4.13 Average score of infection control items (out of 6 items) for child curative care, by facility type

Average

4.2.5 Laboratory Diagnostic Capacity

The 2022 BFHS gathered information on two laboratory tests that are particularly important in diagnosing conditions among children: hemoglobin testing and stool microscopy.

- Among facilities offering outpatient child curative care for sick children, overall, hemoglobin tests (31%) and stool microscopy (5%) tests are available in a lower proportion of public facilities (excluding CCs). However, union-level public facilities are not designed to conduct stool microscopy (Table 4.5).
- Hemoglobin tests are available in almost all of the DHs (94%), UHCs (92%), NGOs (89%), and private facilities (88%) offering outpatient sick child services (Table 4.5).
- Stool microscopy is most available in DHs (61%) and least available in NGOs (17%) offering outpatient sick child services (**Table 4.5**).

4.3 READINESS OF HEALTH FACILITIES TO PROVIDE CHILD CURATIVE CARE

In assessing the overall readiness of health facilities in Bangladesh to provide child curative care, this report used ten tracer indicators from the WHO Service Availability and Readiness Assessment (SARA) (WHO 2013):

- IMCI guidelines:
 - National or other guidelines on IMCI are available at the facility
- IMCI trained staff:
 - At least one provider who ever received in-service training on at least some components of IMCI
- Equipment:

- Child weighing scale
- Thermometer
- Growth chart
- Medicines:
 - ORS
 - Zinc tablets/syrup
 - Amoxicillin syrup/suspension/dispersible tablet
 - Paracetamol syrup/suspension
 - Mebendazole/albendazole

The readiness of facilities to deliver outpatient curative care for sick children is evaluated by the average readiness score. This score is determined by calculating the mean number of items, out of a total of 10 items, that are available on the day of the visit. In essence, it provides insight into the overall capability of the facilities to provide appropriate outpatient care for sick children.

- Among facilities offering sick child services, the highest readiness scores are among DHs (6.9), UHCs (7.9), and CCs (7.2), with an average of nearly seven or more among all ten necessary items to provide outpatient child curative care. Private facilities are less prepared compared to the other facilities, having an average 4.3 items out of ten (**Table 4.6** and **Figure 4.14**).
- The average readiness score has declined in most types of facilities between 2017 and 2022. However, the readiness scores at UHCs has increased and UHFWCs remained unchanged during that time (**Figure 4.14**).

Figure 4.14 Average readiness score of health facilities (out of 10 items) to provide child curative care, by facility type



Average

4.4 BASIC MANAGEMENT AND ADMINISTRATIVE SYSTEM

4.4.1 Personal Supervision and Training

Training and supervision are essential management functions in building capacity and ensuring the quality of services. Periodic in-service training keeps providers up-to-date and helps to refresh their knowledge and skills. Regular supervision is also an important source of support and direction. **Table 4.7** provides information on personal supervision and on interviewed providers of child health services who reported receiving in-service training related to child curative care in the 24 months before the assessment.

- Training related to child health within the 24 months before the survey has declined from one-third (36%) of providers in public facilities in the 2017 BHFS to one-fourth (25%) of providers in the 2022 BHFS (**Table 4.7** and **Figure 4.15**).
- Among child health service providers, about one in three providers at CCs (33%) and NGOs (32%) received in-service training on child health services in the 24 months preceding the survey, which is higher than any other facility type (**Table 4.7** and **Figure 4.15**).

Figure 4.15 Provider training on topics related to child health in the last 24 months

Percent



- Almost all child health service providers in public facilities (93%) have received personal supervision during the six months preceding the survey (Table 4.7 and Figure 4.16).
- The proportion of providers who received personal supervision in the 6 months before the survey is universal in DHs (97%), MCWCs (98%), UHCs (98%), CCs (92%), and NGOs (96%) (**Table 4.7** and **Figure 4.16**).



Figure 4.16 Personal supervision of child health service providers in the six months before the survey

- Around one-quarter of the child health service providers in public facilities (24%) have received both child health related training in the 24 months before the survey and personal supervision in the six months before the survey (**Table 4.7**).
- The percentage of providers who received both training during the last 24 months and personal supervision during the last six months is the highest in NGOs (32%) and lowest in MCWCs (8%) and private hospitals (6%) in 2022 (**Table 4.7**).
- Only 14% of all interviewed child health service providers in Barishal have received both training and personal supervision, which is notably lower than Chattogram and Mymensingh divisions (both 28%) (Table 4.7).

4.4.2 In-Service Training by Topic

Providers of child health services were asked about in-service training on specific topics related to child health. **Table 4.8** presents the percentage of providers who reported receiving in-service training on EPI/cold chain monitoring, IMCI, acute respiratory infection (ARI), and diarrhoea diagnosis or treatment, among interviewed providers who provide child health services.

- Nearly three of ten providers in public facilities reported that they have received in-service training (at any time) on EPI/cold chain monitoring (30%), IMCI (37%), and ARI (29%). However, only a small proportion of them reported receiving training in the two years before the survey on EPI/cold chain monitoring (6%), IMCI (7%), and ARI (6%) (Table 4.8 and Figure 4.17).
- Nearly six in ten child health service providers at CCs (59%) reported that they have received in-service training on diarrhea diagnosis and treatment at any time, which is remarkably higher than providers from DHs (19%) and UHCs (25%) (Table 4.8).

• The percentages of providers in public facilities reported that they received necessary training on child health in the last 24 months preceding the survey, including training on EPI cold chain, IMCI, ARI, and Diarrhoea diagnosis or treatment, have declined since 2017 (**Figure 4.17**).

Figure 4.17 Provider training on topics related to child health in public health facilities during the last 24 months

Percent



CHILD HEALTH SERVICES

LIST OF TABLES

- Table 4.1 Availability of child health services
- Table 4.2
 Guidelines and equipment for outpatient child curative care services
- Table 4.2.1 Equipment for inpatient child curative care services
- Table 4.3 Trained staff for child curative care services
- Table 4.4 Availability of essential and priority medicines
- Table 4.5 Infection control and laboratory diagnostic capacity
- Table 4.6 Readiness of health facilities to provide child curative care services
- Table 4.7 Supportive management for providers of child health services
- Table 4.8
 Training for child health service providers
- Table 4.9 Key indicators for child curative care services in MCWCs

Table 4.1: Availability of child health services

Among all facilities, the percentages offering specific child health services at the facility, by background characteristics Bangladesh HFS 2022

| | Percentage of facilities that offer: | | | | | | | | | |
|---|---|----------------------|-----------------------------------|---|--|--|--|---------------------------|---------------------------------------|---|
| Background characteristic | Outpatient curative care for sick children | Growth | Child vaccination ¹ | Diagnosis of and/or treatment for child nutrition | All 3 basic child health services ² | All 4 basic child health services ³ | Routine vitamin A supplemen- tation | Deworming for children | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type | | | | | | | | | | |
| District and upazila public facilities | 99.1 | 92.9 | 90.9 | 97.1 | 86.7 | 85.9 | 77.5 | 93.0 | 41 | 319 |
| DH MCWC UHC | 100.0 95.1 100.0 | 98.4 68.1 98.1 | 95.2 57.0 98.5 | 100.0 87.1 99.1 | 93.5 42.0 96.6 | 93.5 40.9 95.7 | 83.9 40.5 85.6 | 87.1 90.2 94.6 | 4 7 30 | 62 100 157 |
| Union-level | 95.3 | 74.2 | 79.1 | 86.4 | 59.1 | 55.2 | 28.0 | 81.5 | 310 | 434 |
| UHFWC USC/RD | 94.2 98.1 | 78.2 63.6 | 79.2 79.1 | 86.1 87.1 | 61.6 52.5 | 57.3 49.8 | 24.5 37.4 | 81.6 81.3 | 225 85 | 293 141 |
| Community clinic (CC) | 94.5 | 92.4 | 97.1 | 88.5 | 85.2 | 80.2 | 76.3 | 79.3 | 994 | 488 |
| NGO clinic/hospital Private hospital | 88.4 56.6 | 81.9 22.5 | 71.9 3.0 | 78.6 50.1 | 63.9 1.9 | 60.3 1.9 | 66.3 32.0 | 76.8 42.0 | 29 183 | 127 189 |
| Location | | | | | | | | | | |
| Urban Rural | 65.7 94.6 | 36.9 88.1 | 22.4 92.7 | 59.7 87.8 | 18.4 79.0 | 18.0 74.3 | 41.8 64.8 | 52.3 79.8 | 235 1,322 | 514 1,043 |
| Division | | | | | | | | | | |
| Barishal Chattogram Dhaka | 92.7 97.4 83.8 | 87.6 87.5 75.7 | 84.5 85.1 80.0 | 85.8 92.7 78.0 | 73.9 81.0 62.1 | 69.4 78.5 58.3 | 68.0 66.4 58.8 | 72.7 84.2 68.3 | 111 291 317 | 152 244 264 |
| Khulna Rajshahi Rangpur | 76.7 95.1 95.7 | 76.2 73.9 84 7 | 70.9 82.5 87.3 | 72.2 86.5 88.7 | 60.5 65.3 78.0 | 59.5 61.0 73.8 | 54.7 69.8 61 9 | 64.7 83.1 82.4 | 210 214 193 | 208 212 190 |
| Sylhet Mymensingh | 88.4 94.7 | 82.7 78.8 | 84.0 87.2 | 79.8 83.7 | 71.8 69.1 | 65.0 59.4 | 54.2 50.6 | 70.8 76.4 | 99 122 | 136 151 |
| Total | 90.2 | 80.4 | 82.1 | 83.6 | 69.8 | 65.8 | 61.3 | 75.7 | 1,557 | 1,557 |
| Total excluding CCs | 82.7 | 59.1 | 54.9 | 75.0 | 42.8 | 40.4 | 34.9 | 69.3 | 563 | 1,069 |
| Total public | 94.8 | 88.2 | 93.1 | 88.2 | 79.2 | 74.6 | 65.2 | 80.2 | 1,345 | 1,241 |
| Total public excluding CCs | 95.7 | 76.4 | 80.5 | 87.6 | 62.3 | 58.9 | 33.8 | 82.9 | 351 | 753 |

¹ Facility routinely provides DPT/pentavalent, polio, and measles vaccinations to children.
 ² Outpatient curative care for sick children, growth monitoring, and child vaccination.
 ³ Outpatient curative care for sick children, growth monitoring, child vaccination, and diagnosis of and/or treatment for child nutrition.

Table 4.2: Guidelines and equipment for outpatient child curative care services

Among facilities that offer outpatient curative care for sick children, the percentages having indicated guidelines and equipment, by background characteristics, Bangladesh HFS 2022

| | Gui | delines | Equipment | | | | | | | | | | Number of facilities offering | Number of facilities | |
|---|----------------------|----------------------|---|------------------------------|----------------------|------------------------|--|--|----------------------|----------------------|----------------------|----------------------|--|---|--|
| Background characteristic | IMCI | Growth monitoring | Child weighing scale ¹ | Length or height board | Thermo- meter | Stetho- scope | Average of four equipment items available ² | Infant weighing scale ³ | Growth chart | MUAC tape | Timer ⁴ | Pulse oximeter | outpatient curative care for sick children (weighted) | ottering outpatient curative care for sick children (unweighted) | |
| Facility type | | | | | | | | | | | | | | | |
| District and upazila public facilities | 43.4 | 25.9 | 76.7 | 90.2 | 92.4 | 99.8 | 3.6 | 87.1 | 69.2 | 75.0 | 97.1 | 53.1 | 41 | 314 | |
| DH MCWC UHC | 48.4 9.4 50.5 | 37.1 9.4 28.0 | 82.3 65.1 78.6 | 90.3 75.5 93.5 | 90.3 87.2 93.9 | 100.0 99.0 100.0 | 3.6 3.2 3.7 | 91.9 81.3 87.7 | 64.5 47.6 74.9 | 90.3 28.5 83.5 | 96.8 94.7 97.6 | 50.0 56.3 52.8 | 4 7 30 | 62 95 157 | |
| Union-level | 17.9 | 11.2 | 50.0 | 62.8 | 75.7 | 97.9 | 2.9 | 57.4 | 49.9 | 42.4 | 90.8 | 49.2 | 295 | 413 | |
| UHFWC USC/RD | 19.3 14.1 | 13.2 6.1 | 55.7 35.6 | 68.7 47.8 | 79.4 66.3 | 98.6 96.1 | 3.0 2.5 | 62.4 44.7 | 56.3 33.6 | 46.3 32.5 | 90.2 92.4 | 54.9 34.7 | 212 83 | 274 139 | |
| Community clinic (CC) NGO clinic/hospital Private hospital | 17.0 33.1 4.9 | 25.6 33.3 0.8 | 42.8 66.1 69.8 | 76.7 81.5 39.9 | 77.9 99.2 94.5 | 89.5 100.0 100.0 | 2.9 3.5 3.0 | 50.7 88.2 85.3 | 70.5 62.8 10.5 | 78.8 61.6 10.5 | 84.5 94.9 91.0 | 55.8 53.6 76.5 | 939 25 104 | 461 109 109 | |
| Location | | | | | | | | | | | | | | | |
| Urban Rural | 15.5 17.5 | 9.7 22.3 | 67.1 45.3 | 53.8 73.7 | 93.7 77.7 | 100.0 91.6 | 3.1 2.9 | 82.9 53.2 | 29.4 65.5 | 29.7 70.1 | 90.7 86.4 | 70.0 54.1 | 154 1,250 | 417 989 | |
| Division | | | | | | | | | | | | | | | |
| Barishal Chattogram Dhaka | 26.4 20.4 18.5 | 17.2 26.0 18.1 | 43.0 48.7 45.3 | 69.9 67.3 59.5 | 78.3 83.9 78.6 | 87.3 93.2 95.3 | 2.8 2.9 2.8 | 40.2 55.6 54.9 | 69.3 69.3 59.1 | 72.9 71.9 61.4 | 94.5 90.1 78.2 | 52.6 58.8 60.6 | 103 283 266 | 134 236 234 | |
| Khulna Rajshahi Rangpur | 9.8 14.9 14.5 | 12.2 19.2 25.1 | 43.3 64.5 40.0 | 80.6 67.1 85.1 | 82.1 84.5 72.2 | 96.5 92.0 89.0 | 3.0 3.1 2.9 | 66.2 51.4 62.1 | 73.4 41.8 61.3 | 75.0 44.4 68.8 | 94.8 88.0 93.0 | 60.0 59.0 54.5 | 161 203 185 | 177 198 169 | |
| Sylhet Mymensingh | 17.0 18.5 | 24.8 23.9 | 59.6 35.5 | 82.4 76.4 | 84.0 67.5 | 91.4 90.9 | 3.2 2.7 | 60.8 59.1 | 70.3 53.3 | 79.3 63.0 | 82.6 72.7 | 49.7 36.0 | 88 115 | 118 140 | |
| Total | 17.3 | 20.9 | 47.7 | 71.6 | 79.5 | 92.5 | 2.9 | 56.4 | 61.6 | 65.7 | 86.9 | 55.8 | 1,404 | 1,406 | |
| Total excluding CCs | 18.1 | 11.4 | 57.7 | 61.2 | 82.7 | 98.7 | 3.0 | 67.9 | 43.5 | 39.2 | 91.6 | 55.9 | 465 | 945 | |
| Total public | 18.0 | 22.3 | 45.6 | 73.9 | 77.9 | 91.8 | 2.9 | 53.5 | 65.7 | 70.3 | 86.4 | 54.2 | 1,275 | 1,188 | |
| Total public excluding | 21.0 | 13.0 | 53.3 | 66.2 | 77 7 | 98 1 | 29 | 61.0 | 52.3 | 46 4 | 91.6 | 49 7 | .336 | 727 | |

¹ A scale with a gradation of 250 grams, or a digital standing scale with a gradation of 250 grams or less where an adult can hold a child to be weighed.

² Average number of four equipment items (child weighing scale, length or height board, thermometer, and stethoscope) available for providing child curative care.

³ A scale with a gradation of 100 grams, or a digital standing scale with a gradation of 100 grams where an adult can hold an infant to be weighed.

⁴ A functioning timer or watch with seconds hand or the provider has a watch with seconds hand, or other devices (e.g., cell phone) that can measure seconds.

Table 4.2.1: Equipment for inpatient child curative care services

Among facilities that offer inpatient curative care for sick children, the percentages having the indicated equipment at inpatient service area, by background characteristics, Bangladesh HFS 2022

| | | | | | | Equip | oment | | | | | | Number of facilities | Number of |
|--|--|--|--|--|--|---|--|--|--|--|--|--|---|---|
| Background characteristic | Child weighing scale ¹ | Infant weighing scale ² | Thermometer | Stethoscope | Nebulizer ³ | Timer ⁴ | Torch | Growth chart | Height board | Pulse oximeter | Nasal catheter/ cannula ⁵ | Oxygen | inpatient curative care for sick children (weighted) | inpatient curative care for sick children (unweighted) |
| Facility type | | | | | | | | | | | | | | |
| District and upazila public facilities | 63.4 | 69.5 | 95.8 | 97.6 | 92.6 | 92.0 | 79.5 | 43.7 | 58.4 | 80.7 | 59.0 | 96.4 | 32 | 215 |
| DH MCWC UHC | 83.1 41.0 61.4 | 78.0 70.7 68.1 | 91.5 83.0 97.1 | 98.3 88.7 97.9 | 93.2 30.4 95.5 | 96.6 88.7 91.5 | 88.1 82.7 78.0 | 54.2 53.4 41.6 | 78.0 76.7 54.5 | 83.1 59.4 81.3 | 67.8 36.5 58.7 | 98.3 65.4 97.6 | 4 1 27 | 59 17 139 |
| Union-level public facilities UHFWC USC/RD | - | - | - | - | - | - | - | - | - | - | - | - | 0 0 | 0 0 |
| Community clinic | | - | - | - | - | - | - | - | - | - | - | - | 0 | 0 |
| NGO clinic/hospital Private hospital | 64.6 62.2 | 77.4 77.7 | 100.0 91.1 | 100.0 94.8 | 54.6 91.9 | 67.4 87.1 | 70.3 95.6 | 72.0 18.8 | 70.9 26.8 | 51.8 90.4 | 44.8 69.0 | 32.0 93.3 | 1 68 | 9 61 |
| Location | | | | | | | | | | | | | | |
| Urban Rural | 63.1 60.1 | 75.6 71.2 | 92.1 97.4 | 95.6 97.1 | 91.1 94.0 | 87.8 92.2 | 91.9 77.3 | 26.1 39.3 | 36.5 47.5 | 86.9 84.0 | 66.7 56.3 | 93.3 93.1 | 88 13 | 213 72 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 66.8 64.8 66.1 35.4 58.0 54.8 78.9 58.2 | 86.5 71.0 76.0 78.6 74.6 61.2 97.2 68.4 | 98.3 94.6 87.2 96.1 91.2 100.0 100.0 92.8 | 96.5 100.0 87.2 97.2 100.0 96.0 100.0 100.0 | 88.2 98.8 86.5 92.4 86.8 90.4 82.3 96.3 | 100.0 96.5 79.1 94.4 81.1 96.0 81.7 86.2 | 92.6 87.0 92.9 79.8 98.2 84.4 94.3 80.4 | 66.0 24.5 21.6 35.9 21.2 43.7 40.1 34.4 | 73.7 33.0 37.5 46.6 32.8 47.5 45.8 28.1 | 92.1 88.9 84.2 77.1 88.9 77.1 94.3 83.8 | 57.6 69.9 51.3 75.1 71.9 70.2 79.5 73.1 | 98.3 99.0 88.0 94.9 85.6 96.4 91.5 96.3 | 4 33 31 8 11 5 5 4 | 27 54 58 35 37 28 22 24 |
| Total | 62.6 | 75.0 | 92.8 | 95.8 | 91.5 | 88.4 | 89.8 | 28 | 38.1 | 86.5 | 65.3 | 93.3 | 101 | 285 |
| Total excluding CCs | 62.6 | 75.0 | 92.8 | 95.8 | 91.5 | 88.4 | 89.8 | 28 | 38.1 | 86.5 | 65.3 | 93.3 | 101 | 285 |
| Total public | 63.4 | 69.5 | 95.8 | 97.6 | 92.6 | 92.0 | 79.5 | 43.7 | 58.4 | 80.7 | 59.0 | 96.4 | 32 | 215 |
| Total public excluding CCs | 63.4 | 69.5 | 95.8 | 97.6 | 92.6 | 92.0 | 79.5 | 43.7 | 58.4 | 80.7 | 59.0 | 96.4 | 32 | 215 |

"-" Means facilities do not offer inpatient curative care for sick children

¹ A scale with a gradation of 250 grams, or a digital standing scale with a gradation of 250 grams or less where an adult can hold a child to be weighed.

² A scale with a gradation of 100 grams, or a digital standing scale with a gradation of 100 grams where an adult can hold an infant to be weighed.

³ A functioning nebulizer to administer medication through inhalation for respiratory conditions

⁴ A functioning timer or watch with seconds hand or the provider has watch with seconds hand, or other devices (e.g., cell phone) that can measure seconds.

⁵ A nasal cannula or catheter for delivering supplemental oxygen to children with lower oxygen levels

Table 4.3: Trained staff for child curative care services

Among facilities that offer outpatient curative care for sick children, the percentages having indicated trained staff, by background characteristics, Bangladesh HFS 2022

| | | Train | | Number of Number of | | |
|--|------------------------------|-------------|------------------------------|------------------------|--|--|
| | IMC | | Growth mo | onitoring ² | facilities offering | facilities offering |
| Background characteristic | During the past 24 months | At any time | During the past 24 months | At any time | curative care for sick children (weighted) | curative care for sick children (unweighted) |
| Facility type | | | | | | |
| District and upazila public facilities | 36.2 | 74.5 | 30.1 | 64.8 | 41 | 314 |
| DH | 35.5 | 83.9 | 35.5 | 75.8 | 4 | 62 |
| MCWC | 10.5 | 52.7 | 8.7 | 41.9 | 7 | 95 |
| UHC | 42.2 | 78.2 | 34.3 | 68.4 | 30 | 157 |
| Union-level public facilities | 4.0 | 44.8 | 8.1 | 35.1 | 295 | 413 |
| UHFWC | 3.1 | 45.0 | 8.7 | 38.0 | 212 | 274 |
| USC/RD | 6.3 | 44.3 | 6.8 | 28.0 | 83 | 139 |
| Community clinic (CC) | 10.4 | 52.4 | 15.8 | 56.3 | 939 | 461 |
| NGO clinic/hospital | 23.4 | 53.0 | 13.9 | 42.0 | 25 | 109 |
| Private hospital | 5.4 | 16.2 | 5.4 | 10.9 | 104 | 109 |
| Location | | | | | | |
| Urban | 12.1 | 31.9 | 9.9 | 27.3 | 154 | 417 |
| Rural | 9.4 | 50.9 | 14.3 | 51.1 | 1,250 | 989 |
| Division | | | | | | |
| Barishal | 10.4 | 65.0 | 11.5 | 58.8 | 103 | 134 |
| Chattogram | 16.9 | 49.4 | 18.3 | 54.7 | 283 | 236 |
| Dhaka | 5.7 | 43.1 | 10.1 | 39.6 | 266 | 234 |
| Khulna | 5.1 | 58.3 | 13.7 | 51.3 | 161 | 177 |
| Rajshahi | 6.6 | 41.6 | 5.1 | 40.6 | 203 | 198 |
| Rangpur | 13.4 | 48.6 | 14.8 | 47.2 | 185 | 169 |
| Sylhet | 7.7 | 34.8 | 12.8 | 47.3 | 88 | 118 |
| Mymensingh | 7.6 | 56.4 | 27.6 | 57.1 | 115 | 140 |
| Total | 9.7 | 48.8 | 13.8 | 48.5 | 1,404 | 1,406 |
| Total excluding CCs | 8.3 | 41.5 | 9.8 | 32.7 | 465 | 945 |
| Total public | 9.7 | 51.4 | 14.5 | 51.6 | 1,275 | 1,188 |
| Total public excluding CCs | 8.0 | 48.7 | 10.9 | 39.0 | 336 | 727 |

¹ At least one provider of child health services in the facility reported receiving in-service training in IMCI. Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

² At least one provider of child health services in the facility reported receiving in-service training in growth monitoring (SAM and MAM training). Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

Table 4.4: Availability of essential and priority medicines

Among facilities offering outpatient curative care services for sick children, the percentages where indicated essential and priority medicines to support care for the sick child were observed to be available in the facility on the day of the survey, Bangladesh HFS 2022

| | | | | Essential | medicines | | | | | Priority r | nedicines | | | |
|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|
| Background | ORS ¹ | Amoxicillin syrup, suspension, or dispersible ¹ | Cotrimoxazole syrup, suspension, or dispersible | Paracetamol syrup or suspension ¹ | Vitamin A capsules ¹ | Mebendazole/ albendazole | Zinc tablets or syrup | All 6 essential medicines available ² | Ampicillin powder for injection | Ceftriaxone powder for injection | Gentamycin for injection | Benzathine benzylpenicillin for injection | Number of facilities offering outpatient curative care for sick children (weighted) | Number of facilities otfering outpatient curative care for sick children (unweighted) |
| Facility type | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 86.7 98.4 24.6 99.3 | 67.5 32.3 86.3 68.3 | 34.5 9.7 82.0 27.2 | 85.1 85.5 85.4 84.9 | 34.0 16.1 8.2 42.6 | 75.1 30.6 95.8 76.9 | 68.6 74.2 6.2 82.2 | 18.2 1.6 1.0 24.6 | 32.4 11.3 1.1 42.8 | 74.8 91.9 28.7 83.0 | 69.4 58.1 37.1 78.6 | 11.6 24.2 2.1 12.0 | 41 4 7 30 | 314 62 95 157 |
| Union-level public facilities UHFWC USC/RD | 37.3 18.9 84.4 | 79.8 91.3 50.6 | 65.6 81.2 26.0 | 83.7 91.6 63.6 | 5.8 5.2 7.3 | 83.8 97.7 48.5 | 13.7 6.9 31.1 | 2.4 1.9 3.4 | 0.6 0.3 1.6 | 1.4 0.4 3.8 | 13.5 18.2 1.5 | 2.0 2.2 1.3 | 295 212 83 | 413 274 139 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 98.5 86.4 65.3 | 92.7 29.6 24.3 | 24.5 8.2 11.0 | 95.3 63.0 52.8 | 83.0 37.6 28.4 | 94.8 51.1 48.7 | 81.0 58.9 47.7 | 64.1 10.7 12.4 | 0.5 3.2 10.2 | 0.2 34.8 66.0 | 0.2 13.3 50.7 | 2.5 0.8 10.1 | 939 25 104 | 461 109 109 |
| Location Urban Rural | 73.1 83.8 | 32.9 89.2 | 14.2 34.3 | 59.7 92.4 | 31.4 64.1 | 54.5 91.8 | 52.1 65.0 | 14.5 48.7 | 11.7 1.0 | 60.7 1.6 | 46.0 4.4 | 9.6 2.4 | 154 1,250 | 417 989 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 86.3 86.7 80.8 79.2 78.6 83.4 81.4 84.9 | 87.7 80.1 78.7 85.5 79.9 86.8 88.2 88.2 88.0 | 33.5 35.1 32.7 34.5 28.4 32.1 28.5 28.4 | 92.9 90.3 90.0 91.3 83.2 86.5 87.6 89.9 | 66.7 55.4 66.3 53.9 61.7 60.9 59.9 | 86.9 88.0 86.7 89.9 85.0 88.0 93.4 87.2 | 61.2 65.7 64.5 58.6 58.1 63.9 67.1 72.0 | 44.6 43.6 43.1 49.4 41.9 43.2 48.6 51.9 | 1.5 3.3 3.2 0.9 0.8 2.3 3.5 1.2 | 5.2 10.9 13.4 2.9 8.4 4.8 7.4 4.9 | 5.5 10.3 14.9 7.3 7.0 5.9 8.8 6.2 | 0.8 4.2 5.6 1.4 0.5 3.1 1.7 6.2 | 103 283 266 161 203 185 88 115 | 134 236 234 177 198 169 118 140 |
| Total | 82.6 | 83.0 | 32.1 | 88.8 | 60.5 | 87.7 | 63.6 | 45.0 | 2.2 | 8.1 | 9.0 | 3.2 | 1,404 | 1,406 |
| Total excluding CCs | 50.6 | 63.6 | 47.6 | 75.8 | 15.1 | 73.4 | 28.6 | 6.4 | 5.7 | 24.1 | 26.7 | 4.6 | 465 | 945 |
| Total public | 83.9 | 88.9 | 34.3 | 92.3 | 63.6 | 91.6 | 65.0 | 48.3 | 1.5 | 2.9 | 5.5 | 2.7 | 1,275 | 1,188 |
| Total public excluding CCs | 43.3 | 78.3 | 61.8 | 83.9 | 9.2 | 82.8 | 20.4 | 4.3 | 4.5 | 10.3 | 20.3 | 3.1 | 336 | 727 |

Note: The essential medicines comprise the medicines and commodities indicators for assessing readiness to provide preventative and curative child health services within the health facility assessment methodology proposed by the WHO and USAID (WHO, 2012). ORS = Oral rehydration solution

¹ These medicines and commodities are also in the group of priority medicines for children.

² ORS, amoxicillin syrup/suspension or dispersible, paracetamol syrup/suspension, vitamin A capsules, mebendazole/albendazole, and zinc tablets or syrup.

Table 4.5: Infection control and laboratory diagnostic capacity

Among facilities offering outpatient curative care services for sick children, the percentages with indicated items for infection control observed to be available at the service site on the day of the survey and the percentages with the indicated laboratory diagnostic capacity in the facility, by background characteristics, Bangladesh HFS 2022

| | | | | Items | for infection co | ontrol | | | | Labo diag cap | oratory nostic acity | Number of facilities | Number of facilities |
|--|--|--|--|--|--|--|--|--|--|--|--|---|---|
| Background characteristic | Soap | Running water ¹ | Soap and running water | Alcohol- based hand disin- fectant | Soap and running water or else alcohol- based hand disinfectant | Latex gloves ² | Sharps container | Waste recep- tacle ³ | Average of 6 items available ⁴ | Hemo- globin⁵ | Stool micro- scopy ⁶ | offering outpatient curative care for sick children (weighted) | offering outpatient curative care for sick children (unweighted) |
| Facility type | | | | | | | | | | | | | |
| District and upazila public facilities | 70.1 | 79.4 | 67.5 | 84.5 | 92.3 | 56.8 | 33.9 | 57.8 | 3.8 | 86.8 | 37.2 | 41 | 314 |
| DH MCWC UHC | 75.8 73.7 68.5 | 83.9 83.1 77.9 | 75.8 73.7 64.8 | 90.3 78.9 84.9 | 98.4 89.4 92.1 | 56.5 85.2 50.3 | 38.7 60.1 27.1 | 67.7 58.7 56.1 | 4.1 4.4 3.6 | 93.5 59.9 92.0 | 61.3 - 42.0 | 4 7 30 | 62 95 157 |
| Union-level public facilities UHFWC | 52.9 57.2 | 56.7 60.9 | 49.2 54.0 | 66.8 66.0 | 76.1 76.7 | 68.9 77.2 | 52.1 61.2 | 54.6 58.5 | 3.5 3.8 | 23.2 28.0 | - | 295 212 | 413 274 |
| USC/RD | 42.0 | 46.3 | 37.0 | 69.1 | 74.5 | 47.9 | 29.0 | 44.5 | 2.8 | 11.0 | - | 83 | 139 |
| Community clinic (CC) | 53.3 | 49.4 | 44.1 | 65.8 | 73.5 | 57.5 | 60.0 | 52.4 | 3.4 | 4.2 | - | 939 | 461 |
| NGO clinic/hospital Private hospital | 81.8 79.1 | 80.2 82.0 | 80.2 79.1 | 87.8 89.2 | 94.2 92.7 | 76.9 71.4 | 56.1 28.1 | 67.1 64.6 | 4.5 4.1 | 89.3 88.3 | 16.6 44.2 | 25 104 | 109 109 |
| Location Urban Rural | 74.3 53.9 | 77.8 52.0 | 74.0 46.0 | 87.2 66.4 | 91.4 74.5 | 69.0 60.3 | 33.2 57.9 | 62.2 53.2 | 4.0 3.4 | 84.8 10.2 | 38.2 0.5 | 154 1,250 | 417 989 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh Total Total <i>Total excluding CCs</i> | 37.9 62.3 52.3 62.0 55.5 58.0 56.6 55.7 56.1 61.8 53.8 | 39.1 56.9 53.4 62.8 52.7 58.1 63.1 47.8 54.8 65.7 52.1 | 35.8 52.9 48.8 57.0 46.4 50.5 51.3 42.3 49.1 59.2 46.0 | 56.8 75.7 72.2 72.9 66.6 61.2 64.0 67.6 68.7 74.5 66.7 | 64.7 79.1 75.0 85.5 77.9 74.9 71.7 73.9 76.4 82.2 74.7 | 60.4 60.2 66.5 71.3 52.3 54.8 72.5 55.7 61.2 68.8 60.1 | 44.1 53.5 56.2 55.5 60.2 65.3 56.8 55.2 45.4 57.3 | 49.3 54.8 48.6 62.8 53.8 52.1 58.3 58.6 54.2 57.8 53.1 | 2.9 3.6 3.5 3.8 3.4 3.4 3.4 3.5 3.7 3.7 | 9.2 24.8 25.6 11.8 22.2 7.5 21.3 12.1 18.4 46.9 11.3 | 3.6 6.0 7.6 1.5 6.1 1.8 4.0 2.5 4.7 14.0 1.2 | 103 283 266 161 203 185 88 115 1,404 465 1.275 | 134 236 234 177 198 169 118 140 1,406 945 1.188 |
| i otal public | 53.8 | 52.1 | 46.0 | 66.7 | /4./ | 60.1 | 57.3 | 53.1 | 3.4 | 11.3 | 1.2 | 1,275 | 1,188 |
| l otal public excluding CCs | 55.0 | 59.5 | 51.4 | 69.0 | 78.1 | 67.4 | 49.9 | 55.0 | 3.6 | 31.0 | 4.5 | 336 | 727 |

Note: "-" Means facilities do not have provision of stool microscopy test.

The laboratory diagnostic capacity indicator measures presented in the table comprise the diagnostics domain for assessing readiness to provide preventative and curative child health services within the health facility assessment methodology proposed by the WHO and USAID (WHO, 2012). MCWCs, union-level facilities, and CCs do not provide stool microscopy tests.

¹ Piped water, water in bucket with specially fitted tap, or water in pour pitcher.

² Non-latex equivalent gloves are acceptable.

³ Waste receptacle with plastic bin liner.

⁴ Average number of six items (soap, running water, alcohol-based disinfectant, latex gloves, sharps container, and waste receptacle) available.

⁵ Facility had functioning equipment and reagents for hematology analyser, colorimeter, hemoglobinometer, hemoCue, or tallquist paper.

⁶ Facility had a functioning microscope with glass slides and formal saline (for concentration method), normal saline (for direct method), or Lugol's iodine solution.

Table 4.6: Readiness of health facilities to provide child curative care services

Among facilities that offer outpatient curative care for sick children, the percentages with IMCI guidelines, staff trained in IMCI at any time, basic equipment, and essential medicines available on the day of the survey, and the average readiness score, Bangladesh HFS 2022

| Background characteristic | IMCI guidelines | Staff trained in IMCI (at any time) ¹ | Child weighing scale ² | Thermo- meter | Growth chart | Zinc tablets or syrup | ORS | Amoxicillin syrup, suspension or dispersible | Para- cetamol syrup or suspen- sion | Meben- dazole/ albendazole | Average readiness score (out of 10 items) ³ | Number of facilities offering outpatient curative care for sick children (weighted) | Number of facilities offering outpatient curative care for sick children (unweighted) |
|--|--------------------|--|---|------------------|-----------------|-----------------------------|--------------|--|---|----------------------------------|---|--|--|
| Facility type | | | | | | | | | | | | | |
| District and upazila public facilities | 43.4 | 74.5 | 76.7 | 92.4 | 69.2 | 68.6 | 86.7 | 67.5 | 85.1 | 75.1 | 7.4 | 41 | 314 |
| DH | 48.4 | 83.9 | 82.3 | 90.3 | 64.5 | 74.2 | 98.4 | 32.3 | 85.5 | 30.6 | 6.9 | 4 | 62 |
| MCWC | 9.4 | 52.7 | 65.1 | 87.2 | 47.6 | 6.2 | 24.6 | 86.3 | 85.4 | 95.8 | 5.6 | 7 | 95 |
| UHC | 50.5 | 78.2 | 78.6 | 93.9 | 74.9 | 82.2 | 99.3 | 68.3 | 84.9 | 76.9 | 7.9 | 30 | 157 |
| Union-level public facilities | 17.9 | 44.8 | 50.0 | 75.7 | 49.9 | 13.7 | 37.3 | 79.8 | 83.7 | 83.8 | 5.4 | 295 | 413 |
| UHFWC | 19.3 | 45.0 | 55.7 | 79.4 | 56.3 | 6.9 | 18.9 | 91.3 | 91.6 | 97.7 | 5.6 | 212 | 274 |
| USC/RD | 14.1 | 44.3 | 35.6 | 66.3 | 33.6 | 31.1 | 84.4 | 50.6 | 63.6 | 48.5 | 4.7 | 83 | 139 |
| Community clinic (CC) | 17.0 | 52.4 | 42.8 | 77.9 | 70.5 | 81.0 | 98.5 | 92.7 | 95.3 | 94.8 | 7.2 | 939 | 461 |
| NGO clinic/ hospital Private hospital | 33.1 4.9 | 53.0 16.2 | 66.1 69.8 | 99.2 94.5 | 62.8 10.5 | 58.9 47.7 | 86.4 65.3 | 29.6 24.3 | 63.0 52.8 | 51.1 48.7 | 6.0 4.3 | 25 104 | 109 109 |
| Location | | | | | | | | | | | | | |
| Urban | 15.5 | 31.9 | 67.1 | 93.7 | 29.4 | 52.1 | 73.1 | 32.9 | 59.7 | 54.5 | 5.1 | 154 | 417 |
| Rural | 17.5 | 50.9 | 45.3 | 77.7 | 65.5 | 65.0 | 83.8 | 89.2 | 92.4 | 91.8 | 6.8 | 1,250 | 989 |
| Division | | | | | | | | | | | | | |
| Barishal | 26.4 | 65.0 | 43.0 | 78.3 | 69.3 | 61.2 | 86.3 | 87.7 | 92.9 | 86.9 | 7.0 | 103 | 134 |
| Chattogram | 20.4 | 49.4 | 48.7 | 83.9 | 69.3 | 65.7 | 86.7 | 80.1 | 90.3 | 88.0 | 6.8 | 283 | 236 |
| Dhaka | 18.5 | 43.1 | 45.3 | 78.6 | 59.1 | 64.5 | 80.8 | 78.7 | 90.0 | 86.7 | 6.5 | 266 | 234 |
| Khulna | 9.8 | 58.3 | 43.3 | 82.1 | 73.4 | 58.6 | 79.2 | 85.5 | 91.3 | 89.9 | 6.7 | 161 | 177 |
| Rajshahi | 14.9 | 41.6 | 64.5 | 84.5 | 41.8 | 58.1 | 78.6 | 79.9 | 83.2 | 85.0 | 6.3 | 203 | 198 |
| Rangpur | 14.5 | 48.6 | 40.0 | 72.2 | 61.3 | 63.9 | 83.4 | 86.8 | 86.5 | 88.0 | 6.5 | 185 | 169 |
| Sylhet | 17.0 | 34.8 | 59.6 | 84.0 | 70.3 | 67.1 | 81.4 | 88.2 | 87.6 | 93.4 | 6.8 | 88 | 118 |
| Mymensingh | 18.5 | 56.4 | 35.5 | 67.5 | 53.3 | 72.0 | 84.9 | 88.0 | 89.9 | 87.2 | 6.5 | 115 | 140 |
| Total | 17.3 | 48.8 | 47.7 | 79.5 | 61.6 | 63.6 | 82.6 | 83.0 | 88.8 | 87.7 | 6.6 | 1,404 | 1,406 |
| Total excluding CCs | 18.1 | 41.5 | 57.7 | 82.7 | 43.5 | 28.6 | 50.6 | 63.6 | 75.8 | 73.4 | 5.4 | 465 | 945 |
| Total public | 18.0 | 51.4 | 45.6 | 77.9 | 65.7 | 65.0 | 83.9 | 88.9 | 92.3 | 91.6 | 6.8 | 1,275 | 1,188 |
| Total public excluding CCs | 21.0 | 48.7 | 53.3 | 77.7 | 52.3 | 20.4 | 43.3 | 78.3 | 83.9 | 82.8 | 5.6 | 336 | 727 |

Note: ORS = oral rehydration solution.

¹ At least one provider of child health services in the facility reported receiving in-service training in IMCI. Training refers only to in-service training. The training must have involved ² A scale with gradation of 250 grams, or a digital standing scale with gradation of 250 grams or less where an adult can hold a child to be weighed.

³ Average readiness score is the average number of 10 items (IMCI guideline, staff trained in IMCI at any time, child weighing scale, thermometer, growth chart, zinc tablets/syrup, ORS, amoxicillin syrup/suspension/dispersible, paracetamol syrup/suspension, and mebendazole/albendazole) available that are necessary for providing outpatient curative care for sick children.

Table 4.7: Supportive management for providers of child health services

Among interviewed child health service providers, the percentages who report receiving training related to their work and personal supervision during the specified time periods, by background characteristics, Bangladesh HFS 2022

| | Percentage of interviewed providers who received: | | | | | | | |
|---|--|--|--|--|---|--|--|--|
| Background characteristic | Training related to child health during the 24 months preceding the survey ¹ | Personal supervision during the 6 months preceding the survey ² | Training related to child health during the 24 months and personal supervision during the 6 months preceding the survey | Number of interviewed child health service providers (weighted) | Number of interviewed child health service providers (unweighted) | | | |
| Facility type | | | | | | | | |
| District and upazila public facilities DH MCWC | 18.0 17.4 8.0 | 97.6 96.8 98.0 | 17.8 17.2 8.0 | 960 219 18 | 1,968 394 256 | | | |
| UHC | 18.4 | 97.8 | 18.2 | 723 | 1,318 | | | |
| Union-level public facilities UHFWC USC/RD | 17.3 17.5 16.8 | 86.7 89.4 79.3 | 13.8 14.7 11.3 | 355 259 96 | 644 434 210 | | | |
| Community clinic (CC) | 32.7 | 91.9 | 30.3 | 1,308 | 704 | | | |
| NGO clinic/hospital | 32.4 | 96.2 | 32.4 | 84 | 289 | | | |
| Private hospital | 9.5 | 86.6 | 5.8 | 378 | 233 | | | |
| Location | | | | | | | | |
| Urban | 16.2 | 93.7 | 14.8 | 1,099 | 1,800 | | | |
| Rural | 27.6 | 91.8 | 25.3 | 1,986 | 2,038 | | | |
| Division Barishal | 18.3 | 90.5 | 13.7 | 246 | 418 | | | |
| Chattogram Dhaka | 28.6 19.9 | 92.2 94 5 | 27.7 | 565 690 | 629 676 | | | |
| Khulna Rajshahi Bangur | 23.5 18.1 | 90.9 94.3 | 21.8 16.6 22.8 | 310 456 304 | 469 516 | | | |
| Sylhet Mymensingh | 26.8 33.0 | 93.0 89.3 | 24.2 28.1 | 158 266 | 277 336 | | | |
| Total | 23.5 | 92.5 | 21.6 | 3,085 | 3,838 | | | |
| Total excluding CCs | 16.7 | 93.0 | 15.1 | 1,777 | 3,134 | | | |
| Total public | 25.2 | 93.3 | 23.5 | 2,623 | 3,316 | | | |
| Total public excluding CCs | 17.8 | 94.6 | 16.7 | 1,315 | 2,612 | | | |

¹ Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.
 ² Personal supervision refers to any form of technical support or supervision from a facility-based supervisor or from a visiting supervisor. It may include, but is not limited to, review of records and observation of work, with or without any feedback to the health worker.

Table 4.8: Training for child health service providers

Among interviewed child health service providers, the percentages who report receiving in-service training on topics related to child health during the specified period before the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of providers of child health services who reported that they received in-service training on: | | | | | | | | | Number of | |
|--|--|----------------|---------------------------------|----------------|---------------------------------|----------------|---------------------------------|----------------------|------------------------------------|--------------------------------------|--|
| | EPI/cold | l chain | IMO | CI | AI | RI | Diarrhoea d treati | liagnosis or ment | interviewed child health | interviewed child health | |
| Background characteristic | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | service providers (weighted) | service providers (unweighted) | |
| Facility type | | | | | | | | | | | |
| District and upazila public facilities | 4.9 | 20.9 | 5.7 | 22.5 | 5.0 | 21.9 | 5.8 | 23.6 | 960 | 1,968 | |
| DH | 4.9 | 16.3 | 5.2 | 21.6 | 3.5 | 18.1 | 5.0 | 18.6 | 219 | 394 | |
| MCWC | 2.0 | 24.0 | 3.4 | 29.2 | 3.0 | 28.7 | 1.8 | 27.7 | 18 | 256 | |
| UHC | 5.0 | 22.2 | 5.9 | 22.5 | 5.5 | 22.8 | 6.1 | 25.0 | 723 | 1,318 | |
| Union-level public facilities | 4.7 | 29.1 | 2.5 | 31.8 | 2.7 | 29.2 | 3.6 | 35.6 | 355 | 644 | |
| UHFWC | 3.4 | 30.1 | 2.1 | 32.2 | 2.6 | 29.8 | 3.4 | 35.7 | 259 | 434 | |
| USC/RD | 8.2 | 26.4 | 3.6 | 30.8 | 3.1 | 27.8 | 4.0 | 35.5 | 96 | 210 | |
| Community clinic (CC) | 6.6 | 37.4 | 8.6 | 48.1 | 7.3 | 34.4 | 16.4 | 59.0 | 1,308 | 704 | |
| NGO clinic/hospital | 23.9 | 43.9 | 10.3 | 31.0 | 12.1 | 30.5 | 10.9 | 28.6 | 84 | 289 | |
| Private hospital | 1.2 | 4.5 | 1.9 | 6.0 | 0.9 | 5.6 | 2.4 | 8.3 | 378 | 233 | |
| Location | | | | | | | | | | | |
| Urban | 5.5 | 17.0 | 4.5 | 17.7 | 3.7 | 16.0 | 5.4 | 18.9 | 1.099 | 1.800 | |
| Rural | 5.7 | 33.2 | 7.2 | 40.9 | 6.4 | 31.9 | 12.1 | 49.0 | 1,986 | 2,038 | |
| Division | | | | | | | | | | | |
| Barishal | 7.5 | 30.5 | 7.3 | 47.2 | 7.6 | 31.6 | 11.7 | 48.4 | 246 | 418 | |
| Chattogram | 8.2 | 31.9 | 10.0 | 33.8 | 9.7 | 33.3 | 12.0 | 39.7 | 565 | 629 | |
| Dhaka | 4.6 | 20.6 | 4.5 | 26.9 | 3.6 | 20.2 | 10.4 | 29.2 | 690 | 676 | |
| Khulna | 4.6 | 30.6 | 2.5 | 37.3 | 3.3 | 24.9 | 4.8 | 37.1 | 310 | 469 | |
| Rajshahi | 4.7 | 28.7 | 5.5 | 29.1 | 3.0 | 22.8 | 5.2 | 36.4 | 456 | 516 | |
| Rangpur | 6.4 | 25.1 | 7.2 | 27.0 | 6.1 | 24.8 | 13.6 | 38.5 | 394 | 517 | |
| Sylhet | 4.6 | 20.4 | 7.7 | 27.7 | 7.4 | 26.9 | 8.9 | 37.6 | 158 | 277 | |
| Mymensingh | 3.4 | 34.7 | 5.0 | 43.6 | 3.1 | 31.3 | 9.9 | 54.1 | 266 | 336 | |
| Total | 5.6 | 27.4 | 6.2 | 32.6 | 5.4 | 26.3 | 9.7 | 38.3 | 3,085 | 3,838 | |
| Total excluding CCs | 5.0 | 20.1 | 4.5 | 21.2 | 4.0 | 20.3 | 4.9 | 23.0 | 1,777 | 3,134 | |
| Total public | 5.7 | 30.2 | 6.7 | 36.5 | 5.8 | 29.1 | 10.8 | 42.9 | 2,623 | 3,316 | |
| Total public excluding CCs | 4.8 | 23.1 | 4.8 | 25.0 | 4.4 | 23.9 | 5.2 | 26.9 | 1,315 | 2,612 | |

Note: EPI = Expanded Program on Immunization; IMCI = Integrated Management of Childhood Illness; ARI = Acute respiratory infection

Table 4.9: Key indicators for child curative care services in MCWCs

Among facilities offering outpatient child curative care for sick children, the percentages with key indicators, Bangladesh HFS 2022 $\,$

| | Facility type | | | | | |
|--|--|--|--|--|--|--|
| Indicators | MCWC (district-level) | MCWC (upazila and union-level) | | | | |
| Availability | | | | | | |
| Child curative care Child growth monitoring Child vaccination | 98.3 72.3 66.1 | 89.8 61.1 42.2 | | | | |
| Number of facilities (weighted) | 5 | 3 | | | | |
| Number of facilities (unweighted) | 62 | 38 | | | | |
| Readiness | | | | | | |
| IMCI guidelines Staff trained on IMCI (at any time) ¹ Child weighing scale ² Thermometer Growth chart Zinc tablets/syrup ORS Amoxicillin syrup, suspension or dispersible Paracetamol syrup, suspension Mebendazole/albendazole | 9.8 58.9 70.1 91.7 49.4 9.6 31.4 90.4 92.0 98.4 | 8.5 41.7 56.0 79.3 44.4 0.0 12.5 79.0 73.6 91.1 | | | | |
| Average readiness score ³ | 6.0 | 4.9 | | | | |
| Number of facilities offering outpatient child curative care (weighted) | 4 | 3 | | | | |
| Number of facilities offering outpatient child curative care (unweighted) | 61 | 34 | | | | |

¹ At least one provider of child health services in the facility reported receiving in-service training in IMCI. Training refers only to in-service training. The training must have involved structured sessions and does not include individual instruction that a

Provider might have received during routine supervision.
2 A scale with gradation of 250 grams, or a digital standing scale with gradation of 250 grams or less where an adult can hold a child to be weighed.
3 Average readiness score is the average number of items (out of 10 items) available that are necessary for providing outpatient curative care for sick children.

FAMILY PLANNING SERVICES

Key Findings

- About 8 in 10 public facilities provide at least one modern family planning (FP) method (Table 5.2 and Figure 5.1).
- Modern FP services are universally provided in MCWCs (94%), UHCs (97%), and UHFWCs (92%), but only 10% of private hospitals provide modern FP (Table 5.2 and Figure 5.1).
- The proportion of DHs providing any modern FP services increased remarkably, from 69% in 2017 to 84% in 2022 (Table 5.2 and Figure 5.1).
- Two-thirds of the public facilities, excluding CCs (68%), provide long-acting reversible contraceptives or permanent methods (LARC/PMs) (**Table 5.2**).
- The majority of the DHs (57%), MCWCs (67%), and UHCs (79%) provide male or female sterilization services (Table 5.2 and Figure 5.3).
- More than two-thirds of the public facilities offering any modern FP, excluding CCs (70%), have guidelines available on FP. In most UHCs (89%) guidelines are available, while availability is very low in private facilities (7%) (Table 5.6 and Figure 5.9).
- Almost all MCWCs (95%) and most DHs (81%) and UHCs (87%) offering any modern FP have staff ever trained in FP services. The majority of the public facilities (60%) have staff trained at any time, whereas only one-fifth of them (19%) have staff trained in the preceding 24 months (Table 5.6).
- Among facilities offering any modern FP, a blood pressure apparatus is universally available in MCWCs (97%), UHCs (98%), UHFWCs (90%), NGO clinics (97%), and private hospitals (100%) (Table 5.8 and Figure 5.11).
- The availability of the combined or progestin-only oral pills is universal at UHCs (94%) and UHFWCs (95%) among facilities offering any modern FP methods, but this availability is lower in NGO (74%) and private (34%) facilities. Progestin-only injectables are available in most UHCs (94%) and UHFWCs (93%). Nearly 80% of public health facilities, excluding CCs, have intrauterine contraceptive devices (IUCDs) available on the day of the survey (Table 5.8).
- The average readiness score for FP services in public health facilities that offer any modern FP methods is 4.4 (out of 8 items) (Table 5.8). Comparatively, the readiness score is 6.9 among MCWCs and 7.3 among UHCs. The readiness score has slightly increased in total public health facilities, excluding CCs, from 6.0 in 2017 to 6.1 in 2022 (Table 5.8 and Figure 5.13).

5.1 BACKGROUND

amily planning (FP) is important for the health of a mother and her children, as well as the family's economic situation. FP comprises a group of activities that permit couples to decide freely on the spacing and number of their children. FP improves the health of mothers, children, and entire families and is a key element in upholding reproductive rights. Therefore, wherever maternal health, reproductive health, or child health services are provided, facilities should strive to increase the appropriate use of FP and contraceptive services and to provide client education.

The Government of Bangladesh has a National Population Policy (MOHFW 2012) that seeks to reduce fertility to replacement level, and the 4th Health, Population and Nutrition Sector Program (HPNSP) has embraced strategies for making FP services available, accessible, acceptable, and affordable for all women and men of reproductive age at public, private, and nongovernmental (NGO) health facilities. The HPNSP strategies aimed to increase overall FP use to 75% by 2022 (MOHFW 2012).

Over the past four decades, Bangladesh has made remarkable progress in increasing the utilization of modern FP methods among currently married women. According to the 2022 Bangladesh Demographic and Health Survey (BDHS), the contraceptive prevalence rate (CPR) increased from 56% to 64% between 2007 and 2022. Between 2017–18 and 2022, use of modern contraceptive methods increased from 52% to 55%. The 4th HPNSP aimed to reach 50% use of modern methods in Chattogram and Sylhet by 2022. Since 2014, use of modern methods in Sylhet increased from 41% to 44%; however, there was a slight increase in Chattogram, from 47% to 49% (NIPORT, et al. 2014; NIPORT, et al. 2023). The 4th HPNSP aimed to reach a modern contraceptive use prevalence of 60% in Chattogram and Sylhet by 2022 (Government of Bangladesh 2017).

- Availability of FP services: Section 5.2, including Tables 5.1–5.5 and Figures 5.1–5.3, examines the availability of FP services. FP services are considered available if a facility provides FP methods to clients at the facility, prescribes methods that the client must obtain elsewhere, or counsels about FP.
- Availability of FP Commodities: Section 5.3, including Tables 5.5 and 5.8; and Figures 5.4–5.8, examines the availability of FP commodities. It includes oral pills, injectables, condoms, IUCDs, and implants.
- Availability of guidelines, trained staff, equipment, and infection control items: Section 5.4, including Tables 5.6 and 5.7; and Figures 5.9–5.12, addresses the extent to which facilities that offer FP services have the capacity to support quality services, including the necessary service guidelines, trained staff, equipment, and infection control materials.
- Readiness of health facilities to provide FP services: Section 5.5, including Table 5.8 and Figure 5.13, describes the average readiness score of health facilities to offer FP services using the WHO-specified items or tracer indicators that facilities must have to be considered ready to offer such services.
- Basic management and administrative systems: Section 5.6, including Tables 5.9 and 5.10, evaluates aspects of training and supervision that are important to support the delivery of high-quality FP services.

5.2 FAMILY PLANNING SERVICES

FP methods differ in their functions, effectiveness, side effects, and modes of use. The acceptability and desirability of FP methods also differ among users. Thus, a facility that offers a wide variety of FP methods is best able to meet client needs. Although a broad mix of methods is optimal, facilities are expected to differ in the exact mix of methods they offer because of differences in provider qualifications, training, and the infrastructure required to safely provide certain methods. Methods that can be offered safely with minimal

training are pills, injectables, and condoms, along with counseling on FP methods. Safely offering implants, IUCDs, female sterilization, and male sterilization requires a higher level of skill and a more developed infrastructure.

The 2022 BHFS obtained information on the availability of FP services (modern FP methods, long-acting reversible contraceptives or permanent methods [LARC/PMs], and male or female sterilization) at each of the public, private, and NGO health facilities in the survey sample. This section of the report uses the following definitions in assessing the availability of FP services:

- A facility is considered to **offer** FP services if the facility reports that it provides a specified FP method, prescribes the method for clients to obtain elsewhere, or counsels clients on the method. Facilities in this category do not necessarily provide FP methods to clients.
- A facility is said to **provide** FP services if the facility reports that it stocks a specific method and makes it available to clients when they visit the facility. Facilities in this category provide FP methods to clients.

5.2.1 Availability of Family Planning Services

Table 5.1 shows the availability of FP services at facilities that provide a method, prescribe the method for clients to obtain elsewhere, or counsel clients on the method (i.e., facilities that do not necessarily provide an FP method to the client).

- Eight in ten public health facilities offer (provide, prescribe, or counsel clients) any modern FP (including an emergency contraceptive) in 2022. Almost all DHs (94%), MCWCs (95%), UHCs (98%), and UHFWCs (96%) offer these services, whereas it is offered by only one-fourth of private hospitals. The services are more available at the facilities in rural (83%) areas compared to the facilities in urban areas (39%) (**Table 5.1**).
- Seventy-nine percent of public facilities, excluding CCs, offer long-acting or permanent methods of family planning service. Almost all of the MCWCs (93%), UHCs (97%), and most of the UHFWCs (89%) offer LARC/PMs; however, less than half of USC/RDs (45%) and only one-fourth of the private hospitals (24%) offer these services (Table 5.1).
- Twenty-nine percent (34%, excluding CCs) of public health facilities offer male or female sterilization. The majority of MCWCs (77%) and most UHCs (87%) offer male and female sterilization; however, only 17% of private hospitals offer these services (**Table 5.1**).
- Ninety-one percent of public facilities, excluding CCs, offer postpartum family planning (PPFP) services. UHCs (99%), UHFWCs (99%), MCWCs (98%), DH (97%), NGOs (90%), and CCs (89%), universally offer PPFP services; private hospitals (47%) are least likely to offer these services (Table 5.1).
- Modern FP services (with or without stock) are universally offered by DHs (95%), MCWCs (99%), UHCs (98%), and UHFWCs (100%), whereas it is provided by 54% of USC/RDs and 30% of private hospitals (Table 5.1).
- Modern FP services (with stock) are available in almost all MCWCs (94%), UHCs (97%) and UHFWCs (92%). Almost two-thirds of NGOs (60%) have availability, whereas it is available in only 10% of private facilities (Table 5.1).

5.2.2 Facilities Providing Modern Family Planning Methods and Services

This section describes the extent to which modern FP methods are provided (i.e., the facility stocks the specific method and makes it available for clients) at different types of facilities (**Table 5.2**).

- More than three-quarters of public facilities (77%) provide at least one modern FP method in 2022 (Table 5.2 and Figure 5.1).
- Methods of modern FP are universally provided by MCWCs (94%), UHCs (97%), and UHFWCs (92%). The majority of CCs (75%) and NGO clinics (60%) provide any modern FP methods; however, only one-tenth of private hospitals offer these services (Table 5.2 and Figure 5.1).
- The proportion of public facilities providing any modern FP methods decreased from 88% in 2017 to 77% in 2022. However, availability of modern FP services increased remarkably in DHs, from 69% in 2017 to 84% in 2022 (**Table 5.2** and **Figure 5.1**).
- The percentage of FP service availability varies by location. Urban health facilities provided a lower proportion of modern FP services (26%) than the rural facilities (76%) (**Table 5.2**).



Figure 5.1 Facilities providing modern FP methods and services, by facility type

- The majority of the public facilities, excluding CCs, (68%) in Bangladesh provide LARC/PMs (Table 5.2).
- About 8 in 10 DHs (79%) and more than ninety percent of the MCWCs (91%) and UHCs (95%) are providing LARC/PM services in 2022 (Table 5.2 and Figure 5.2).
- The proportion of DHs providing LARC/PMs increased remarkably from 66% in 2017 to 79% in 2022 and remained unchanged in UHCs (95%); however, it decreased notably in USC/RDs and NGO facilities over the past five years (**Figure 5.2**).
- The LARC/PMs are provided more in urban (23%) facilities than rural facilities (18%) (Table 5.2).





■2014 BHFS ■2017 BHFS ■2022 BHFS

- The majority of the DHs (57%), MCWCs (67%), and UHCs (79%) are providing male or female sterilization services in 2022 (Table 5.2 and Figure 5.3). Since 2017, there have been notable increases in sterilization services in DHs (from 47% in 2017 to 57% in 2022) and UHCs (from 70% in 2017 to 79% in 2022) (Figure 5.3).
- The provision of male or female sterilization has markedly increased 23 percentage points in DHs, from 34% in 2014 to 57% in 2022 (**Figure 5.3**).



Figure 5.3 Facilities providing male or female sterilization services, by facility type



5.2.3 Frequency of Availability of Family Planning Services

Percent

- Fifty-six percent of public facilities reported that they provide FP services every day. Most of the DHs (83%), MCWC (87%), UHCs (84%), private hospitals (86%), and NGOs (92%) that offer FP services provide these services every day (**Table 5.4**).
- However, relatively smaller proportions of union-level facilities (56%) and CCs (55%) offer FP services every day (**Table 5.4**).
- Frequency of FP service availability varies by location. More urban facilities (86%) provide FP services every day than rural facilities (56%) (**Table 5.4**).
- By division, the proportion of all facilities offering FP services every day ranges from 40% in Mymensingh to 74% in Barishal (**Table 5.4**).

5.3 AVAILABILITY OF FP COMMODITIES ON THE DAY OF ASSESSMENT

Stock-outs of FP methods can put a woman at risk of an unintended pregnancy. To obtain information on stockouts, the 2022 BHFS assessed the availability on the day of survey of each method reportedly provided. Table 5.5 shows the stock of FP commodities which were observed to be available on the day of survey among facilities that reported that they provide the specific commodity.

The majority of the public facilities (79%) providing FP commodities (oral pills, injectables, condoms, IUCDs, implants) actually had stock of every method they provide available on the day of survey (Table 5.5).

- Nine in ten public health facilities, excluding CCs, (91%) providing FP commodities (oral pills, injectables, condoms, IUCDs, and implants) actually had stock of every method they provide available on the day of the survey (**Table 5.5**).
- UHCs are most likely (92%) to have every FP commodity available in stock (**Table 5.5**).

5.3.1 Availability of Oral Pills

- Among facilities that offer (provide, prescribe, or counsel clients) any modern FP methods, the availability of oral pills is universal at UHCs (94%) and UHFWCs (95%). Most of the CCs (87%) have oral pills available on the day of the survey; however, only 34% of private facilities have oral pills, a decrease from 44% in 2014 (**Table 5.8** and **Figure 5.4**).
- Seventy-nine percent of DHs have oral pills available among facilities that reported that they offer any modern family planning methods (Table 5.8). Whereas, among facilities providing oral pills, almost all DHs (98%) were observed to have stock of oral pills on the day of the survey (Table 5.5).



Figure 5.4 Availability of oral pills, by facility type

5.3.2 Availability of Injectables

- Ninety percent of public facilities, excluding CCs, had injectables on the day of the survey among facilities that offer (provide, prescribe, or counsel clients) any modern FP methods (Table 5.8 and Figure 5.5).
- Injectables are universally available in UHCs (94%) and UHFWCs (93%). The availability of injectables in NGO facilities has declined twenty-two percentage points, from 88% in 2017 to 66% in 2022 (Table 5.8 and Figure 5.5).

The availability of injectables in DHs is 71% among facilities offering any modern FP methods (Table 5.8); however, stock of injectables was observed on the day of the survey in 89% of the DHs providing injectables (Table 5.5).



Figure 5.5 Availability of injectables, by facility type

5.3.3 Availability of Condoms

- Although there has been an overall decline in the availability of male condoms in public facilities (from 91% in 2014 to 79% in 2022), they are universally available in UHCs (94%) and UHFWCs (94%) on the day of the survey among facilities that offer (provide, prescribe, or counsel clients) any modern FP methods (**Table 5.8** and **Figure 5.6**).
- Thirty-two percent of private hospitals have male condoms available among facilities offering any modern FP methods (Table 5.8 and Figure 5.6). However, 96% of the private hospitals were observed to have stock of male condoms on the day of the survey among facilities providing condoms (Table 5.5).



Figure 5.6 Availability of condoms, by facility type

5.3.4 Availability of Intrauterine Contraceptive Devices

- Seventy-nine percent of public health facilities, excluding CCs, have IUCDs available in 2022, an increase from 75% in 2017 among facilities that offer (provide, prescribe, or counsel clients) any modern FP methods. Availability of IUCDs for contraception are lowest in private facilities (3%). More than 9 in 10 UHCs (91%) and 8 in 10 MCWCs (85%) had IUCDs available on the day of visit (Table 5.8 and Figure 5.7).
- The availability of IUCDs in Chattogram division is 24% among facilities offering any modern FP methods (**Table 5.8**). Whereas, stock of IUCDs are observed in 99% of all facilities (including NGOs and private) in Chattogram division among facilities providing any modern FP methods (**Table 5.5**).



Figure 5.7 Availability of IUCDs, by facility type



5.3.5 Availability of Implants (One Rod or Two Rod)

- Nineteen percent of public health facilities, excluding CCs, have implants available in 2022, among facilities that offer (provide, prescribe, or counsel clients) any modern FP methods, which increased from 15% in 2017. In the last five years, the availability of implants increased in all types of facilities except MCWCs, UHCs, and private facilities (**Table 5.8** and **Figure 5.8**).
- Two-thirds of MCWCs had implants available on the day of the survey, among facilities offering any modern FP methods (Table 5.8). However, almost all MCWCs (94%) were observed to have the stock of implants among facilities providing condoms on the day of survey (Table 5.5).


Figure 5.8 Availability of Implants in health facilities, by facility type



5.4 AVAILABILITY OF SERVICE GUIDELINES, TRAINED STAFF, AND EQUIPMENT

To address the extent to which facilities that offer FP services have the capacity to support quality services, the 2022 BHFS gathered data on the percentage of facilities having guidelines, trained staff, and basic equipment for FP services.

5.4.1 Availability of Guidelines for Family Planning Services

- Almost 39% of public facilities offering FP had guidelines available on the day of the survey. UHCs are most likely (89%) and private hospitals are least likely (7%) to have FP guidelines in 2022 (Table 5.6 and Figure 5.9).
- Between 2017 and 2022, the availability of guidelines at DHs increased sharply (from 61% to 74%), while the percentage of MCWCs with guidelines remained unchanged. UHCs and UHFWCs showed small decreases in the availability of FP guidelines (Figure 5.9).
- The overall availability of FP guidelines in public facilities decreased remarkably since 2017 (from 69% to 39%), mainly due to a decrease in the availability at CCs (from 39% to 27%) and NGOs (87% to 77%) (Figure 5.9).

Percent



Figure 5.9 Availability of FP guidelines in health facilities, by facility type Percent

5.4.2 Availability of Trained Staff in Family Planning

Training is also an important management function to support health care providers. In particular, periodic inservice training can keep providers up to date and help them refresh their knowledge and skills.

- The majority of the public facilities (60%) have staff trained in FP at any time, and 19% have staff trained during the last 24 months. Ninety-five percent of MCWCs and 87% of UHCs have staff trained at any time, but much lower percentages (58% and 39%, respectively) of these facilities have staff trained during the preceding 24 months (**Table 5.6**).
- Since 2017, the availability of trained staff at any time in public health facilities increased by five percentage points; however, it decreased slightly at UHCs, UHFWCs, and NGO facilities (**Figure 5.10**).
- Private hospitals (39%) and CCs (53%) are the least likely to have ever trained staff members in FP (Table 5.6).



Figure 5.10 Availability of staff trained in FP at any time in health facilities, by facility type

5.4.3 Availability of Equipment for Family Planning Services

- Most public health facilities (80%) offering modern FP methods had a functioning BP apparatus available on the day of the visit. A BP apparatus was universally available at MCWCs (97%), UHCs (98%), UHFWCs (90%), NGOs (97%), and private hospitals (100%) (Table 5.8 and Figure 5.11).
- Overall, 80% of public facilities (90% excluding CCs) have a functioning BP apparatus, while few public facilities have a pelvic model (17%) for IUCD demonstration and a model for showing condom use (18%) (Table 5.6).
- Most public facilities, excluding CCs, (85%) have an examination bed or couch, and the majority of public facilities, excluding CCs, (68%) have an examination light (Table 5.6).



Figure 5.11 Availability of functioning blood pressure apparatus in health facilities, by facility type

5.4.4 Infection Control During Provision of Family Planning

Infection control is essential in FP procedures such as insertion or removal of IUCDs, injectables, and implants. The 2022 BHFS collected data on the percentage of facilities offering six infection control items (soap, running water, alcohol-based hand disinfectant, latex gloves, sharps container, and waste receptacle) during provision of modern FP methods.

- The majority of the public health facilities have alcohol-based hand disinfectant (66%), latex gloves (62%), and soap (57%) available at the time of survey (**Table 5.7**).
- The average number of infection control items available at FP service sites in public health facilities is 3.5 out of 6 in 2022. Among public facilities, excluding CCs, the average number of infection control items increased slightly, from 3.6 in 2017 to 4.0 in 2022 (**Table 5.7** and **Figure 5.12**).
- Among facilities offering any modern FP methods, UHCs have the highest average number of items for infection control-4.9 items-in the 2022 BHFS (**Table 5.7** and **Figure 5.12**).



NGO

Private

CCs

Figure 5.12 Average number of items (out of six) available for infection control in family planning service areas, by facility type

5.5 READINESS OF HEALTH FACILITIES TO PROVIDE FAMILY PLANNING SERVICES

UHC

■2014 BHFS ■2017 BHFS ■2022 BHFS

(N=1,359)

UHFWC USC/RD

(N=1,182)

MCWC

(N=1,260)

The WHO has specified a set of items or tracer indicators that facilities must have to be considered ready to offer FP services (WHO 2013, Service Availability and Readiness Assessment). Data from the 2014, 2017, and 2022 BHFS are used to construct a slightly less restrictive measure of FP service readiness that is appropriate for the Bangladesh context. The following eight items/indicators are included in this measure of the readiness of health facilities to provide FP services:

Trained staff:

Total

public

Total

public excluding CCs DH

- At least one staff person who has ever received in-service FP training
- Guidelines:
 - National or any other FP guidelines
- Equipment:
 - Blood pressure apparatus
- Commodities:
 - Oral pills
 - Injectable
 - Condom
 - IUCD
 - Implant (one rod or two rod)

The readiness of facilities for providing FP services is assessed through an average readiness score. This score represents the mean number of available items or tracer indicators from the set of eight adapted items for FP services on the day of visit.

- The average readiness score for providing FP services in public health facilities is 4.4 in 2022 (Table 5.8). It increased slightly in total public health facilities (excluding CCs), from 6.0 in 2017 to 6.1 in 2022 (Figure 5.13).
- Among facilities offering any modern FP, MCWCs (6.9) and UHCs (7.3) have the highest average score for FP services in the 2022 BHFS. However, private facilities have an average of 2.2 items (Table 5.8 and Figure 5.13).

Figure 5.13 Average readiness score of health facilities (out of 8 items) to provide FP services, by facility type



5.6 **BASIC MANAGEMENT AND ADMINISTRATIVE SYSTEMS**

5.6.1 Training and Supervision

Average

Supportive management for FP providers, including in-service training and personal supervision, is needed to ensure quality services. Training related to FP involves in-service training with structured sessions, while personal supervision includes technical support or supervision from a facility-based supervisor or from a visiting supervisor. Table 5.9 presents the percentage of FP service providers who reported having training and supervision according to facility type, location, and division.

- Eighteen percent of service providers in public facilities (excluding providers at CCs) reported receiving training related to FP services during the two years before the survey (Table 5.9).
- More than 9 in 10 providers (93%) at public health facilities received personal supervision during the • six months before the survey (Table 5.9).

- Overall, 16% of providers at public health facilities received both training (related to FP during the past two years) and personal supervision during the past six months (**Table 5.9**).
- Service providers at MCWCs (28%), NGOs (24%), and DHs (23%) have received more in-service training and supervision, compared to providers at UHCs (14%) and CCs (15%) who are the least likely to have received training and supervision (**Table 5.9**).

5.6.2 In-Service Training for Family Planning Service Providers by Topic

The types of in-service training and the extent of training received by FP service providers incorporates FP counseling, FP-related clinical issues, insertion or removal of IUCDs and implants, FP for HIV-positive clients, postpartum FP, vasectomy, tubal ligation, and emergency contraception.

- Approximately half (46%) of providers in public health facilities have received training on FP counseling at some point, but only 12% received such training during the past 24 months of the survey (Table 5.10).
- Similarly, one-fourth of the providers have received training at any time on FP-related clinical issues in public health facilities, but only 6% received training during the past 24 months (**Table 5.10**).
- The proportion of providers receiving training at any time before the survey on insertion or removal of IUCDs (22%) is almost twice as high as the proportion receiving training on insertion or removal of implants (13%) in public health facilities (**Table 5.10**).
- In public health facilities, nearly 30% of providers have ever received training on postpartum FP, with 7% reporting such training during the last 24 months (**Table 5.10**).
- In general, providers in public health facilities in urban areas are more likely than those in rural areas to have received training during the last 24 months. For example, 8% of urban providers received training on FP-related clinical issues, compared with 5% of rural providers. Similarly, 9% of providers in urban areas reported having training on insertion or removal of IUCDs, compared with 3% in rural areas (Table 5.10).
- Six percent of providers in public health facilities in urban areas reported they have been trained on insertion or removal of implants, compared with 3% in rural areas during the last 24 months. The percentages of providers receiving training on postpartum FP in the last 24 months are also higher in urban (11%) than rural areas (6%) (**Table 5.10**).

FAMILY PLANNING SERVICES

LIST OF TABLES

- Table 5.1 Availability of family planning services
- Table 5.2 Methods of family planning provided
- Table 5.3 Methods of family planning provided
- Table 5.4 Frequency of availability of family planning services
- Table 5.5 Availability of family planning commodities
- Table 5.6 Guidelines, trained staff, and basic equipment for family planning services
- Table 5.7 Items for infection control during provision of family planning
- Table 5.8 Readiness of health facilities to provide family planning services
- Table 5.9 Supportive management for providers of family planning services
- Table 5.10 Training for family planning service providers
- Table 5.11 Key indicators for family planning services in MCWCs

Table 5.1: Availability of family planning services

Among all facilities, the percentages that offer — provide, prescribe, or counsel clients — family planning methods, by background characteristics, Bangladesh HFS 2022

| | | Ν | lethods of family p | lanning (FP) | | | | |
|--|--|--|--|--|---|--|---|--|
| Background characteristic | Percentage offering any modern FP (including an emergency contraceptive) ¹ | Percentage offering any long- acting or permanent methods ² | Percentage offering male or female sterilization ³ | Percentage offering post- partum FP services ⁴ | Percentage offering modern FP methods and services with/without stock | Percentage offering modern FP methods and services with stock | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type District and upazila public facilities DH MCWC | 97.1 93.5 95.1 | 95.3 87.2 93.2 | 84.3 79.0 76.9 | 98.8 96.8 97.9 | 98.0 95.2 99.0 | 95.3 83.9 94.0 | 41 4 7 | 319 62 100 |
| Union-level public facilities UHFWC USC/RD | 83.5 96.0 50.3 | 76.6 88.8 44.5 | 27.7 32.1 16.1 | 89.4 98.9 64.2 | 87.5 100.0 54.2 | 78.3 91.8 42.4 | 310 225 85 | 434 293 141 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 82.1 74.8 25.0 | 32.7 66.0 23.6 | 26.8 44.1 16.9 | 89.4 89.9 47.1 | 88.7 86.4 29.5 | 75.3 59.9 10.3 | 994 29 183 | 488 127 189 |
| Location Urban Rural | 38.6 82.6 | 36.1 43.8 | 27.9 27.6 | 58.7 89.3 | 43.6 88.5 | 26.1 76.0 | 235 1,322 | 514 1,043 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 82.0 80.9 70.7 70.3 75.4 79.3 74.4 78.6 | 45.9 37.0 35.5 41.0 55.2 51.5 36.7 43.5 | 32.4 21.7 22.1 25.2 34.9 41.3 22.4 26.0 | 90.6 88.0 80.3 83.4 85.3 86.9 76.1 87.0 | 90.1 86.2 76.3 77.1 79.6 83.5 78.5 89.0 | 79.3 71.3 65.8 63.9 60.0 73.0 72.7 71.0 | 111 291 317 210 214 193 99 122 | 152 244 264 208 212 190 136 151 |
| Total | 75.9 | 42.7 | 27.6 | 84.7 | 81.7 | 68.5 | 1,557 | 1,557 |
| Total excluding CCs | 65.0 | 60.2 | 29.2 | 76.4 | 69.3 | 56.5 | 563 | 1,069 |
| Total public | 82.9 | 44.8 | 28.8 | 89.7 | 88.7 | 76.6 | 1,345 | 1,241 |
| Total public excluding CCs | 85.1 | 78.8 | 34.4 | 90.5 | 88.7 | 80.3 | 351 | 753 |

Note: A facility is considered to offer FP services if the facility reports that it provides a specified FP method, prescribes the method for clients to obtain elsewhere, or counsels clients on the method. Facilities in this category do not necessarily provide FP methods to clients.

¹ Facility provides, prescribes, or counsels clients on any of the following: contraceptive pills (combined or progestin-only), injectables (progestin-only), one-rod implant, two-rod implant (zadell), IUCDs, male condoms, female sterilization (tubal ligation) or male sterilization (vasectomy), and emergency contraceptive pills. ² Facility provides, prescribes, or counsels clients on any of the following long-term and permanent methods of FP: one-rod implant, two-rod implant, IUCDs, female ³ Providers in the facility perform male or female sterilization (vasectomy).
 ⁴ Facility provides, prescribes, or counsel clients with postpartum family planning (PPFP) services.

Table 5.2: Methods of family planning provided

Among all facilities, the percentages that provide clients with specific modern family planning methods by background characteristics, Bangladesh HFS 2022

| | Meth | ods of family planning | g (FP) | | |
|-------------------------------|--|--|--|---------------------------------------|--------------------------------------|
| Background characteristic | Percentage that provide any modern FP (including an emergency contraceptive) ¹ | Percentage that provide any long- acting or permanent methods ² | Percentage that provide male or female sterilization ³ | Number of facilities (weighted) | Number of facilities (unweighted) |
| Facility type | | | | | |
| District and upazila | | | | | |
| public facilities | 95.3 | 92.5 | 74.1 | 41 | 319 |
| DH | 83.9 | 79.0 | 56.5 | 4 | 62 |
| MCWC | 94.0 | 91.1 | 66.9 | 7 | 100 |
| UHC | 97.3 | 94.8 | 78.5 | 30 | 157 |
| Union-level public facilities | 78.3 | 64.3 | - | 310 | 434 |
| UHFWC | 91.8 | 76.6 | - | 225 | 293 |
| USC/RD | 42.4 | 31.7 | - | 85 | 141 |
| Community clinic (CC) | 75.3 | - | - | 994 | 488 |
| NGO clinic/hospital | 59.9 | 45.3 | 20.0 | 29 | 127 |
| Private hospital | 10.3 | 10.0 | 10.0 | 183 | 189 |
| Location | | | | | |
| Urban | 26.1 | 22.9 | 18.3 | 235 | 514 |
| Rural | 76.0 | 17.5 | 2.3 | 1,322 | 1,043 |
| Division | | | | | |
| Barishal | 79.3 | 18.7 | 2.8 | 111 | 152 |
| Chattogram | 71.3 | 20.6 | 5.5 | 291 | 244 |
| Dhaka | 65.8 | 21.1 | 7.8 | 317 | 264 |
| Khulna | 63.9 | 17.1 | 3.3 | 210 | 208 |
| Rajshahi | 60.0 | 14.5 | 2.7 | 214 | 212 |
| Rangpur | 73.0 | 16.1 | 3.9 | 193 | 190 |
| Sylhet | 72.7 | 19.8 | 5.6 | 99 | 136 |
| Mymensingh | 71.0 | 16.2 | 2.9 | 122 | 151 |
| Total | 68.5 | 18.3 | 4.7 | 1,557 | 1,557 |
| Total excluding CCs | 56.5 | 47.8 | 11.7 | 563 | 1,069 |
| Total public | 76.6 | 18.9 | 3.6 | 1,345 | 1,241 |
| Total public excluding CCs | 80.3 | 67.6 | 11.9 | 351 | 753 |

Note: To be considered as providing a specific FP method, the facility must report that it stocks the method and makes it available to clients without clients having to go elsewhere to obtain it. In the case of male or female sterilization, the facility reports that providers in the facility perform the procedures. "-" Means the specific FP methods are not provided at union-level public facilities or at community clinics.

¹ Facility provides any of the following: contraceptive pills (combined or progestin-only), injectables (progestin-only), one-rod implant, two-rod implant (zadell), IUCDs, male condoms, female sterilization (tubal ligation), or male sterilization (vasectomy), and emergency contraceptive pills. ² Facility provides any of the following long-term and permanent FP methods: one-rod implant, two-rod implant, IUCDs, female

sterilization (tubal ligation), or male sterilization (vasectomy). ³ Perform male or female sterilization in the facility.

Table 5.3: Methods of family planning provided¹

Facility type District and Unionupazila level Commu-NGO Total Total public public public nity clinic clinic/ Private excluding excluding Methods provided facilities DH MCWC UHC facilities UHFWC USC/RD (CC) hospital hospital Total CCs Total public CCs Combined or progestin-only 90.3 74.2 85.0 93.9 78.0 91.5 42.4 73.1 53.4 6.7 66.3 54.5 74.7 79.5 oral pill 90.1 74.2 84.1 93.9 75.8 89.5 39.5 55.1 50.6 2.0 53.8 51.6 61.0 77.5 Progestin-only injectable Male condom 89.9 71.0 85.0 93.9 76.7 90.5 40.3 69.2 51.0 6.7 63.5 53.6 71.5 78.3 Intrauterine contraceptive 86.5 71.0 74.5 40.1 43.4 17.7 65.6 81.1 90.1 62.8 31.7 1.4 16.2 device -75.6 38.7 65.9 83.4 9.1 10.1 6.6 27.2 1.3 4.7 12.4 4.7 17.0 Implant (one or two rods) -10.5 10.6 Tubal ligation 73.1 56.5 66.9 77.1 12.9 10.0 4.3 2.3 ----64.5 6.5 58.9 74.5 14.3 0.7 2.1 5.7 2.0 7.6 Vasectomy At least 1 modern method 95.3 83.9 94.0 97.3 78.0 91.5 42.4 75.3 59.2 10.3 68.4 56.3 76.5 80.1 At least 2 temporary modern methods² 90.4 75.8 85.0 93.9 77.8 91.5 41.5 68.6 54.5 6.7 63.5 54.4 71.4 79.3 At least 4 temporary modern methods² 87.1 67.1 81.1 91.5 63.0 75.4 30.3 0.9 37.1 1.3 16.3 43.4 17.9 65.9 Any LAPM 92.5 79.0 91.1 94.8 64.3 76.6 31.7 1.6 45.3 10.0 18.3 47.8 18.9 67.6 Male or female sterilization 74.1 56.5 66.9 78.5 3.6 4.7 0.9 0.7 20.0 10.0 4.7 11.7 3.6 11.9 Number of facilities (weighted) 41 4 7 30 310 225 85 994 29 183 1,557 563 1,345 351 Number of facilities 319 62 100 157 434 293 127 189 141 488 1,557 1,069 1,241 753 (unweighted)

Among all facilities, the percentages that provide clients with specific modern family planning methods, by facility type, Bangladesh HFS 2022

Note: "-" Means the specific FP methods are not provided at this type of facility.

¹ To be considered as providing a specific FP method, the facility must report that it stocks the method and makes it available to clients without clients having to go elsewhere to obtain it. In the case of vasectomy and tubal ligation, the facility reports that providers in the facility perform the procedures.

² Any methods other than male or female sterilization.

Table 5.4: Frequency of availability of family planning services

Among facilities that offer any modern family planning services, the percentages that offer the services every day or less frequently than every day, by background characteristics, Bangladesh HFS 2022

| | Percentage of faci | lities offering FP | Number of | Number of |
|---|-----------------------------|------------------------|---------------------|---------------------|
| | services ¹ where | e services are | facilities offering | facilities offering |
| | offered on ind | icated days: | any modern FP | any modern FP |
| Background characteristic | Provides but not | Provides | services | services |
| | every day ² | every day ² | (weighted) | (unweighted) |
| Facility type | | | | |
| District and upazila public facilities DH | 15.5 | 84.5 | 40 | 307 |
| | 17.2 | 82.8 | 4 | 58 |
| MCWC | 12.6 | 87.4 | 7 | 95 |
| UHC | 15.9 | 84.1 | 29 | 154 |
| Union-level public facilities | 44.1 | 55.9 | 258 | 350 |
| UHFWC | 42.0 | 58.0 | 216 | 280 |
| USC/RD | 55.0 | 45.0 | 42 | 70 |
| Community clinic (CC) | 44.8 | 55.2 | 816 | 400 |
| NGO clinic/hospital | 7.6 | 92.4 | 22 | 91 |
| Private hospital | 14.1 | 85.9 | 46 | 49 |
| Location | | | | |
| Urban | 14.3 | 85.7 | 91 | 334 |
| Rural | 44.1 | 55.9 | 1,091 | 863 |
| Division | | | | |
| Barishal | 26.5 | 73.5 | 91 | 120 |
| Chattogram | 30.9 | 69.1 | 235 | 201 |
| Dhaka | 39.2 | 60.8 | 224 | 203 |
| Khulna | 43.6 | 56.4 | 148 | 155 |
| Rajshahi | 50.6 | 49.4 | 161 | 161 |
| Rangpur | 50.2 | 49.8 | 153 | 143 |
| Sylhet | 38.9 | 61.1 | 74 | 103 |
| Mymensingh | 60.5 | 39.5 | 96 | 111 |
| Total | 41.8 | 58.2 | 1,182 | 1,197 |
| Total excluding CCs | 35.1 | 64.9 | 366 | 797 |
| Total public | 43.6 | 56.4 | 1,114 | 1,057 |
| Total public excluding CCs | 40.3 | 59.7 | 298 | 657 |

¹ Includes services for contraceptive pills (combined or progestin-only), injectables (progestin-only), implants, IUCDs, male condoms, tubal ligation, vasectomy, and emergency contraceptive pills.
² Every day refers to all working days when the facility is open.

Table 5.5: Availability of family planning commodities

Among facilities that provide¹ the indicated modern family planning method, the percentages where the commodity was observed to be available on the day of the survey, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Combined or progestin-only oral pills | Progestin-only injectable | Male condom | Intrauterine contraceptive device | Implant (one rod or two rod) | Every method provided by facility was available on day of survey |
|--|---|------------------------------|-------------|---|------------------------------------|---|
| Facility type | | | | | | |
| District and upazila public facilities | 98.8 | 98.1 | 97.8 | 97.4 | 95.7 | 90.5 |
| DH | 97.8 | 89.1 | 95.5 | 88.6 | 79.2 | 78.7 |
| MCWC | 100.0 | 100.0 | 98.8 | 98.8 | 93.6 | 89.2 |
| UHC | 98.7 | 98.7 | 97.8 | 98.1 | 97.2 | 92.2 |
| Union-level public facilities | 98.4 | 96.6 | 97.8 | 96.2 | 73.2 | 90.9 |
| UHFWC | 98.9 | 97.6 | 98.2 | 96.4 | 75.5 | 92.0 |
| USC/RD | 95.2 | 90.8 | 95.0 | 94.7 | 64.1 | 84.8 |
| Community clinic (CC) | 95.7 | 79.1 | 87.0 | - | - | 75.0 |
| NGO clinic/hospital | 97.2 | 94.8 | 97.8 | 91.3 | 85.8 | 76.3 |
| Private hospital | 100.0 | 19.2 | 95.9 | 34.2 | 30.4 | 71.5 |
| Location | | | | | | |
| Urban | 99.6 | 92.0 | 98.6 | 91.5 | 87.3 | 83.7 |
| Rural | 96.4 | 84.5 | 89.8 | 94.5 | 73.7 | 79.0 |
| Division | | | | | | |
| Barishal | 98.9 | 96.7 | 88.0 | 92.3 | 98.0 | 87.9 |
| Chattogram | 93.9 | 90.3 | 93.6 | 98.7 | 90.9 | 83.0 |
| Dhaka | 94.8 | 86.9 | 92.1 | 92.9 | 69.7 | 82.1 |
| Khulna | 99.7 | 80.4 | 93.3 | 93.5 | 77.5 | 78.8 |
| Rajshahi | 99.3 | 79.8 | 92.2 | 93.0 | 98.7 | 76.9 |
| Rangpur | 94.7 | 82.3 | 81.2 | 96.5 | 83.4 | 71.0 |
| Sylhet | 97.6 | 83.4 | 91.8 | 96.9 | 76.9 | 84.1 |
| Mymensingh | 97.3 | 77.9 | 85.7 | 83.4 | 45.2 | 67.9 |
| Total | 96.5 | 84.9 | 90.3 | 94.1 | 79.2 | 79.2 |
| Total excluding CCs | 98.4 | 95.7 | 97.7 | 95.5 | 83.3 | 89.3 |
| Total public | 96.5 | 85.0 | 90.1 | 94.8 | 80.2 | 79.3 |
| Total public excluding CCs | 98.4 | 96.8 | 97.8 | 96.4 | 85.0 | 90.9 |

Note: The denominators for each characteristic method combination are different and are not shown in the table. "-" Means the specific FP methods are not provided at this type of facility.

¹ To be considered as providing a specific FP method, the facility must report that it stocks the method and makes it available to clients without clients having to go elsewhere to obtain it.

The combined or progestin-only oral contraceptive pills, injectable contraceptives, and male condom measures presented in the table comprise the medicines and commodities domain for assessing readiness to provide FP services within the health facility assessment methodology proposed by WHO and USAID (2012). Each commodity or method shown in this table was observed to be available in the service area or location where commodities are stored, and at least one of the observed commodities or methods was valid—that is, within expiration date.

Table 5.6: Guidelines, trained staff, and basic equipment for family planning services

Among facilities offering any modern family planning methods, the percentages having family planning guidelines, having at least one staff member recently trained on family planning service delivery, and with the indicated equipment observed to be available on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentag modern fa | e of facilities o mily planning a | ffering any Ind having: | | Equipment | | | | | | | Number of facilities |
|--|--|---|--|--|--|--|--|--|--|--|--|--|
| Background characteristic | Guidelines on family planning ¹ | Staff trained in family planning during past 24 months ² | Staff trained in family planning at any time ² | Blood pressure apparatus ³ | Examination light | Examination bed or couch | Samples of family planning methods | Pelvic model for IUCD ⁴ | Model for showing condom use | Other family planning- specific visual aid ⁵ | offering any modern family planning methods (weighted) | offering any modern family planning methods (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 85.0 74.1 75.5 88.8 | 42.3 43.1 57.8 38.6 | 87.8 81.0 94.8 87.1 | 96.7 84.5 96.8 98.4 | 81.3 60.3 71.5 86.6 | 92.8 86.2 89.5 94.5 | 83.7 63.8 80.8 87.2 | 66.1 20.7 71.6 71.3 | 71.7 31.0 73.4 77.1 | 89.8 81.0 87.2 91.6 | 40 4 7 29 | 307 58 95 154 |
| Union-level public facilities UHFWC USC/RD | 68.1 70.8 54.5 | 24.8 26.3 16.9 | 74.8 77.7 59.7 | 88.8 89.6 84.7 | 66.3 66.1 67.4 | 83.9 86.7 69.5 | 65.5 67.3 56.2 | 52.2 54.2 42.4 | 50.3 52.9 36.8 | 68.6 71.1 56.4 | 258 216 42 | 350 280 70 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 27.2 77.0 7.0 | 16.2 45.3 20.4 | 53.4 76.8 38.7 | 76.7 96.5 100.0 | 46.4 89.0 88.7 | 64.1 88.2 96.2 | 16.9 71.3 22.4 | 3.3 59.0 7.9 | 5.1 61.4 4.3 | 30.9 84.2 14.1 | 816 22 46 | 400 91 49 |
| Location Urban Rural | 41.4 38.3 | 30.3 18.7 | 60.1 59.0 | 98.5 79.8 | 87.4 51.6 | 91.6 69.4 | 48.7 29.6 | 32.9 16.0 | 34.4 16.9 | 52.6 40.6 | 91 1,091 | 334 863 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 29.0 41.6 45.3 32.8 40.4 32.9 40.2 34.5 | 13.6 25.1 25.2 13.5 12.5 17.1 18.9 24.9 | 48.7 65.6 61.2 54.8 55.7 59.0 60.9 59.0 | 72.0 82.5 86.0 79.0 80.5 76.2 89.1 82.7 | 38.4 58.8 65.4 46.4 57.5 48.2 62.3 43.1 | 63.8 64.2 69.4 78.3 81.0 77.1 73.3 60.1 | 29.0 33.3 35.6 23.6 23.5 35.0 43.7 24.9 | 11.2 19.8 22.4 14.1 11.5 18.9 19.9 15.5 | 11.2 15.1 26.8 14.0 12.3 22.4 23.0 18.7 | 36.9 43.9 41.4 40.0 42.1 41.9 43.0 39.7 | 91 235 224 148 161 153 74 96 | 120 201 203 155 161 143 103 111 |
| Total | 38.3 | 19.6 | 59.1 | 81.3 | 54.3 | 71.1 | 31.0 | 17.3 | 18.2 | 41.5 | 1,182 | 1,197 |
| Total excluding CCs | 62.9 | 34.3 | 76.9 | 91.5 | 72.1 | 86.7 | 62.5 | 48.6 | 47.6 | 65.1 | 366 | 797 |
| Total public | 38.8 | 19.1 | 59.6 | 80.2 | 52.2 | 69.8 | 30.6 | 16.9 | 18.0 | 41.8 | 1,114 | 1,057 |
| Total public excluding CCs | 70.4 | 27.2 | 76.5 | 89.8 | 68.3 | 85.1 | 67.9 | 54.1 | 53.2 | 71.5 | 298 | 657 |

Note: The measures presented in the table with guidelines for FP and staff trained in FP comprise the staff and training domains, and blood pressure apparatus comprises the equipment domain, for assessing readiness to provide FP services within the health facility assessment methodology proposed by WHO and USAID (2012).

¹ National guidelines/manual or any other guidelines/instructions/job aid/checklist on FP.
 ² The facility had at least one interviewed staff member providing the service who reports in-service training in some aspect of FP. The training must involve structured sessions and does not include individual instruction that a provider might have received during routine supervision.
 ³ A functioning digital blood pressure apparatus or a manual sphygmomanometer with a stethoscope.
 ⁴ IUCD = intrauterine contraceptive device.
 ⁵ Flip charts or leaflets.

Table 5.7: Items for infection control during provision of family planning

Among facilities offering any modern family planning methods, the percentages with indicated items for infection control observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| _ | Percentage of facilities offering any modern family planning methods that have items for infection control | | | | | | | | | | |
|--|--|--|--|--|---|--|--|--|---|--|--|
| Background characteristic | Soap | Running water ¹ | Soap and running water | Alcohol- based hand disinfectant | Soap and running water or else alcohol- based hand disinfectant | Latex gloves ² | Sharps container | Waste receptacle ³ | Average items available ⁴ | Number of facilities offering any modern family planning methods (weighted) | Number of facilities offering any modern family planning methods (unweighted) |
| Facility type | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 83.6 77.6 74.9 86.5 | 86.6 84.5 82.1 88.0 | 81.0 75.9 74.9 83.2 | 81.6 81.0 72.6 83.8 | 91.6 91.4 85.2 93.1 | 91.7 86.2 86.5 93.7 | 69.9 56.9 67.6 72 3 | 68.8 67.2 63.0 70.3 | 4.8 4.5 4.5 | 40 4 7 29 | 307 58 95 |
| Union-level public facilities UHFWC USC/RD | 63.0 65.8 48.7 | 67.9 70.1 57.0 | 59.4 62.2 45.5 | 65.3 63.7 73.5 | 77.5 76.6 81.6 | 76.9 78.5 68.4 | 61.6 65.2 43.1 | 56.7 58.9 45.9 | 3.9 4.0 3.4 | 258 216 42 | 350 280 70 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 53.7 81.7 83.8 | 49.3 79.1 85.8 | 44.1 79.1 79.8 | 65.8 88.0 93.8 | 73.9 94.4 95.3 | 55.8 83.8 81.0 | 61.8 73.2 26.9 | 48.0 76.0 69.7 | 3.3 4.8 4.4 | 816 22 46 | 400 91 49 |
| Location Urban Rural | 79.0 56.7 | 80.2 54.6 | 75.7 48.7 | 89.4 65.9 | 94.3 75.0 | 87.6 61.1 | 51.6 61.7 | 69.6 50.5 | 4.6 3.5 | 91 1,091 | 334 863 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 39.2 64.9 54.7 62.0 63.1 59.1 61.1 53.5 | 41.4 57.3 56.0 61.2 62.0 58.9 65.2 44.1 | 36.9 53.1 51.9 56.4 53.7 50.3 54.2 39.8 | 50.1 78.7 72.0 64.8 74.5 55.0 61.9 65.5 | 60.0 81.4 76.0 79.9 85.3 72.5 68.2 73.4 | 61.4 63.5 73.1 66.2 56.0 56.1 71.9 52.1 | 50.5 60.5 63.3 54.4 61.1 66.9 65.3 62.7 | 39.9 54.7 45.6 55.1 50.5 52.0 65.3 58.9 | 2.8 3.6 3.6 3.7 3.5 3.9 3.4 | 91 235 224 148 161 153 74 96 | 120 201 203 155 161 143 103 111 |
| Total | 58.4 | 56.6 | 50.8 | 67.7 | 76.5 | 63.1 | 60.9 | 52.0 | 3.6 | 1,182 | 1,197 |
| Total excluding CCs | 69.0 | 72.9 | 65.5 | 72.0 | 82.2 | 79.4 | 58.8 | 60.8 | 4.1 | 366 | 797 |
| Total public | 57.0 | 55.0 | 49.0 | 66.3 | 75.4 | 62.0 | 62.0 | 50.8 | 3.5 | 1,114 | 1,057 |
| Total public excluding CCs | 65.8 | 70.4 | 62.3 | 67.5 | 79.4 | 78.9 | 62.7 | 58.3 | 4.0 | 298 | 657 |

¹ Piped water, water in bucket with specially fitted tap, or water in pour pitcher.
 ² Non-latex equivalent gloves are acceptable.
 ³ Waste receptacle with plastic bin liner.
 ⁴ Average number of items (soap, running water, alcohol-based hand disinfectant, latex gloves, sharps container, and waste receptacle) available for infection control.

Table 5.8: Readiness of health facilities to provide family planning services

Among facilities that offer any modern family planning methods, the percentages with family planning guidelines, with at least one staff member recently trained on family planning services, with the indicated contraceptive commodities available on the day of the survey, and the average readiness score by background characteristics, Bangladesh HFS 2022

| | Percentage of facilities offering any modern family planning and having: | | | | | | | | | | Number of facilities offering |
|--|--|---|---|---|------------------------------|----------------|---|--------------------------------|--|--|--|
| Background characteristic | Guidelines on family planning ¹ | Staff trained in family planning any time ² | Blood pressure apparatus ³ | Combined or progestin-only oral pills | Progestin-only injectable | Male condom | Intrauterine contraceptive device | Implant (one or two rod) | Average readiness score (out of 8) ⁴ | any modern family planning methods (weighted) | any modern family planning methods (unweighted) |
| Facility type | | | | | | | | | | | |
| District and upazila public facilities | 85.0 | 87.8 | 96.7 | 92.0 | 91.2 | 90.5 | 87.6 | 77.9 | 7.1 | 40 | 307 |
| DH | 74.1 | 81.0 | 84.5 | 79.3 | 70.7 | 72.4 | 69.0 | 37.9 | 5.7 | 4 | 58 |
| MCWC UHC | 75.5 88.8 | 94.8 87.1 | 96.8 98.4 | 89.4 94.4 | 89.4 94.4 | 88.3 93.6 | 85.3 90.7 | 66.1 86.3 | 6.9 7.3 | 7 29 | 95 154 |
| Union-level | | | | | | | | | | | |
| public facilities | 68.1 | 74.8 | 88.8 | 92.8 | 89.4 | 91.2 | 77.9 | 10.3 | 5.9 | 258 | 350 |
| | 70.8 | //./ 59.7 | 89.6 84.7 | 95.3 80.2 | 93.0 71.3 | 93.9 78.0 | 80.9 | 10.6 8.4 | 6.1 5.0 | 216 42 | 280 |
| 000/110 | 04.0 | 55.7 | 04.7 | 00.2 | 71.0 | 70.0 | 02.0 | 0.4 | 0.0 | 74 | 10 |
| Community clinic | | | | | | | | | | | |
| (CC) NGO clinic/ | 27.2 | 53.4 | 76.7 | 86.5 | 55.3 | 74.3 | - | - | 3.8 | 816 | 400 |
| hospital Private hospital | 77.0 7.0 | 76.8 38.7 | 96.5 100.0 | 74.3 34.3 | 66.2 1.6 | 71.9 32.0 | 50.2 2.6 | 31.1 1.6 | 5.4 2.2 | 22 46 | 91 49 |
| Location | | | | | | | | | | | |
| Urban | 41.4 | 60.1 | 98.5 | 61.7 | 44.6 | 59.6 | 38.8 | 29.3 | 4.3 | 91 | 334 |
| Rural | 38.0 | 59.0 | 79.8 | 87.8 | 63.5 | 78.3 | 21.9 | 3.9 | 4.3 | 1,091 | 863 |
| Division | | | | | | | | | | | |
| Barishal | 29.0 | 48.7 | 72.0 | 94.2 | 76.6 | 77.0 | 24.9 | 3.7 | 4.3 | 91 | 120 |
| Chattogram | 41.6 | 65.6 | 82.5 | 83.6 | 62.7 | 80.3 | 23.6 | 9.3 | 4.5 | 235 | 201 |
| Khulna | 40.3 | 01.2 54.8 | 86.0 79.0 | 04.0 01.5 | 62.0 | 79.9 | 20.0 | 0.2 3.1 | 4.5 1 3 | 224 | 203 |
| Raishahi | 40.4 | 55.7 | 80.5 | 78.8 | 55.9 | 68.2 | 20.8 | 4.2 | 4.0 | 161 | 161 |
| Rangpur | 32.9 | 59.0 | 76.2 | 85.1 | 59.7 | 69.4 | 19.1 | 3.6 | 4.0 | 153 | 143 |
| Sylhet | 40.2 | 60.9 | 89.1 | 90.6 | 62.7 | 88.0 | 29.1 | 7.7 | 4.7 | 74 | 103 |
| Mymensingh | 34.5 | 59.0 | 82.7 | 86.0 | 58.7 | 71.3 | 17.7 | 3.1 | 4.1 | 96 | 111 |
| Total | 38.3 | 59.1 | 81.3 | 85.8 | 62.1 | 76.9 | 23.2 | 5.9 | 4.3 | 1,182 | 1,197 |
| Total excluding CCs | 62.9 | 76.9 | 91.5 | 84.3 | 77.3 | 82.6 | 67.9 | 17.8 | 5.6 | 366 | 797 |
| Total public | 38.8 | 59.6 | 80.2 | 88.2 | 64.5 | 78.8 | 23.5 | 5.5 | 4.4 | 1,114 | 1,057 |
| Total public excluding CCs | 70.4 | 76.5 | 89.8 | 92.7 | 89.7 | 91.1 | 79.2 | 19.4 | 6.1 | 298 | 657 |

Note: The measures presented in the table on guidelines for FP and staff trained in FP comprise the staff and training domains, and blood pressure apparatus the equipment domain, for assessing readiness to provide FP services within the health facility assessment methodology proposed by WHO and USAID (2012). "-" Means the specific FP methods are not provided at this type of facility.

¹ National guidelines/manual or any other guidelines/instructions/job aid/checklist on FP.

² The facility had at least one interviewed staff member providing the service who reports receiving in-service training in some aspect of FP. The training must involve structured sessions and does not include individual instruction that a provider might have received during routine supervision. ³ A functioning digital blood pressure apparatus or a manual sphygmomanometer with a stethoscope.

⁴ Average readiness score is the average number of 8 items (guidelines on FP, staff trained in FP any time, blood pressure apparatus, combined or progestin-only oral pills, progestin-only injectable, male condom, intrauterine contraceptive device, and implant) available for providing FP services.

Table 5.9: Supportive management for providers of family planning services

Among interviewed family planning service providers, the percentage who report receiving training related to their work and personal supervision during the specified time periods, by background characteristics, Bangladesh HFS 2022

| | Percentage of | f interviewed providers | | | |
|--|-----------------------------------|-----------------------------------|------------------------|-----------------------|-----------------------|
| | | | Training related to | | |
| | | | family planning during | Number of | Number of |
| | Training related to | | the 24 months and | interviewed providers | interviewed providers |
| | family planning during | Personal supervision | personal supervision | of family planning | of family planning |
| Background | the 24 months | during the 6 months | during the 6 months | services | services |
| characteristic | preceding the survey ¹ | preceding the survey ² | preceding the survey | (weighted) | (unweighted) |
| Facility type | | | | | |
| District and upazila public facilities | 16.7 | 97.5 | 16.4 | 540 | 1,277 |
| DH | 22.5 | 96.2 | 22.5 | 129 | 219 |
| MCWC | 27.6 | 97.7 | 27.6 | 23 | 314 |
| UHC | 14.2 | 97.9 | 13.7 | 388 | 744 |
| Union-level public facilities | 21.0 | 88.6 | 17.2 | 285 | 506 |
| UHFWC | 21.7 | 87.9 | 17.3 | 244 | 406 |
| USC/RD | 16.6 | 92.6 | 16.4 | 41 | 100 |
| Community clinic (CC) | 15.4 | 90.9 | 14.6 | 949 | 513 |
| NGO clinic/hospital | 24.1 | 96.7 | 24.1 | 65 | 229 |
| Private hospital | 16.6 | 85.1 | 15.7 | 153 | 94 |
| Location | | | | | |
| Urban | 17.6 | 94.3 | 17.4 | 566 | 1.170 |
| Rural | 16.6 | 91.2 | 15.3 | 1,426 | 1,449 |
| Division | | | | | |
| Barishal | 13.0 | 86.7 | 11.1 | 162 | 272 |
| Chattogram | 22.8 | 93.2 | 21.8 | 369 | 446 |
| Dhaka | 23.2 | 94.2 | 22.5 | 414 | 435 |
| Khulna | 8.8 | 86.8 | 6.3 | 215 | 301 |
| Rajshahi | 9.9 | 91.7 | 9.9 | 286 | 375 |
| Rangpur | 11.5 | 95.0 | 11.1 | 265 | 340 |
| Sylhet | 21.1 | 95.5 | 20.6 | 112 | 226 |
| Mymensingh | 20.4 | 90.0 | 17.8 | 169 | 224 |
| Total | 16.9 | 92.1 | 15.9 | 1,992 | 2,619 |
| Total excluding CCs | 18.3 | 93.2 | 17.0 | 1,043 | 2,106 |
| Total public | 16.3 | 92.5 | 15.6 | 1,774 | 2,296 |
| Total public excluding CCs | 18.2 | 94.4 | 16.7 | 825 | 1,783 |

¹ Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might ¹ Personal supervision refers to any form of technical support or supervision from a facility-based supervisor or from a visiting supervisor. It may include, but

is not limited to, review of records and observation of work, with or without any feedback to the health worker.

Table 5.10: Training for family planning service providers

Among interviewed family planning service providers, the percentages who report receiving in-service training on topics related to family planning during the specified time periods preceding the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of providers of FP services who report receiving in-service training on: | | | | | | | | | | | Number of Number of | Number of | | | | | | | |
|--|---|--|---|--|--|--|---|---|--|--|--|--|--|--|--|--|---|--|---|--|
| | Counselir | ng on FP | FP-relate issu | d clinical es ¹ | Insertion of IL | /removal JCD | Insertion of Im | /removal plant | FP for HI | /+ clients | Postpar | tum FP | Vased | ctomy | Tubal li | gation | Emerg | gency ception | interviewed providers of family | providers of family |
| Background characteristic | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | planning services (weighted) | planning services (unweighted) |
| Facility type | | | | | | | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 11.5 18.1 12.2 9.3 | 38.0 43.5 64.7 34.5 | 6.4 9.8 11.0 4.9 | 24.5 30.9 46.2 21.0 | 8.5 10.9 15.1 7.3 | 32.4 40.7 62.3 27.8 | 4.7 7.2 5.3 3.8 | 16.4 18.2 30.7 15.0 | 4.3 7.9 4.1 3.1 | 12.3 18.4 26.1 9.4 | 9.6 14.7 18.4 7.3 | 29.8 39.5 63.0 24.5 | 3.3 4.9 5.2 2.7 | 11.4 12.8 27.4 9.9 | 3.4 4.6 5.2 2.8 | 12.0 14.2 28.4 10.3 | 7.2 12.1 7.3 5.6 | 24.2 33.1 51.0 19.6 | 540 129 23 388 | 1,277 219 314 744 |
| Union-level public facilities UHFWC USC/RD | 11.4 11.3 12.3 | 58.2 58.6 55.6 | 5.1 5.0 5.7 | 36.0 37.9 24.4 | 4.8 4.6 6.2 | 37.5 38.5 31.8 | 1.7 1.0 5.3 | 13.7 12.8 18.7 | 3.5 3.6 3.1 | 17.3 18.5 10.4 | 7.1 6.7 9.6 | 45.5 46.9 37.2 | 1.0 0.8 2.2 | 8.5 8.1 10.6 | 0.7 0.6 1.0 | 9.4 9.3 9.6 | 8.5 8.9 5.8 | 38.2 39.2 32.6 | 285 244 41 | 506 406 100 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 11.9 22.5 15.6 | 46.3 49.2 28.9 | 5.3 14.9 8.9 | 21.8 36.1 15.0 | 1.7 14.1 8.8 | 11.5 40.0 19.0 | 2.2 13.2 6.1 | 11.2 33.4 15.4 | 3.5 9.2 5.7 | 13.7 21.9 15.1 | 5.8 15.8 10.0 | 24.6 38.3 17.3 | 1.6 8.3 1.2 | 9.5 19.4 5.0 | 1.6 8.4 8.6 | 10.2 19.5 18.7 | 4.4 15.8 11.3 | 22.6 40.4 19.1 | 949 65 153 | 513 229 94 |
| Location Urban Rural | 13.8 11.8 | 37.2 47.4 | 7.9 5.4 | 24.6 24.5 | 9.2 3.3 | 31.5 18.8 | 5.7 2.6 | 18.2 12.3 | 5.3 3.6 | 14.6 14.0 | 10.6 6.4 | 28.6 29.0 | 3.1 1.8 | 10.8 9.4 | 5.1 1.7 | 14.9 10.2 | 8.8 5.8 | 25.4 25.6 | 566 1,426 | 1,170 1,449 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 7.5 16.8 18.0 4.8 7.4 8.9 18.2 13.1 | 35.3 57.9 42.8 36.5 42.8 38.8 52.3 44.7 | 2.0 9.6 11.8 1.9 1.4 3.7 8.8 4.0 | 13.2 33.8 22.3 19.4 23.9 26.0 29.9 22.0 | 4.0 6.1 7.5 2.1 2.5 2.9 7.2 6.7 | 18.0 29.0 22.1 22.1 20.7 18.7 25.4 19.8 | 0.6 5.0 1.2 1.7 2.1 4.1 5.3 | 8.6 18.4 13.7 10.9 15.8 10.1 16.3 15.7 | 1.2 6.9 5.1 0.8 1.7 4.2 7.0 3.9 | 8.2 21.7 13.2 12.2 9.0 15.2 19.7 11.9 | 7.0 13.2 9.3 3.3 2.0 5.0 8.5 10.5 | 20.0 36.7 24.2 30.5 29.8 24.9 34.1 30.6 | 0.2 5.4 2.2 0.8 0.2 1.8 2.9 2.0 | 8.1 12.8 6.9 6.6 13.0 10.3 11.4 9.1 | 0.3 5.3 4.4 0.9 0.2 1.8 4.0 2.1 | 8.4 15.0 10.1 10.6 13.0 11.2 12.3 9.3 | 3.0 11.2 9.9 4.4 2.3 4.3 9.4 4.1 | 14.8 34.9 26.3 23.5 23.7 25.7 27.1 18.1 | 162 369 414 215 286 265 112 169 | 272 446 435 301 375 340 226 224 |
| Total | 12.4 | 44.5 | 6.1 | 24.5 | 5.0 | 22.4 | 3.5 | 14.0 | 4.1 | 14.2 | 7.6 | 28.9 | 2.1 | 9.8 | 2.7 | 11.5 | 6.6 | 25.6 | 1,992 | 2,619 |
| Total excluding CCs | 12.8 | 42.9 | 6.9 | 26.9 | 7.9 | 32.3 | 4.6 | 16.6 | 4.6 | 14.6 | 9.3 | 32.8 | 2.7 | 10.2 | 3.7 | 12.7 | 8.7 | 28.3 | 1,043 | 2,106 |
| Total public | 11.7 | 45.7 | 5.6 | 24.9 | 4.3 | 22.0 | 2.9 | 13.2 | 3.7 | 13.8 | 7.1 | 29.5 | 2.0 | 9.9 | 2.0 | 10.6 | 5.9 | 25.6 | 1,774 | 2,296 |
| Total public excluding CCS | 11.5 | 45.0 | 5.9 | 28.4 | 7.2 | 34.1 | 3.6 | 15.5 | 4.0 | 14.0 | 8.7 | 35.2 | 2.5 | 10.4 | 2.4 | 11.1 | 7.7 | 29.0 | 825 | 1,783 |

Note: Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision. IUCD = Intrauterine contraceptive device

¹ Any training on clinical management of family planning methods, including managing side effects.

Table 5.11: Key indicators for family planning services in MCWCs

Among facilities offering any modern family planning methods, the percentages with key indicators, Bangladesh HFS 2022

| | Facility type | | | | | | |
|--|--|--|--|--|--|--|--|
| Indicators | MCWC (district-level) | MCWC (upazila and union-level) | | | | | |
| Availability | | ` | | | | | |
| Provide any modern FP (including an emergency contraceptive) ¹ Provide any long-acting and permanent methods ² Provide male or female sterilization ³ Number of facilities (weighted) | 96.8 96.8 82.4 5 | 89.5 81.7 41.5 3 | | | | | |
| Number of facilities (unweighted) | 62 | 38 | | | | | |
| Readiness Guideline on family planning ⁴ Staff trained in family planning at any time ⁵ Blood pressure apparatus ⁶ Combined or progestin-only oral pills Progestin-only injectables Male condoms IUCD Implant | 82.9 98.5 98.3 86.6 86.6 84.9 86.6 84.9 | 62.6 88.6 94.4 94.2 94.2 94.2 83.0 33.7 | | | | | |
| Average readiness score ⁷ | 7.1 | 6.4 | | | | | |
| Number of facilities offering any modern family planning method (weighted) Number of facilities offering any modern family | 4 | 3 | | | | | |
| planning method (unweighted) | 60 | 35 | | | | | |

¹ Facility provides any of the following: contraceptive pills (combined or progestin-only), injectables (progestin-only), one-rod implant, two-rod implant (zadell), IUCDs, male condoms, female sterilization (tubal ligation), or male sterilization (vasectomy), and ² Facility provides any of the following long-term and permanent FP methods: one-rod

implant, two-rod implant, IUCDs, female sterilization (tubal ligation), or male sterilization (vasectomy).

 ³ Perform male or female sterilization in the facility.
 ⁴ National guidelines/manual or any other guidelines/instructions/job aid/checklist on FP.

^{FF}. ⁵ The facility had at least one interviewed staff member providing the service who reports receiving in-service training in some aspect of FP. The training must involve structured sessions and does not include individual instruction that a provider might

have received during routine supervision. ⁶ A functioning digital blood pressure apparatus, or a manual sphygmomanometer with a stethoscope. ⁷ Average readiness score is the average number of items (out of 8 items) available for

providing FP services.

ANTENATAL CARE SERVICES

Key Findings

- The availability of ANC services is universal in public health facilities (100%). All DHs, UHCs, UHFWCs, CCs, and NGOs offer ANC services (100% each), whereas the service is available in seven in ten private facilities (72%) (**Table 6.1** and **Figure 6.1**).
- Overall, the availability of ANC services in public health facilities providing ANC services remained almost unchanged in the last five years. However, the availability of ANC services in private facilities has experienced a notable decrease, dropping 23 percentage points between 2017 and 2022 (**Figure 6.1**).
- One-third (34%) of total public health facilities providing ANC services had guidelines available on the day of survey (**Table 6.2** and **Figure 6.2**).
- Two-thirds (67%) of public health facilities that offer ANC have at least one staff with in-service ANC training at any time, which has increased by 13 percentage points between 2017 and 2022 (**Table 6.2** and **Figure 6.3**).
- Seventeen percent of ANC providers among public facilities received ANC training during the 24 months preceding the survey (Table 6.7). Around four in ten providers in public facilities ever received in-service training on ANC counseling (45%), pregnancy complications (42%), and family planning (38%) (Table 6.8).
- The availability of blood pressure apparatus is nearly universal in all public health facilities excluding CCs (93%), private (91%), and NGO facilities (99%). Blood pressure apparatus is available in almost all of the DHs (100%), UHCs (99%), and MCWCs (96%) (**Table 6.2** and **Figure 6.4**).
- Among the facilities offering ANC, almost one-third of public facilities excluding CCs have the capacity to conduct tests for hemoglobin (32%), whereas every four in ten have urine protein (40%) and urine glucose (39%) tests available (**Table 6.3**).
- Total hemoglobin testing capacity has increased in public health facilities (excluding CCs) from 23% in 2017 to 32% in 2022. The availability of urine protein test is almost universal in DHs (89%). Moreover, most UHCs (80%), NGOs (84%), and private facilities (80%) have the capacity to perform urine protein test **(Table 6.3** and **Figures 6.6-6.7).**
- Every 9 in 10 public facilities offering ANC services have essential medicines (iron, folic acid, or combined iron and folic acid tablets) for ANC available. The availability of iron or folic acid tablets for ANC is universal (99%) in MCWCs and UHFWCs, whereas it is relatively lower in private facilities (60%) (**Table 6.4** and **Figure 6.9**).
- The average readiness score for ANC items (guidelines, trained staff, blood pressure apparatus, hemoglobin test, urine protein test, and iron or folic acid tablets) in total public facilities is 3.0 (out of 6 items) in 2022. The ANC readiness score is 4.8 for NGOs and 3.4 for private facilities (**Table 6.6** and **Figure 6.11**).

6.1 BACKGROUND

ntenatal care (ANC) is the gateway for many critical maternal, newborn, and child health care services. ANC is also often the first opportunity for a woman and her family to interact with the health system. The provision of ANC by a medically trained provider is intended to monitor the status of a pregnancy, identify the complications associated with the pregnancy, and prevent adverse pregnancy outcomes. For the most effective outcomes, there should be regular ANC throughout pregnancy.

The results of the 2022 Bangladesh Demographic and Health Survey (BDHS) showed almost universal coverage (88%) in the percentage of women age 15-49 with a live birth in the 2 years preceding the survey received at least one ANC visit with a medical provider. It was reported 82% in the 2017 BDHS and 65% in the 2014 BDHS. However, between 2017 and 2022, the proportion of women who had the recommended four or more ANC visits in the 2 years preceding the survey during their most recent pregnancy decreased from 46% to 41%. Therefore, Bangladesh needs more attention to evaluate the situation of the 4th Health, Population and Nutrition Sector Program's target of 50% of pregnant women completing at least four ANC visits by 2022 (Ministry of Health and Family Welfare [MOHFW], January 2017- June 2024).

Data from the 2022 BHFS related to delivery of ANC services will be useful in the country's continuing efforts to improve ANC care. This chapter explores the results of the 2022 BHFS in the following key areas related to provision of ANC services at health facilities in Bangladesh:

- Availability of ANC services: Section 6.2, including Tables 6.1–6.5 and Figures 6.1–6.10, examines the availability of ANC services at health facilities, including the availability of basic amenities and equipment, diagnostic capacity, essential medicines, and infection control processes.
- Readiness of health facilities to provide ANC services: Section 6.3, including Table 6.6 and Figure 6.11, addresses the average readiness score of facilities to provide quality ANC services.
- Basic management and administrative systems: Section 6.4, including Tables 6.7 and 6.8, considers supervision and recent in-service training of providers that support quality services.

6.2 AVAILABILITY OF ANTENATAL CARE (ANC) SERVICES

6.2.1 ANC Service Provision

- The availability of ANC services is universal in public health facilities (100%) in 2022 (**Table 6.1** and **Figure 6.1**).
- All of the DHs, UHCs, UHFWCs, CCs, and NGOs offer ANC services (100% each); and the service is available in seven in ten private facilities (72%) (**Table 6.1** and **Figure 6.1**)
- Overall, the availability of ANC services in public health facilities providing ANC services remained almost unchanged in the last five years. However, the availability of ANC services in private facilities has experienced a notable decrease, dropping 23 percentage points between 2017 and 2022 (**Table 6.1** and **Figure 6.1**).
- Eight in ten public facilities offer ANC services during all working days in a week (**Table 6.1**).
- About 94% of DHs, 93% of UHCs, 83% of CCs, 96% of NGO facilities, and 84% of private facilities offer ANC services on all working days. However, 65% of union-level public facilities offer the services every day (**Table 6.1**).



Figure 6.1 Availability of ANC services in health facilities, by facility type

6.2.2 Availability of ANC Guidelines, Trained Staff, and Equipment

The availability of ANC service guidelines, appropriately trained providers, and specific supplies and equipment all support the provision of high-quality services. Moreover, the ability to provide regularly prescribed medications and perform basic tests for diagnosis improves services.

- Around one-third of the public health facilities (34%) providing ANC had guidelines on ANC available on the day of survey. The availability of ANC guidelines is the highest (60%) in DHs, followed by NGOs (52%), while it is the lowest in private (9%) facilities (**Table 6.2** and **Figure 6.2**).
- The availability of ANC guidelines in public health facilities decreased from 49% in 2014 to 34% in 2022. A notable decline has been observed among all facilities in the last eight years regarding the availability of ANC guidelines (**Table 6.2** and **Figure 6.2**).



Figure 6.2 Availability of ANC guidelines in health facilities, by facility type Percent

- Two-thirds (67%) of public health facilities that offer ANC have at least one staff with in-service ANC training at any time, an increase of 13 percentage points in the last five years (Table 6.2 and Figure 6.3).
- MCWCs are most likely (88%) to have at least one staff person with training on ANC at any time, and private facilities are least likely (27%) to have at least one trained staff person at any time (Table 6.2 and Figure 6.3).
- The proportion of facilities offering ANC services that have at least one ever trained staff varies by region, from 55% in Sylhet to 71% in Khulna and Mymensingh (**Table 6.2**).
- Only 21% of public facilities, 15% of private facilities, and 37% of NGOs had at least one staff trained within 24 months (Table 6.2).





Percent

• The availability of blood pressure apparatus is universal in total public health facilities excluding CCs (93%), private hospitals (91%), and NGO facilities (99%). Blood pressure apparatus is available in almost all of the DHs (100%), UHCs (99%), and MCWCs (96%) (**Table 6.2** and **Figure 6.4**).

• Eighty-one percent of the CCs were observed to have a blood pressure (BP) apparatus on the day of survey, which is the lowest among all other facilities (**Table 6.2** and **Figure 6.4**).

Among the public facilities offering ANC, 84% have an adult weighing scale, 91% have a stethoscope, and 71% have an examination bed or couch. Forty-one percent of public facilities have measuring tape available and only 7% have a fetal stethoscope (Table 6.2).



Figure 6.4 Availability of blood pressure (BP) apparatus to support ANC services

- The average score of equipment out of 6 items (blood pressure apparatus, stethoscope, adult weighing scale, fetal stethoscope, measuring tape, and examination bed or couch) is 4.2 in public facilities excluding CCs, while it is 4.6 in private facilities (**Table 6.2** and **Figure 6.5**).
- The average score of equipment is relatively higher in UHCs (5.2), DHs (5.1), and NGO facilities (5.0). The score is lowest in CCs (3.6) and USC/RDs (3.7) (**Table 6.2** and **Figure 6.5**).
- The average score of equipment has increased in all health facilities except CCs and private facilities in the last five years (**Figure 6.5**).



Figure 6.5 Average score of equipment (out of 6 items) to provide ANC services, by facility type

6.2.3 Availability of Laboratory Diagnostic Tests

Performing basic laboratory tests onsite saves time for both the client and the provider. It also makes it much more likely that the client receives the test results and the provider can act on them.

- Among facilities offering ANC services, the hemoglobin test is universally available in DHs (94%) and UHCs (92%). Moreover, most NGOs (87%) and private hospitals (83%) perform hemoglobin test in the facility (Table 6.3 and Figure 6.6).
- Hemoglobin testing capacity has increased in total public health facilities (excluding CCs) from 23% in 2017 to 32% in 2022. Most facilities showed an increase in the capacity to conduct hemoglobin test for ANC services during the same period (Table 6.3 and Figure 6.6).
- Among facilities offering ANC, the availability of urine glucose testing is higher in DHs (86%), UHCs (80%), and NGO facilities (83%) (**Table 6.3**).





- The availability of urine protein tests is almost universal in DHs (89%). Moreover, most UHCs (80%), NGOs (84%), and private facilities (80%) have the capacity to perform urine protein test (Table 6.3 and Figure 6.7).
- There is a sharp increase (from 24% in 2017 to 40% in 2022) is observed in the availability of urine protein test in public health facilities (excluding CCs) offering ANC (**Table 6.3** and **Figure 6.7**).

Figure 6.7 Availability of urine protein test in health facilities to provide ANC services, by facility type



• The average score of laboratory diagnostic tests (hemoglobin, urine protein, urine glucose, and blood grouping and Rhesus factor tests) is 1.2 out of 4 tests in public facilities excluding CCs to provide ANC services (**Table 6.3** and **Figure 6.8**).

- The average score of laboratory diagnostic tests is highest in DHs (3.6), followed by UHCs (3.3) and NGO facilities (3.3) (**Table 6.3** and **Figure 6.8**).
- The average score of laboratory diagnostic tests has remained the same in total public facilities between 2017 and 2022. The average score of laboratory diagnostic tests has increased in all health facilities except CCs in the last five years (**Figure 6.8**).

Figure 6.8 Average score of laboratory diagnostic tests (out of 4 tests) to provide ANC services, by facility type



6.2.4 Availability of Medicines for Routine ANC

Pregnant women should take iron supplements and/or folic acid to combat anemia and improve pregnancy outcomes.

- Every 9 out of 10 public facilities providing ANC services had essential medicines (iron, folic acid, combined iron and folic acid, iron or folic acid tablets) available on the day of the survey. Private facilities are less likely to have these medicines than district and upazila public facilities, union-level public facilities, community clinics, and NGO facilities (**Table 6.4**).
- The availability of iron or folic acid tablets for ANC is universal in MCWCs and UHFWCs (99% each). (Table 6.4 and Figure 6.9).
- The availability of iron or folic acid tablets remains unchanged in total public health facilities since 2017 (both 93%), though the availability has decreased slightly from 2014 (96% to 93%) (**Table 6.4** and **Figure 6.9**).
- Rural facilities are more likely than urban facilities to have medicines for routine ANC. About 94% of rural facilities have iron or folic acid tablets, compared to only 67% of urban facilities (**Table 6.4**).



Figure 6.9 Availability of iron or folic acid in health facilities to provide ANC services

6.2.5 Availability of Infection Control Items

Infection control is very important to ensure the overall quality of ANC services.

- More than half of the public facilities have alcohol-based hand disinfectant (65%), latex gloves (62%), sharps container (60%), soap (55%), running water (53%), and waste receptacle (52%) on ANC service site for infection control (Table 6.5).
- The average score of infection control items (soap, running water, alcohol-based hand disinfectant, latex gloves, sharps container, and waste receptacle) is 3.6 in public facilities excluding CCs (Table 6.5 and Figure 6.10).
- The average score of infection control items is relatively highest in NGO facilities (4.9), followed by DHs and private facilities (4.2), MCHCs (4.1), and UHCs (4.0). The score is lowest in USC/RDs (2.8) (Table 6.5 and Figure 6.10).

Figure 6.10 Average score of infection control items (out of 6 items) for ANC services, by facility type



6.3 READINESS OF HEALTH FACILITIES TO PROVIDE ANC SERVICES

The WHO has identified a set of items/tracer indicators that a facility needs in order to offer quality ANC services (WHO 2013, Service Availability and Readiness Assessment [SARA]). Data from the BHFSs were used to construct a slightly less restrictive and Bangladesh-context-appropriate version of the WHO-recommended service readiness measure for ANC.

The measure requires the following six items/tracer indicators to be available for a health facility to be considered ready to offer quality ANC service:

- Trained staff:
 - At least one provider ever received in-service ANC training
- Guidelines:
 - National or other ANC guidelines at the facility
- Equipment:
 - Blood pressure apparatus
- Diagnostic capacity:
 - Hemoglobin test
 - Urine protein test
- Medicines:
 - Iron or folic acid tablets

The assessment of facilities' readiness to provide antenatal care (ANC) services is done based on the average readiness score, which is interpreted as the average number of items or tracer indicators available from the adapted six items for ANC on the day of the visit.

- The average readiness score for public facilities is 3.0 (out of 6) in 2022 (**Table 6.6** and **Figure 6.11**).
- The average readiness score is relatively higher in DHs (5.1), MCWCs (4.7), UHCs (5.2), and NGO facilities (4.8). The score is lowest in USC/RDs (2.8) and CCs (2.7) (**Table 6.6** and **Figure 6.11**).
- The average readiness score for providing ANC services in public facilities (excluding CCs) increased from 3.2 to 3.8 between 2014 and 2022 (**Table 6.6** and **Figure 6.11**).

Figure 6.11 Average readiness score of health facilities (out of 6 items) to provide ANC services, by facility type



6.4 BASIC MANAGEMENT AND ADMINISTRATIVE SYSTEMS

6.4.1 Recent In-service Training and Supervision

In-service training and personal supervision related to ANC are key elements in ensuring supportive management of staff who provide ANC services. Training is critical to ensuring that personnel have the knowledge and skills to provide quality ANC services. Personal supervision helps sustain health worker capacity because it identifies a provider's strengths and weaknesses. **Table 6.7** presents information on recent in-service training (during the 24 months preceding the survey) and recent personal supervision (during the 6 months preceding the survey) of ANC providers. Providers who have received recent in-service training can be expected to have more up-to-date knowledge about their particular service area.

• Among public facilities, 17% of ANC providers had received ANC training during the 24 months preceding the survey (**Table 6.7**).

- Twenty-two percent of ANC providers at CCs received recent in-service ANC training, along with 16% of providers at NGOs, 13% of providers at district and upazila public facilities, and 9% of providers at both union-level facilities and private hospitals (**Table 6.7**).
- Providers at rural health facilities (18%) are more likely than providers at urban facilities (12%) to have received recent ANC training. This difference can be attributed to the concentration of CCs in rural areas (Table 6.7).
- The proportion of providers received ANC training during the 24 months preceding the survey varies by region, from 27% in Mymensingh to 5% in Khulna (**Table 6.7**).
- The great majority of health workers who provide ANC services are supervised on a routine basis. About 9 in 10 providers at public facilities (98% at district and upazila facilities, 86% at union-level public facilities, and 92% at CCs) reported receiving personal supervision during the 6 months preceding the survey (**Table 6.7**).
- Ninety-seven percent of providers at NGO facilities and 90% of providers at private hospitals received recent personal supervision (**Table 6.7**).

6.4.2 In-service Training by Topic

Table 6.8 shows the percentages of providers of ANC services who reported ever receiving training on specific topics related to ANC and who reported receiving training on a given topic during the 24 months preceding the survey.

- Around four in ten providers at public facilities reported having ever received training on ANC counseling (45%), pregnancy complications (42%), and family planning (38%) (**Table 6.8**).
- One-third of providers (34%) at public facilities have received in-service training on ANC screening at any time (**Table 6.8**).
- Less than one in ten providers (9%) of public facilities have ever received training on sexually transmitted diseases (**Table 6.8**).
- The proportion of providers of ANC services who reported ever receiving training on ANC counseling, ANC screening, complications of pregnancy, family planning, and sexually transmitted infections is relatively higher among union-level public facilities than district and upazila public facilities (Table 6.8).
- The percentage of ANC service providers who received any type of ANC in-service training at any time is relatively higher in rural areas than urban areas (**Table 6.8**).

ANTENATAL CARE SERVICES

LIST OF TABLES

- Table 6.1 Availability of antenatal care services
- Table 6.2 Guidelines, trained staff, and basic equipment for antenatal care services
- Table 6.3 Diagnostic capacity
- Table 6.4 Availability of medicines for routine antenatal care
- Table 6.5 Items for infection control during provision of antenatal care
- Table 6.6 Readiness of health facilities to provide antenatal care services
- Table 6.7 Supportive management for providers of antenatal care services
- Table 6.8 Training for antenatal care service providers
- Table 6.9 Key indicators for antenatal care services in MCWCs

Table 6.1: Availability of antenatal care services

Among all facilities, the percentage offering antenatal care (ANC) services and, among facilities offering ANC services, the percentages offering the service on the indicated number of days per week, by background characteristics, Bangladesh HFS 2022

| | Percentage of | Number of | of Number of facilities | Percentage of fa ANC where AN offered on inc | acilities offering C services are licated days ¹ | Number of facilities | Number of facilities |
|--|------------------------------|--------------------------|----------------------------|--|---|----------------------------|---------------------------|
| Background characteristic | facilities that offer ANC | facilities (weighted) | facilities (unweighted) | Provides but not every day ¹ | Provides every day ¹ | offering ANC (weighted) | offering ANC (unweighted) |
| Facility type | | | | | | | |
| District and upazila public facilities | 99.8 | 41 | 319 | 9.6 | 90.4 | 41 | 318 |
| DH | 100.0 | 4 | 62 | 6.5 | 93.5 | 4 | 62 |
| MCWC | 99.0 | 7 | 100 | 22.4 | 77.6 | 7 | 99 |
| UHC | 100.0 | 30 | 157 | 7.0 | 93.0 | 30 | 157 |
| Union-level public facilities | 98.3 | 310 | 434 | 35.4 | 64.6 | 305 | 427 |
| UHFWC | 100.0 | 225 | 293 | 35.8 | 64.2 | 225 | 293 |
| USC/RD | 93.7 | 85 | 141 | 34.1 | 65.9 | 80 | 134 |
| Community clinic (CC) | 99.8 | 994 | 488 | 16.7 | 83.3 | 992 | 487 |
| NGO clinic/hospital | 100.0 | 29 | 127 | 3.6 | 96.4 | 29 | 127 |
| Private hospital | 72.4 | 183 | 189 | 16.1 | 83.9 | 132 | 141 |
| Location | | | | | | | |
| Urban | 79.3 | 235 | 514 | 13.0 | 87.0 | 187 | 467 |
| Rural | 99.3 | 1,322 | 1,043 | 21.0 | 79.0 | 1,312 | 1,033 |
| Division | | | | | | | |
| Barishal | 98.8 | 111 | 152 | 13.9 | 86.1 | 110 | 148 |
| Chattogram | 97.4 | 291 | 244 | 15.5 | 84.5 | 283 | 238 |
| Dhaka Khuda a | 95.4 | 317 | 264 | 23.3 | /6./ | 303 | 254 |
| Knuina | 90.5 | 210 | 108 | 12.3 | 86.8 | 190 | 197 |
| Rajshani Pangnur | 97.8 | 214 | 212 | 24.5 | 75.5 75.2 | 209 | 205 |
| Sylbet | 97.7 | 00 | 136 | 24.0 | 78.0 | 07 | 132 |
| Mymensingh | 97.5 | 122 | 150 | 22.0 | 78.6 | 118 | 146 |
| Mymonolign | 01.0 | 122 | 101 | 2 | 10.0 | 110 | 110 |
| Total | 96.3 | 1,557 | 1,557 | 20.0 | 80.0 | 1,499 | 1,500 |
| Total excluding CCs | 90.1 | 563 | 1,069 | 26.4 | 73.6 | 507 | 1,013 |
| Total public | 99.5 | 1,345 | 1,241 | 20.7 | 79.3 | 1,338 | 1,232 |
| Total public excluding CCs | 98.5 | 351 | 753 | 32.3 | 67.7 | 346 | 745 |
| 1 | | | | | | | |

¹ Every day refers to all working days when the facility is open

Table 6.2: Guidelines, trained staff, and basic equipment for antenatal care services

Among facilities offering antenatal care (ANC) services, the percentage having guidelines, at least one staff member trained on ANC service delivery, and the indicated equipment observed to be available on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of facilities offering ANC that have: | | | Equipment | | | | | | | | |
|--|---|---|---|---|-----------------------------|-----------------------------|----------------------------|-----------------------------|----------------------------------|--|---|---|
| Background characteristic | Guidelines on ANC ¹ | Staff trained for ANC during the past 24 months ² | Staff trained for ANC at any time ² | Blood pressure apparatus ³ | Stetho- scope | Adult weighing scale | Fetal stetho- scope | Measuring tape⁴ | Exami- nation bed or couch | Average equipment score (out of 6) ⁵ | Number of facilities offering ANC (weighted) | Number of facilities offering ANC (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public facilities DH | 47.8 59.7 | 42.0 43.5 | 82.8 80.6 | 98.7 100.0 | 99.7 100.0 | 94.5 98.4 | 45.3 33.9 | 77.6 75.8 | 98.3 98.4 | 5.1 5.1 4 7 | 41 4 7 | 318 62 |
| UHC | 43.4 | 44.1 | 82.0 | 90.0 99.1 | 100.0 | 94.4 | 48.4 | 83.8 | 99.1 | 5.2 | 30 | 157 |
| Union-level public facilities UHFWC USC/RD | 30.9 35.8 17.0 | 11.2 10.6 13.0 | 57.2 60.4 48.1 | 92.2 92.6 91.0 | 96.7 96.7 96.7 | 83.7 89.3 67.9 | 14.0 16.0 8.1 | 43.9 47.1 35.0 | 82.4 86.7 70.2 | 4.1 4.3 3.7 | 305 225 80 | 427 293 134 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 33.8 52.4 9.2 | 22.4 37.2 14.6 | 69.9 65.1 27.4 | 80.6 98.6 91.1 | 88.4 99.7 91.1 | 83.1 96.8 87.2 | 3.7 43.1 49.3 | 38.2 69.5 49.5 | 66.7 96.1 92.4 | 3.6 5.0 4.6 | 992 29 132 | 487 127 141 |
| Location Urban Rural | 19.6 33.5 | 20.5 20.2 | 41.0 67.1 | 93.6 83.5 | 93.6 90.5 | 88.5 83.6 | 45.3 7.0 | 54.8 40.2 | 93.0 70.9 | 4.7 3.8 | 187 1,312 | 467 1,033 |
| Division Barishal | 25 1 | 21.9 | 67.5 | 78 1 | 89.5 | 78.9 | 10 7 | 37.6 | 67 4 | 3.6 | 110 | 148 |
| Chattogram Dhaka Khulna | 36.0 32.0 27.1 | 23.6 17.3 13.0 | 67.8 57.5 70.5 | 83.4 87.6 84.8 | 89.7 94.5 92.2 | 82.1 86.2 84.6 | 16.9 13.3 8.0 | 42.7 47.4 41.1 | 66.5 70.2 84.0 | 3.8 3.9 3.9 | 283 303 190 | 238 254 197 |
| Rajshahi Rangpur Sylhet | 32.1 24.1 37.5 | 15.3 24.5 20.0 | 58.2 65.5 55.2 | 88.9 79.5 87.6 | 91.2 87.5 90.7 | 90.8 84.6 87.4 | 11.4 7.4 13.5 | 37.2 39.3 53.9 | 81.4 79.8 79.2 | 4.0 3.8 4.1 | 209 189 97 | 205 180 132 |
| Mymensingh | 41.5 | 32.3 | 70.6 | 85.1 | 88.1 | 73.6 | 8.2 | 35.6 | 60.4 | 3.5 | 118 | 146 |
| Total | 31.8 | 20.3 | 63.8 | 84.7 | 90.9 | 84.2 | 11.7 | 42.0 | 73.6 | 3.9 | 1,499 | 1,500 |
| Total excluding CCs | 27.8 | 16.1 | 51.9 | 92.8 | 95.6 | 86.3 | 27.4 | 49.6 | 87.1 | 4.4 | 507 | 1,013 |
| Total public | 33.5 | 20.5 | 67.4 | 83.8 | 90.6 | 83.6 | 7.3 | 40.7 | 71.3 | 3.8 | 1,338 | 1,232 |
| Total public excluding CCs | 32.9 | 14.9 | 60.2 | 93.0 | 97.0 | 85.0 | 17.7 | 47.9 | 84.3 | 4.2 | 346 | 745 |

Note: The guidelines for ANC and staff trained in ANC comprise the training domain and the blood pressure apparatus indicator comprises the equipment domain for assessing readiness to provide ANC services within the health facility assessment methodology proposed by WHO and USAID (2012). ¹ National ANC guidelines/protocol/manual or other guidelines/protocol/manual relevant to ANC ² Facility has at least one interviewed staff member providing ANC services who reports receiving in-service training in any of the following: ANC screening, counseling, nutritional

assessment of pregnant women, management of pre-eclampsia/eclampsia, and antenatal corticosteroid for threatened preterm labor. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision

³ Functioning digital blood pressure apparatus or else a functioning manual sphygmomanometer and a stethoscope

⁴ For measuring fundal height

⁵ Average number of equipment (blood pressure apparatus, stethoscope, adult weighing scale, fetal stethoscope, measuring tape, examination bed or couch) available for offering ANC services
Table 6.3: Diagnostic capacity

Among facilities offering antenatal care (ANC) services, the percentages with the capacity to conduct the indicated tests in the facility, by background characteristics, Bangladesh HFS 2022

| | Percentage of | facilities offering | ANC that have t | _ | | | |
|--|---|--|--|---|---|---|---|
| Background characteristic | Hemoglobin ¹ | Urine protein ² | Urine glucose ³ | Blood grouping and Rhesus factor ⁴ | Average number of tests available (out of 4) ⁵ | Number of facilities offering ANC (weighted) | Number of facilities offering ANC (unweighted) |
| Facility type | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 86.5 93.5 59.4 92.0 | 76.9 88.7 59.0 79.5 | 76.8 85.5 59.9 79.5 | 69.3 97.8 6.0 80.5 | 3.1 3.6 1.8 3.3 | 41 4 7 30 | 318 62 99 157 |
| Union-level public facilities UHFWC USC/RD | 24.3 28.8 11.5 | 35.2 43.5 11.5 | 33.5 41.6 10.6 | - - - | 0.9 1.1 0.3 | 305 225 80 | 427 293 134 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 4.2 87.0 83.4 | 2.7 84.4 80.1 | 2.9 83.3 79.4 | - 77.3 79.4 | 0.1 3.3 3.2 | 992 29 133 | 487 127 141 |
| Location Urban Rural | 81.5 10.3 | 77.2 11.6 | 76.7 11.3 | 74.8 1.3 | 3.1 0.3 | 187 1,313 | 467 1,033 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 10.9 25.1 24.5 14.4 23.0 9.2 22.7 12.7 | 12.1 24.3 25.9 15.4 20.2 12.6 24.6 13.7 | 13.0 24.8 24.5 15.5 20.3 12.6 25.3 10.9 | 7.1 14.0 15.3 8.1 12.0 5.0 7.3 5.4 | 0.4 0.9 0.5 0.8 0.4 0.8 | 110 283 303 190 209 189 97 118 | 148 238 254 197 205 180 132 146 |
| Total | 19.1 | 19.7 | 19.5 | 10.5 | 0.7 | 1,499 | 1,500 |
| Total excluding CCs Total public Total public excluding CCs | 48.4 11.3 31.7 | 53.1 12.4 40.2 | 51.9 12.1 38.7 | 30.9 2.2 8.5 | 1.8 0.4 1.2 | 507 1,338 346 | 1,013 1,232 745 |

"-" The tests are not offered at union-level public facilities or at community clinics

Note: The hemoglobin and urine protein measures presented in the table comprise the diagnostics domain for assessing readiness to provide ANC services within the health facility assessment methodology proposed by WHO and USAID (2012).

¹ Capacity to conduct any hemoglobin test in the facility

² Dip sticks for urine protein

³ Dip sticks for urine glucose

⁴ Anti-A, anti-B, and anti-D reagents; and glass slides ⁵ Average number of tests (hemoglobin, urine protein, urine glucose, and blood grouping and Rhesus factor) available for offering ANC services

Table 6.4: Availability of medicines for routine antenatal care

Among facilities offering antenatal care (ANC) services, percentages with essential medicines for ANC observed to be available on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage o | f facilities offering Al | | | | |
|----------------------------|--------------|--------------------------|------------------------------|----------------------------|--|--|
| Background characteristic | Iron tablets | Folic acid tablets | Combined iron and folic acid | Iron or folic acid tablets | Number of facilities offering ANC (weighted) | Number of facilities offering ANC (unweighted) |
| Facility type | | | | | | |
| District and upazila | | | | | | |
| public facilities | 85.4 | 85.2 | 80.2 | 86.3 | 41 | 318 |
| DH | 79.0 | 75.8 | 69.4 | 80.6 | 4 | 62 |
| MCWC | 94.7 | 98.0 | 87.5 | 99.0 | 7 | 99 |
| UHC | 84.1 | 83.6 | 80.1 | 84.1 | 30 | 157 |
| Union-level public | | | | | | 427 |
| facilities | 94.3 | 96.0 | 90.6 | 96.4 | 305 | |
| UHFWC | 96.0 | 98.9 | 91.8 | 98.9 | 225 | 293 |
| USC/RD | 89.2 | 87.7 | 87.1 | 89.2 | 80 | 134 |
| Community clinic (CC) | 91.6 | 92.5 | 89.4 | 92.6 | 992 | 487 |
| NGO clinic/hospital | 82.0 | 84.2 | 78.7 | 85.5 | 29 | 127 |
| Private hospital | 51.1 | 60.3 | 50.8 | 60.3 | 133 | 141 |
| Location | | | | | | |
| Urban | 59.5 | 66.4 | 58.0 | 66.7 | 187 | 467 |
| Rural | 92.3 | 93.3 | 89.7 | 93.6 | 1,313 | 1,033 |
| Division | | | | | | |
| Barishal | 98.4 | 98.3 | 95.8 | 98.4 | 110 | 148 |
| Chattogram | 88.1 | 91.6 | 88.0 | 91.6 | 283 | 238 |
| Dhaka | 86.5 | 87.8 | 84.3 | 88.3 | 303 | 254 |
| Khulna | 87.5 | 88.8 | 86.5 | 88.8 | 190 | 197 |
| Rajshahi | 81.2 | 85.3 | 74.8 | 85.3 | 209 | 205 |
| Rangpur | 90.0 | 90.0 | 86.6 | 90.0 | 189 | 180 |
| Sylhet | 90.2 | 91.4 | 87.1 | 91.8 | 97 | 132 |
| Mymensingh | 92.5 | 92.8 | 90.3 | 94.4 | 118 | 146 |
| Total | 88.2 | 90.0 | 85.8 | 90.2 | 1,499 | 1,500 |
| Total excluding CCs | 81.6 | 85.1 | 78.7 | 85.5 | 507 | 1,013 |
| Total public | 92.0 | 93.0 | 89.4 | 93.3 | 1,338 | 1,232 |
| Total public excluding CCs | 93.2 | 94.7 | 89.3 | 95.2 | 346 | 745 |

Note: The medicines presented in the table comprise the medicines and commodities domain for assessing readiness to provide ANC services within the health facility assessment methodology proposed by WHO and USAID (2012).

Table 6.5: Items for infection control during provision of antenatal care

Among facilities offering antenatal care (ANC) services, the percentages with indicated items for infection control observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of facilities offering ANC that have items for infection control | | | | | | | | | | |
|----------------------------|---|--------------------|--------------|--------------|---------------|---------------------|--------------|-------------------------|-------------------------|------------|--------------|
| | | | | | Soap and | | | | | | |
| | | | | | running water | | | | | | |
| | | | Coop and | Alcohol- | or else | | | | Avorago itomo | Number of | Number of |
| | | Running | supping | based | hased hand | Latov | Sharne | Wasto | available | | facilities |
| Background characteristic | Soap | water ¹ | water | disinfectant | disinfectant | gloves ² | container | receptacle ³ | (out of 6) ⁴ | (weighted) | (unweighted) |
| Facility type | | | | | | | | | | | |
| District and upazila | | | | | | | | | | | |
| public facilities | 71.8 | 80.2 | 69.2 | 83.5 | 91.3 | 74.5 | 34.5 | 61.0 | 4.1 | 41 | 318318 |
| DH | 77.4 | 79.0 | 75.8 | 88.7 | 95.2 | 72.6 | 35.5 | 62.9 | 4.2 | 4 | 62 |
| MCWC | 68.7 | 78.6 | 66.6 | 72.6 | 82.8 | 80.9 | 57.0 | 55.7 | 4.1 | 7 | .99 |
| UHC | 71.7 | 80.8 | 68.8 | 85.3 | 92.7 | 73.2 | 29.0 | 62.0 | 4.0 | 30 | 157 |
| Union-level public | | | | | | | | | | | |
| facilities | 54.2 | 58.1 | 50.8 | 63.8 | 74.2 | 72.3 | 56.4 | 52.2 | 3.6 | 305 | 427427 |
| UHFWC | 59.9 | 62.6 | 56.1 | 62.0 | 74.2 | 79.5 | 62.8 | 56.8 | 3.8 | 225 | 293 |
| USC/RD | 37.9 | 45.4 | 35.7 | 68.9 | 74.1 | 52.1 | 38.2 | 39.2 | 2.8 | 80 | 134 |
| Community clinic (CC) | 54.1 | 50.2 | 45.2 | 65.1 | 73.5 | 58.4 | 61.7 | 51.0 | 3.4 | 992 | 487 |
| NGO clinic/hospital | 84.7 | 83.2 | 82.5 | 89.3 | 94.4 | 87.0 | 64.8 | 76.4 | 4.9 | 29 | 127 |
| Private hospital | 79.7 | 83.7 | 78.7 | 82.6 | 89.2 | 78.1 | 32.4 | 62.2 | 4.2 | 133 | 141 |
| Location | | | | | | | | | | | |
| Urban | 76.6 | 80.1 | 75.2 | 82.8 | 89.0 | 80.4 | 39.5 | 62.5 | 4.2 | 187 | 467 |
| Rural | 54.7 | 52.8 | 47.2 | 65.1 | 74.1 | 61.7 | 59.8 | 51.6 | 3.5 | 1,313 | 1,033 |
| Division | | | | | | | | | | | |
| Barishal | 38.9 | 39.8 | 36.9 | 56.4 | 63.0 | 63.7 | 49.3 | 48.8 | 2.9 | 110 | 148 |
| Chattogram | 62.2 | 56.2 | 52.1 | 75.9 | 79.2 | 66.1 | 56.4 | 54.1 | 3.7 | 283 | 238 |
| Dhaka | 55.5 | 57.1 | 52.8 | 70.1 | 74.9 | 68.2 | 58.4 | 46.6 | 3.6 | 303 | 254 |
| Knuina | 63.2 | 64.9 | 58.4 | 66.2 | 81.9 | 66.4 | 52.4 | 58.6 | 3.7 | 190 | 197 |
| Rajshani Bongnur | 20.Z | 55.1 57.1 | 47.9 51.2 | 62.4 | 78.Z 75.4 | 50.3 50.1 | 58.5 60.2 | 49.0 | 3.4 | 209 | 205 |
| Sylbot | 50.7 60.2 | 66.4 | 55.2 | 64.5 | 75.4 | 74.2 | 66.6 | 50.7 | 3.0 | 07 | 132 |
| Mymensingh | 56.9 | 47.4 | 42 A | 64.5 | 73.2 | 59.6 | 57.7 | 62.7 | 3.5 | 118 | 146 |
| Wymensingn | 50.5 | 47.4 | 42.4 | 04.0 | 13.2 | 55.0 | 57.7 | 02.7 | 5.5 | 110 | 140 |
| Total | 57.4 | 56.2 | 50.7 | 67.3 | 75.9 | 64.0 | 57.3 | 53.0 | 3.6 | 1,499 | 1,500 |
| Total excluding CCs | 64.0 | 68.0 | 61.4 | 71.8 | 80.6 | 74.8 | 48.8 | 56.9 | 3.8 | 507 | 1,0131,013 |
| Total public | 54.6 | 52.9 | 47.2 | 65.3 | 74.2 | 62.1 | 59.6 | 51.6 | 3.5 | 1,338 | 1,232 |
| Total public excluding CCs | 56.3 | 60.7 | 53.0 | 66.1 | 76.2 | 72.6 | 53.8 | 53.3 | 3.6 | 346 | 745745 |

¹ Piped water, water in bucket with specially fitted tap, or water in pour pitcher

² Non-latex equivalent gloves are acceptable ³ Waste receptacle with plastic bin liner ⁴ Average number of items (Soap, running water, alcohol-based hand disinfectant, latex gloves, sharps container, and waste receptacle) available for infection control

Table 6.6: Readiness of health facilities to provide antenatal care services

Among facilities that offer antenatal care (ANC) services, the percentages with the indicated items considered important for the provision of quality ANC services, and the average readiness score by background characteristics, Bangladesh HFS 2022

| Background characteristics | Guidelines on ANC ¹ | Staff trained for ANC any time ² | Blood pressure apparatus ³ | Hemoglobin testing capacity | Urine protein testing capacity | Iron or folic acid tablets | Average readiness score (out of 6) ⁴ | Ultrasono- graphy | Number of facilities offering ANC (weighted) | Number of facilities offering ANC (unweighted) |
|-------------------------------|-----------------------------------|---|---|-----------------------------------|---|-------------------------------|--|----------------------|---|---|
| Facility type | | | | | | | | | | |
| District and upazila | | | | | | | | | | |
| public facilities | 47.8 | 82.8 | 98.7 | 86.5 | 76.9 | 86.3 | 5.1 | 10.7 | 41 | 318 |
| DH | 59.7 | 80.6 | 100.0 | 93.5 | 88.7 | 80.6 | 5.1 | 16.1 | 4 | 62 |
| MCWC | 43.4 | 87.7 | 96.0 | 59.4 | 59.0 | 99.0 | 4.7 | 4.0 | 7 | 99 |
| UHC | 47.1 | 82.0 | 99.1 | 92.0 | 79.5 | 84.1 | 5.2 | 11.5 | 30 | 157 |
| Union level public | | | | | | | | | | |
| facilities | 30.9 | 57.2 | 92.2 | 24.3 | 35.2 | 96.4 | 3.6 | - | 305 | 427 |
| UHFWC | 35.8 | 60.4 | 92.6 | 28.8 | 43.5 | 98.9 | 3.9 | - | 225 | 293 |
| USC/RD | 17.0 | 48.1 | 91.0 | 11.5 | 11.5 | 89.2 | 2.8 | - | 80 | 134 |
| Community clinic (CC) | 33.8 | 69.9 | 80.6 | 4.2 | 2.7 | 92.6 | 2.7 | - | 992 | 487 |
| NGO clinic/hospital | 52.4 | 65.1 | 98.6 | 87.0 | 84.4 | 85.5 | 4.8 | 15.2 | 29 | 127 |
| Private hospital | 9.2 | 27.4 | 91.1 | 83.4 | 80.1 | 60.3 | 3.4 | 17.2 | 133 | 141 |
| Location | | | | | | | | | | |
| Urban | 19.6 | 41.0 | 93.6 | 81.5 | 77.2 | 66.7 | 3.8 | 16.0 | 187 | 467 |
| Rural | 33.5 | 67.1 | 83.5 | 10.3 | 11.6 | 93.6 | 3.0 | 0.1 | 1,313 | 1,033 |
| Division | | | | | | | | | | |
| Barishal | 25.1 | 67.5 | 78.1 | 10.9 | 12.1 | 98.4 | 2.9 | 0.5 | 110 | 148 |
| Chattogram | 36.0 | 67.8 | 83.4 | 25.1 | 24.3 | 91.6 | 3.3 | 3.1 | 283 | 238 |
| Dhaka | 32.0 | 57.5 | 87.6 | 24.5 | 25.9 | 88.3 | 3.2 | 3.7 | 303 | 254 |
| Khulna | 27.1 | 70.5 | 84.8 | 14.4 | 15.4 | 88.8 | 3.0 | 0.3 | 190 | 197 |
| Rajshahi | 32.1 | 58.2 | 88.9 | 23.0 | 20.2 | 85.3 | 3.1 | 2.2 | 209 | 205 |
| Rangpur | 24.1 | 65.5 | 79.5 | 9.2 | 12.6 | 90.0 | 2.8 | 0.6 | 189 | 180 |
| Sylhet | 37.5 | 55.2 | 87.6 | 22.7 | 24.6 | 91.8 | 3.1 | 3.3 | 97 | 132 |
| Mymensingh | 41.5 | 70.6 | 85.1 | 12.7 | 13.7 | 94.4 | 3.0 | 1.2 | 118 | 146 |
| Total | 31.8 | 63.8 | 84.7 | 19.1 | 19.7 | 90.2 | 3.1 | 2.1 | 1,499 | 1,500 |
| Total excluding CCs | 27.8 | 51.9 | 92.8 | 48.7 | 53.1 | 85.5 | 3.8 | 6.2 | 507 | 1,013 |
| Total public | 33.5 | 67.4 | 83.8 | 11.3 | 12.4 | 93.3 | 3.0 | 2.3 | 1,338 | 1,232 |
| Total public excluding CCs | 32.9 | 60.2 | 93.0 | 31.7 | 40.2 | 95.2 | 3.8 | 1.3 | 346 | 745 |

"-" The tests are not offered at union-level facilities and community clinics

Note: The guidelines for ANC and staff trained in ANC comprise the training domain and the blood pressure apparatus indicator comprises the equipment domain, for assessing readiness to provide ANC services within the health facility assessment methodology proposed by WHO and USAID (2012).

¹ National ANC guidelines or other guidelines relevant to ANC

² Facility has at least one interviewed staff member providing ANC services who reports receiving in-service training in any of the following: ANC screening, counseling, nutritional assessment of pregnant women, management of pre-eclampsia/eclampsia, and antenatal corticosteroid for threatened preterm labor. The training must have involved structured sessions, and does not include individual instruction that a provider might have received during routine supervision

³ Functioning digital blood pressure apparatus or a functioning manual sphygmomanometer and a stethoscope

⁴ Average readiness score is the average number of 6 items (guidelines on ANC, staff trained for ANC any time, blood pressure apparatus, hemoglobin testing capacity, urine protein testing capacity, and iron or folic acid tablets) available for providing antenatal care services

Table 6.7: Supportive management for providers of antenatal care services

Among interviewed antenatal care (ANC) providers, the percentages who received training related to their work and personal supervision during the specified time periods, by background characteristics, Bangladesh HFS 2022

| | Percentage of i | interviewed provider | | | |
|----------------------------|---------------------|----------------------|--|-------------------|-------------------|
| | Training related to | Personal | Training related to ANC during the 24 months and | | |
| | ANC during the | supervision during | supervision during | Number of | Number of |
| | 24 months | the 6 months | the 6 months | interviewed ANC | interviewed ANC |
| Background | preceding the | preceding the | preceding the | service providers | service providers |
| characteristic | survey ¹ | survey ² | survey | (weighted) | (unweighted) |
| Facility type | | • | | , , , | (|
| District and upazila | | | | | |
| public facilities | 12.7 | 97.8 | 12.3 | 813 | 1,833 |
| DH | 15.5 | 97.4 | 14.3 | 206 | 334 |
| MCWC | 15.3 | 98.5 | 15.3 | 25 | 335 |
| UHC | 11.6 | 97.9 | 11.5 | 582 | 1,164 |
| Union-level public | | | | | |
| facilities | 9.4 | 86.4 | 8.5 | 342 | 638 |
| UHFWC | 9.2 | 88.8 | 7.9 | 247 | 429 |
| USC/RD | 10.0 | 80.3 | 9.9 | 95 | 209 |
| Community clinic (CC) | 22.3 | 92.2 | 20.9 | 1,276 | 681 |
| NGO clinic/hospital | 16.2 | 96.5 | 16.2 | 96 | 326 |
| Private hospital | 8.9 | 90.0 | 8.0 | 614 | 338 |
| Location | | | | | |
| Urban | 11.5 | 94.4 | 10.8 | 1,253 | 1,857 |
| Rural | 18.4 | 91.7 | 17.2 | 1,888 | 1,959 |
| Division | | | | | |
| Barishal | 13.0 | 91.1 | 12.5 | 233 | 404 |
| Chattogram | 23.2 | 93.3 | 22.3 | 530 | 623 |
| Dhaka | 11.9 | 94.4 | 11.3 | 785 | 666 |
| Khulna | 5.3 | 88.4 | 4.7 | 339 | 461 |
| Rajshahi | 10.9 | 94.3 | 9.8 | 454 | 513 |
| Rangpur | 18.7 | 92.6 | 18.0 | 393 | 511 |
| Sylhet | 22.8 | 94.4 | 20.6 | 162 | 304 |
| Mymensingh | 26.8 | 89.4 | 24.7 | 245 | 334 |
| Total | 15.6 | 92.7 | 14.7 | 3,141 | 3,816 |
| Total excluding CCs | 11.0 | 93.1 | 10.4 | 1,865 | 3,135 |
| Total public | 17.3 | 93.3 | 16.3 | 2,431 | 3,152 |
| Total public excluding CCs | 11.7 | 94.4 | 11.2 | 1,155 | 2,471 |

¹ Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision ² Personal supervision refers to any form of technical support or supervision from a facility-based supervisor or from a visiting supervisor. It may include, but is not limited to, review of records and observation of work, with or without any feedback to the health worker

Table 6.8: Training for antenatal care service providers

Among interviewed antenatal care (ANC) service providers, the percentages who reported receiving in-service training on topics related to ANC during the specified period before the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of interviewed providers of ANC who reported receiving in-service training on: | | | | | | | | | | | |
|--|---|-----------------------------|---------------------------------|-----------------------------|-------------------------------|-----------------------------|---------------------------------|-----------------------------|---|-----------------------------|--|--|
| | ANC counseling | | ANC screening | | Complications of pregnancy | | Family planning ¹ | | Sexually transmitted infections ² | | Number of | Number of |
| Background characteristic | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | ANC service providers (weighted) | ANC service providers (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public facilities | 8.3 | 26.9 | 6.8 | 22.9 | 6.6 | 24.4 | 10.4 | 27.4 | 1.0 | 8.8 | 813 | 1,833 |
| DH MCWC UHC | 9.7 9.8 7.7 | 34.3 47.3 23.4 | 6.7 6.1 6.9 | 28.4 41.0 20.2 | 6.6 8.0 6.5 | 28.4 45.6 22.0 | 13.6 24.4 8.7 | 31.8 72.9 23.9 | 0.7 2.0 1.0 | 8.2 18.7 8.6 | 206 25 582 | 334 335 1.164 |
| Union-level public | | | | | | | • | | | | | ., |
| facilities UHFWC USC/RD | 5.8 5.5 6.6 | 41.2 45.6 29.6 | 3.8 2.8 6.5 | 33.0 36.3 24.5 | 5.6 5.1 7.2 | 37.8 41.6 28.0 | 16.2 19.5 7.9 | 53.5 63.9 26.7 | 1.8 2.0 1.3 | 14.3 15.6 10.9 | 342 247 95 | 638 429 209 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 13.3 12.9 5.2 | 57.3 30.2 8.6 | 9.0 8.9 6.1 | 40.9 24.0 9.2 | 11.7 12.6 4.9 | 54.2 28.0 10.7 | 14.7 16.5 6.0 | 41.2 40.9 10.3 | 1.0 4.6 1.7 | 7.5 18.6 3.7 | 1,276 96 614 | 681 326 338 |
| Location | | | | | | | | | | | | |
| Urban Rural | 7.1 11.3 | 18.5 49.8 | 6.4 7.9 | 16.3 36.9 | 5.8 10.2 | 17.8 47.1 | 8.6 14.4 | 20.5 41.1 | 1.6 1.1 | 7.2 8.8 | 1,253 1,888 | 1,857 1,959 |
| Division | | | | | | | | | | | | |
| Barishal Chattogram | 10.1 16.1 | 41.4 46.4 | 9.1 10.9 | 35.3 35.6 | 9.7 11.7 | 39.4 41.2 | 8.4 20.4 | 28.5 44.0 | 2.2 1.8 | 6.6 9.8 | 233 530 | 404 623 |
| Dnaka Khulna Raishahi | 7.1 3.7 4 7 | 25.6 38.0 35.1 | 5.8 2.3 3.0 | 21.7 29.6 25.1 | 6.4 2.1 4.3 | 28.0 33.1 31.9 | 13.9 6.7 5.8 | 27.9 32.2 28.6 | 0.9 0.5 0.9 | 5.2 11.4 6.3 | 785 339 454 | 461 513 |
| Rangpur Sylhet | 10.9 18.2 | 34.7 44.6 | 9.9 13.9 | 26.6 37.7 | 11.3 18.3 | 33.6 41.2 | 7.4 18.0 | 28.7 42.2 | 1.9 3.2 | 12.4 6.4 | 393 162 | 511 304 |
| Mymensingh | 12.4 | 54.1 | 8.8 | 32.2 | 11.5 | 51.6 | 14.7 | 38.8 | 0.2 | 9.1 | 245 | 334 |
| I OTAI | 9.6 | 37.3 | 7.3 | 28.7 | 8.4 | 35.4 | 12.1 | 32.9 | 1.3 | 8.2 | 3,141 | 3,816 |
| Total excluding CCs | 7.1 | 23.6 44.0 | 6.1 7.5 | 20.3 | 6.1 0.1 | 22.5 41.0 | 10.3 12.5 | 27.2 | 1.5 | 8.6 8.0 | 1,865 2,116 | 3,135 |
| Total public excluding CCs | 7.5 | 44.9 31.1 | 7.5 5.9 | 33.0 25.9 | 9.1 6.3 | 41.9 28.4 | 12.1 | 35.1 | 1.1 | 0.9 10.4 | 719 | 2,471 |

Note: Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

¹ Includes training in any of the following: general counseling for family planning, insertion and/or removal of intrauterine contraceptive device (IUCD), insertion and/or removal of implants, performing vasectomy, performing tubal ligation, clinical management of family planning methods including managing side effects, family planning for HIV-positive women, postpartum family planning, injectable contraceptives, and emergency contraceptive pills

women, postpartum family planning, injectable contraceptives, and emergency contraceptive pills ² Includes training in any of the following: diagnosing and treating sexually transmitted infections (STIs), the syndromic approach to diagnosing and managing STIs, and treatment of drug-resistant STIs

Table 6.9: Key indicators for antenatal care services in MCWCs

Among facilities offering antenatal care (ANC) services, the percentages with key indicators, Bangladesh $\mbox{HFS}\ 2022$

| | Facility type | | | | |
|---|--|---|--|--|--|
| Indicators | MCWC (district-level) | MCWC (upazila and union-level) | | | |
| Availability | | | | | |
| Antenatal care (ANC) | 100.0 | 97.2 | | | |
| Number of facilities (weighted) | 5 | 3 | | | |
| Number of facilities (unweighted) | 62 | 38 | | | |
| Readiness | | | | | |
| Guideline on ANC ¹ Staff trained for ANC at any time ² Blood pressure apparatus ³ Hemoglobin testing capacity Urine protein testing capacity Iron or folic acid tablets Average readiness score⁴ | 48.2 93.4 98.3 67.6 66.4 100.0 4.7 | 35.4 78.1 92.0 45.7 46.5 97.2 3.9 | | | |
| Number of facilities offering ANC services (weighted) | 5 | 3 | | | |
| Number of facilities offering ANC services (unweighted) | 62 | 37 | | | |

¹ National ANC guidelines or other guidelines relevant to ANC

² Facility has at least one interviewed staff member providing ANC services who reports receiving in-service training in any of the following: ANC screening, counseling, nutritional assessment of pregnant women, management of pre-eclampsia/eclampsia, and antenatal corticosteroid for threatened preterm labor. The training must have involved structured sessions, and does not include individual instruction that a provider might have received during routine supervision

³ Functioning digital blood pressure apparatus or a functioning manual

sphygmomanometer and a stethoscope ⁴ Average readiness score is the average number of items (out of 6 items) available for providing antenatal care services

Key Findings

Delivery Care Services

- More than half (55%) of the public facilities (excluding CCs) offer normal delivery services (Table 7.1 and Figure 7.1.1)
- Normal delivery service is universally available in DHs (100%) and UHCs (100%). Moreover, 89% of the private facilities, 87% of the MCWCs, and 5% of the CCs offer normal delivery services. Additionally, 34% of the NGO facilities provide normal delivery services (Table 7.1 and Figure 7.1.1)
- Almost all of the DHs (100%) and private facilities (93%), and about half of the UHCs (53%), perform cesarean section, while only one-fifth of the NGOs (21%) offer this service (Table 7.1 and Figure 7.1.2).
- About half of public health facilities have at least one staff member trained in delivery care (46%) and one in ten have guidelines related to BEmOC and CEmOC (10%). Most of these facilities have a delivery pack on site (80%) (Table 7.2 and Figures 7.2.1, 7.2.2, and 7.2.6).
- About one-fourth of public facilities have at least one staff member trained in integrated management of pregnancy and childbirth (IMPAC) and post-abortion care (PAC). Less than half of public facilities have a staff member trained in routine labor and delivery care or active management of the third stage of labor (AMTSL) (Table 7.3).
- Since 2017, the availability of essential medicines among public facilities offering normal delivery services has increased— including oxytocin (23% to 40%), injectable antibiotics (9% to 15%), and magnesium sulphate (8% to 20%) (Table 7.4 and Figures 7.3.1–7.3.3).
- Most public facilities (excluding CCs) offering normal delivery services have delivery packs (91%), gloves (88%), and delivery beds (81%) available (Table 7.2). Partograph and examination light availability increased markedly between 2017 and 2022 (Figure 7.2.5).
- Overall, the average readiness score of public facilities (excluding CCs) for providing normal delivery services is 6.6 (among the WHO-specified 13 items). The average readiness score increased in almost all types of facilities from 2017 to 2022; nevertheless, the CCs experienced a decrease (Table 7.7 and Figure 7.6).
- Nearly half of the DHs performed all seven signal functions (BEmOC) in the last three months, whereas about one-fourth of the UHCs (26%) and one in seven MCWCs (15%) performed all seven signal functions (BEmOC) during that period (Table 7.6 and Figure 7.4).
- CEmOC signal functions (all nine) are performed in less than half of the DHs (44%); and less than one-third of NGOs (28%) and private hospitals (27%) (Table 7.6 and Figure 7.5).

Newborn Care Services

- Nearly one-third of the providers in public facilities (36%) have received in-service training on newborn resuscitation using a neonatal bag and mask at any time. Around one in three providers in public facilities (34%) reported that they received training on umbilical cord care (use of 7.1% chlorhexidine) (Table 7.13).
- The availability of equipment in public health facilities has increased markedly from 2017 to 2022: infant weighing scale (37% to 76%), functioning fetal stethoscope (6% to 28%), neonatal bag and mask (49% to 72%), and suction apparatus (20% to 32%) (Table 7.10 and Figure 7.8.1–7.8.4).

- The primary nine newborn signal functions are performed in half of the public facilities (excluding CCs) offering normal delivery services (**Table 7.10.1**).
- The majority of DHs (82%), UHCs (79%), NGOs (76%), and private facilities (51%) performed all nine primary newborn signal functions. However, less than one-fifth of CCs (15%) provided all nine primary newborn signal functions in the last three months (Table 7.10.1 and Figure 7.9.1).

7.1 BACKGROUND

B angladesh showed remarkable progress in reducing the maternal mortality rate (MMR) and the neonatal mortality rate (NMR) in the era of the Millennium Development Goals (MDGs). In accordance with the Sustainable Development Goals (SDGs), the Government of Bangladesh had set the ambitious target of reaching an MMR of 121 per 100,000 live births and an NMR of 18 per 100,000 live births by 2022 in the 4th Health, Population and Nutrition Sector Program (HPNSP), with an emphasis on maternal and newborn care. The MMR decreased from 322 per 100,000 live births in 2001 to 156 per 100,000 live births in 2022 (SVRS 2022), and the NMR fell from 87 per 1,000 live births in 1990 to 20 per 1,000 live births in 2022 (NIPORT 2022).

Continuous care during pregnancy and childbirth is important in reducing both maternal and newborn deaths. There is definitive global evidence that the availability of emergency obstetric and neonatal care (EmONC) and skilled attendance at childbirth are crucial to saving the lives of mothers and newborns. Institutional delivery care saves lives because a skilled birth attendant is immediately available to manage complications during labor and delivery or to refer the mother to the next level of care.

Hemorrhage and eclampsia alone are responsible for 55% of maternal deaths in Bangladesh. A significant number of women die during the pregnancy and delivery period; the pregnancy-related mortality rate is 206 per 100,000 live births. The Bangladesh Maternal Mortality and Health Care Survey (BMMS) 2016 has emphasized that after a significant reduction in the MMR from 2001 to 2010, this decline has begun to stall. No important differences in the MMR were found between the 2010 BMMS (194 maternal deaths per 100,000 live births) and 2016 BMMS (196 maternal deaths per 100,000 live births). However, the MMR has declined notably to 156 maternal deaths per 100,000 live births according to SVRS 2022. Only 41% of pregnant women had four or more antenatal care (ANC) visits as recommended by WHO (NIPORT 2022). Nearly two-thirds of births occurred at health facilities and were attended by medically-trained providers. It is important to note that 45% of births were cesarean section (C-section) (NIPORT 2022). Overall, however, the percentage of women receiving the complete continuum of maternity care (ANC, delivery care, and postnatal care from medically-trained providers) increased significantly between 2010 and 2016, from 19% to 43% (NIPORT 2016).

Although there have been remarkable reductions in NMRs worldwide, there are existing challenges to achieving SDG and national targets. In the least developed countries, the frequency of postnatal care (PNC) within two days of birth is very low, inhibiting physical examinations of newborns for danger signs and symptoms. In Bangladesh, the major causes of neonatal deaths include prematurity, pneumonia/infections, and intrapartum complications such as birth asphyxia. These preventable deaths can be eliminated with increases in recommended ANC, births attended by medically-trained providers, and PNC visits. Bangladesh is committed to addressing preventable causes of child mortality and to reaching the SDG target of reducing the NMR to 12 per 1,000 live births by 2030. The development of the "Promise Renewed Declaration: Bangladesh Call to Action 2013" and the inclusion of the National Newborn Health Program (NNHP) in the 4th HPNSP are examples of the Bangladesh government's continuous efforts to reduce the country's NMR.

Bangladesh has also reviewed its Maternal, Neonatal, Child, and Adolescent Health (MNC&AH) operational plan under the 4th HPNSP. The HPNSP focuses on strengthening safe deliveries at home, providing around-theclock EmONC services at the upazila level, and developing a functional referral system from community facilities to upazila-level hospitals. To deliver high-quality MNC&AH services and to achieve national outcomes, the Government of Bangladesh articulated the HNPSP and the Bangladesh Every Newborn Action Plan (BENAP) within the Health System Strengthening (HSS) framework.

This chapter explores the following key issues related to provision of quality delivery and newborn care services at health facilities in Bangladesh:

- Availability of maternal health services: Section 7.2, including Tables 7.1–7.5 and Figures 7.1.1–7.3.6, examines the availability of maternal health services, including the availability of service guidelines, staff with up-to-date training, basic items that support quality provision of delivery services, and items for infection control.
- Signal functions for emergency obstetric and newborn care: Section 7.3, including Table 7.6 and Figures 7.4–7.5, explores the extent to which facilities that provide normal delivery care performed nine signal functions at least once during the three months before the survey.
- Readiness of health facilities to provide normal delivery services: Section 7.4, including Table 7.7 and Figure 7.6, provides information on the readiness of health facilities to provide normal delivery services according to the WHO criteria, and average readiness score to provide normal delivery services.
- **Newborn care practices:** Section 7.5, including **Tables 7.8–7.10** and **Figures 7.7.1–7.8.5**, provides information on routine newborn care practices in health facilities and the availability of essential medicines for newborns.
- Newborn signal functions: Section 7.6, including Table 7.10.1 and Figures 7.9.1–7.9.4, explores the extent to which facilities that provide delivery care performed 20 newborn signal functions at least once during the three months before the survey.
- Basic management and administrative systems: Section 7.7, including Tables 7.11–7.13 and Figures 7.10.1–7.10.2, considers the extent to which essential management and administrative systems are in place to support quality services, including personal supervision and in-service training for providers of delivery and newborn care.

7.2 AVAILABILITY OF MATERNAL HEALTH SERVICES

7.2.1 Service Provision

Table 7.1 and **Figures 7.1.1–7.1.2** provide information on the availability of various maternal health services. Although ANC and PNC are offered universally across all types of public and NGO health facilities, they are comparatively lower in the private health facilities (**Table 7.1**).

- More than half (55%) of the public facilities (excluding CCs) offer normal delivery services.
- Normal delivery services are universally available in DHs (100%) and UHCs (100%). Moreover, 89% of the private facilities, 87% of the MCWCs, and 34% of the NGO facilities offer normal delivery services. Additionally, around 5% of the CCs provide normal delivery services (Table 7.1 and Figure 7.1.1).

- The availability of normal delivery services in UHCs reached 100% in 2022, an increase from 95% in 2017 (Figure 7.1.1).
- Normal delivery service availability is five times higher in urban areas (82%), compared to rural areas (17%) (Table 7.1).



Figure 7.1.1 Availability of normal delivery services in health facilities, by facility type

- Availability of cesarean delivery services is universal in DHs (100%), and over 9 in 10 private facilities (93%) are offering the service in 2022 (**Table 7.1** and **Figure 7.1.2**).
- Cesarean deliveries are available in more than half of the UHCs (53%); and one in five NGOs (21%), which has nearly doubled since 2017 (**Figure 7.1.2**).
- Eighty percent of urban facilities perform cesarean deliveries, whereas the service is available in 1% of rural facilities (**Table 7.1**).





■ 2014 BHFS ■ 2017 BHFS ■ 2022 BHFS

7.2.2 Trained Staff, Guidelines, and Equipment for Delivery Services

The availability of service trained staff, guidelines, and certain basic equipment are the key elements in the provision of quality delivery services. **Table 7.2** reports the extent to which these items were available on the day of the survey in facilities that offer normal delivery services.

- The availability of trained staff at any time in delivery care is nearly half (46%) in public health facilities offering normal delivery services (**Table 7.2**).
- The availability of trained staff at any time has increased markedly from 2017 to 2022 in MCWCs (66% to 82%), USC/RDs (27% to 42%), CCs (36% to 57%), and private hospitals (16% to 23%) (Figure 7.2.1).





- Overall, 13% of public health facilities (excluding CCs) offering normal delivery services have any guidelines related to BEmOC and CemOC (Table 7.2 and Figure 7.2.2).
- The availability of guidelines on BemOC and CemOC on site is highest in NGOs (29%), in comparison . with other types of facilities (Figure 7.2.2).
- Rural facilities (9%) are more likely than urban facilities (7%) to have any of the guidelines for BemOC or CemOC (Table 7.2).



Figure 7.2.2 Availability of guidelines on maternal health in health facilities, by facility type

- Emergency transport may be critical if a woman needs to be referred to another facility for more specialized care. Among facilities offering normal delivery services, district and upazila level public facilities (86%) are more likely to have emergency transport compared to NGOs (42%) and private (32%) health facilities (Table 7.2).
- Almost all of the DHs (97%) and UHCs (89%) that offer normal delivery services universally have emergency transportation (**Table 7.2 and Figure 7.2.3**).

Figure 7.2.3 Availability of emergency transport in health facilities that offer normal delivery services, by facility type



■ 2014 BHFS □ 2017 BHFS ■ 2022 BHFS

- The availability of delivery beds is more than 80% in public health facilities (excluding CCs) in 2022, which has increased gradually from 74% in the 2014 BHFS (**Table 7.2** and **Figure 7.2.4**).
- NGO clinics/hospitals have shown a rise in the availability of delivery beds, from 84% in the 2014 BHFS to 95% in the 2022 BHFS (Figure 7.2.4).



Figure 7.2.4 Availability of delivery beds in health facilities, by facility type

- Overall, the availability of functioning examination lights in public health facilities offering normal delivery services increased from 57% in 2017 to 71% in 2022 (Figure 7.2.5).
- More than 90% of DHs (94%), MCWCs (91%), NGO clinics (93%), and private facilities (91%) offering normal delivery services have an examination light (Table 7.2 and Figure 7.2.5).



Figure 7.2.5 Availability of functioning examination light in health facilities, by facility type

- Since 2014, there has been a 32-percentage point increase in the availability of delivery packs (from 59% to 91%) in public health facilities (excluding CCs) offering normal delivery services (Figure 7.2.6). Availability of delivery packs means either the facility has a sterile delivery pack available at the delivery site or all the following individual equipment must be present: cord clamp, episiotomy scissors, scissors/blade to cut cord, suture material with needle, and needle holder.
- Most of the facilities offering normal delivery services have delivery packs; however, only one-third of the CCs that perform normal deliveries had delivery packs available on the day of survey (Table 7.2 and Figure 7.2.6).



Figure 7.2.6 Availability of delivery pack in health facilities, by facility type

- Half of public health facilities (excluding CCs) offering normal delivery services have partographs on site in 2022, which has increased notably from about one-fifth in 2017 (Table 7.2 and Figure 7.2.7).
- Among facilities offering normal delivery services, partographs are universally available in DHs (90%), UHCs (90%), and NGOs (92%); however, it is found in only about one in five private hospitals (Table 7.2 and Figure 7.2.7).

92 90 90 77 65 59 59 50 47 47 44 4342 41 39 40 31 26 23 21 2221 1716 1615 15 6 5 MCWC UHFWC USC/RD Total Total DH UHC CCs NGO Private public public excluding CCs ■ 2014 BHFS ■ 2017 BHFS ■ 2022 BHFS (N=280) (N=358) (N=412)

Figure 7.2.7 Availability of partograph in health facilities, by facility type

- Among facilities offering normal delivery services, most public facilities (85%) have gloves available in labor rooms. Gloves are available at 100% of the NGOs and 93% of the private hospitals. CCs are least likely (71%) to have gloves (**Table 7.2** and **Figure 7.2.8**).
- Availability of gloves in facilities offering normal delivery services increased in DHs (82% to 98%), MCWCs (83% to 95%), UHCs (72% to 95%), and private facilities (69% to 93%) from 2017 to 2022 (Figure 7.2.8).



Figure 7.2.8 Availability of gloves in health facilities, by facility type

Table 7.3 presents information on the availability of staff with in-service training in areas related to delivery or newborn care. The table considers staff who ever received in-service training and staff who have received training within the 24 months before the survey, among facilities offering normal delivery services.

- Only 5% of the public facilities (excluding CCs) that offer normal delivery services have providers who received in-service training in integrated management of pregnancy and childbirth (IMPAC) during the 24 months preceding the survey; however, 24% of all public facilities have at least one provider who has ever received in-service training in IMPAC (Table 7.3).
- Only 8% of the total public and private facilities have providers who had received in-service training on routine care for labor and delivery during the past 24 months; however, 43% of the public facilities and 26% of private facilities have at least one provider who has ever received in-service training (Table 7.3).
- Fifty-nine percent of public facilities reported having at least one staff member who had ever received in-service training in neonatal resuscitation, but only 18% of public facilities have at least one staff member with recent (past 24 months) in-service training on this topic (**Table 7.3**).
- Most of the NGO facilities (80%) and less than three in every ten private hospitals (28%) have at least one provider ever trained in neonatal resuscitation (Table 7.3).

- Thirty-eight percent of the public facilities have a staff member who has ever received in-service training in active management of the third stage of labor (AMTSL) (**Table 7.3**).
- Even fewer facilities have staff with in-service training in post-abortion care or emergency obstetric care at any time. Only one-fourth of the public facilities have at least one staff member who has ever received training in post-abortion care, while 22% of the public facilities (excluding CCs) have at least one staff member who has received training in emergency obstetric care (**Table 7.3**).

7.2.3 Medicines and Commodities for Delivery Care

Table 7.4 describes the availability of essential medicines and commodities for delivery. The table also provides information on the availability of priority medicines for mothers, as defined by the WHO.

Essential Medicines for Delivery

- Nearly half (49%) of the public facilities (excluding CCs) that offer normal delivery services have injectable uterotonics (oxytocin) which is required for active management of the third stage of labor and management of postpartum hemorrhage (**Table 7.4** and **Figure 7.3.1**).
- The availability of oxytocin is relatively higher among NGOs (96%), followed by UHCs (91%), DHs (84%), and private facilities (82%). The availability of injectable uterotonics is lowest among CCs (<1%), where normal delivery service is available (**Table 7.4** and **Figure 7.3.1**).



Figure 7.3.1 Availability of oxytocin in health facilities, by facility type

Injectable antibiotics are required for the management of puerperal sepsis. The availability of injectable antibiotics (injectable penicillin, injectable gentamicin, injectable ampicillin, or injectable ceftriaxone) is highest in DHs (98%). More than two-thirds of the NGO clinics (70%) and private facilities (72%) have injectable antibiotics (Table 7.4 and Figure 7.3.2).





- One in four public facilities (excluding CCs) have injectable magnesium sulphate, which is essential for the management of eclampsia. Sixty-five percent of DHs, 62% of private facilities, 34% of MCWCs, and 56% of UHCs have injectable magnesium sulphate available on site (Table 7.4 and Figure 7.3.3).
- The availability of magnesium sulphate is the highest in NGO hospitals (68%). However, less than 1% of CCs have injectable magnesium sulphate for the management or pre-referral management of eclampsia (**Table 7.4** and **Figure 7.3.3**).

Figure 7.3.3 Availability of magnesium sulphate in health facilities, by facility type



- Skin disinfectant is available in 56% of the public health facilities (excluding CCs) in 2022, which increased from 49% in 2017 (**Table 7.4** and **Figure 7.3.4**).
- The availability of skin disinfectant is relatively higher in NGO facilities (89%), followed by private facilities (83%) and UHCs (80%), in comparison with other types of facilities. The availability of skin disinfectant is the least common among CCs (20%) that offer normal delivery services (**Table 7.4** and **Figure 7.3.4**).



Figure 7.3.4 Availability of skin disinfectant in health facilities, by facility type

- More than one-third of the public health facilities (excluding CCs) offering normal delivery services have intravenous fluids with infusion set (**Table 7.4** and **Figure 7.3.5**).
- Among facilities offering normal delivery services, the availability of IV fluids with infusion set is higher among the DHs (66%), UHCs (64%), NGOs (64%), and private hospitals (52%) (Table 7.4 and Figure 7.3.5).



Figure 7.3.5 Availability of intravenous fluids with infusion set in health facilities, by facility type

Priority Medicines for Mothers

- Table 7.4 also provides information on availability of the WHO-defined priority medicines for mothers on the day of the survey. In general, these priority medicines are not widely available, and they are much more likely to be available in higher-level than in lower-level facilities. Moreover, even among higher-level facilities, there was considerable variation in priority medicine availability. Sodium chloride injectable solution is one of the widely available (51%) priority medicines in public facilities (excluding CCs) offering normal delivery services; 73% of NGOs and 60% of private hospitals, reported having this medication (Table 7.4).
- Sixty-seven percent of public facilities offering normal delivery services have misoprostol capsules or tablets, which are important for the management of postpartum hemorrhage. About three-fourths (74%) of union-level facilities have misoprostol capsules or tablets. They are available at 48% of DHs and 62% of UHCs, and also in around three-fourths of NGO clinics (77%) and half of the private hospitals (51%) (Table 7.4).

7.2.4 Items for Infection Control during Provision of Delivery Care

Infection control is vital during delivery care. The basic items for infection control in the delivery service area are soap, running water, alcohol-based hand disinfectant, latex gloves, sharps container, and waste receptacle. **Table 7.5** shows the proportion of facilities that had the indicated items and the average number of items considered important for infection control available at the service site on the day of the survey visit.

• Overall, almost all public facilities (87%) that offer normal delivery services have either soap and running water or alcohol-based hand disinfectant. The availability of these items is almost universal in DHs, MCWCs, UHCs, NGO clinics, and private hospitals (**Table 7.5**).

- Latex gloves (85%), sharps containers (61%), and waste receptacles (70%) are observed in the majority of public facilities (Table 7.5).
- Between 2017 and 2022, the average number of items available in public facilities (excluding CCs) offering normal delivery services for infection control has improved from 3.9 to 4.5. All types of facilities have seen an increase in the overall score for the average items related to infection control, except private facilities (**Figure 7.3.6**).



Figure 7.3.6 Average Items available (out of six) for infection control in delivery service area

7.3 SIGNAL FUNCTIONS FOR EMERGENCY OBSTETRIC CARE

Complications of labor and delivery can be expected to occur in a certain percentage of deliveries. It is estimated that approximately 15% of mothers develop life-threatening complications at the time of delivery, even if the mothers are otherwise healthy during the antenatal period. In such situations, facilities must be equipped to provide Emergency Obstetric and Neonatal Care (EmOC). Within EmOC, there are nine signal functions, layered in three levels: Obstetric first aid, Basic Emergency Obstetric and Neonatal Care (EEmOC). Facilities are considered as BEmOC facilities if they perform the first seven signal functions over the designated three-month period. The facilities are considered as CEmOC facilities if they perform all nine signal functions over a designated three-month period. Each of the facilities that provide normal delivery care were asked whether they had performed any number of the nine signal functions at least once during the three months preceding the survey.

- Among public facilities that offer normal delivery services, only 5% performed seven signal functions (BEmOC) in the last three months. Nearly half of the DHs (47%) performed the basic seven signal functions. The estimates are much lower in MCWCs (15%) and UHCs (26%) (Table 7.6).
- Only 1% of union level facilities and less than 1% of CCs performed the seven basic signal functions in the last three months. Basic signal function availability is 38% in private facilities and 32% in NGO

clinics that offer normal delivery services (**Table 7.6**). A notable increase is observed in NGO clinics, from 9% in 2017 to 32% in 2022(**Figure 7.4**).

• Urban facilities (37%) are more likely than rural facilities (2%) to perform the basic seven signal functions for emergency obstetric care (**Table 7.6**).



Figure 7.4 Trends in BEmOC signal functions by facility type

- Less than half of DHs (44%) and about one-quarter of the private facilities (27%) have performed all nine signal functions (CEmOC) during the past three months. However, only 11% of UHCs and 4% of MCWCs performed all CEmOC signal functions (Table 7.6 and Figure 7.5) during that time.
- The proportion of facilities that performed all nine signal functions increased notably from 2014 to 2022 among NGO clinics (9% to 28%), UHCs (5% to 11%), and private facilities (16% to 27%), and slightly decreased in MCWCs (6% to 4%) (**Figure 7.5**).



Figure 7.5 Trends in CEmOC signal functions by facility type

7.4 READINESS OF HEALTH FACILITIES TO PROVIDE NORMAL DELIVERY SERVICES

The WHO assesses service readiness for normal delivery based on the availability of specific items/tracer indicators in health facilities. In this section of the report, data from the 2022 BFHS are used to construct a slightly less restrictive and Bangladesh-context-appropriate version of the WHO measures. The measure of readiness to provide normal delivery services includes the following 13 items/tracer indicators:

Trained staff:

Percent

- At least one provider trained in delivery care at any time
- Guidelines:
 - National or other BEmOC or CEmOC guidelines available at the facility
- Equipment:
 - Examination light
 - Delivery pack
 - Suction apparatus
 - Neonatal bag and mask
 - Partograph
 - Gloves

Medicines and commodities:

- Injectable oxytocin
- Injectable antibiotic

- Magnesium sulphate
- Skin disinfectant
- Intravenous solution with infusion set

Table 7.7 and **Figure 7.6** show the average readiness score of thirteen items/tracer indicators to assess a health facility's readiness for normal delivery services.

- Among all public health facilities, the average readiness score for normal delivery services increased from 4.7 in 2014 to 5.9 in 2022 (**Figure 7.6**).
- DHs and NGO facilities have an average of 10.3 items necessary for normal delivery, the highest of all facilities (**Table 7.7** and **Figure 7.6**).
- Except in CCs, the average readiness score improved in all facility types since 2014. The readiness score declined in CCs from 3.4 in 2014 to 2.8 in 2022 (Figure 7.6).

Figure 7.6 Average readiness score of health facilities (out of 13 items) to provide normal delivery services, by facility type

Average



7.5 NEWBORN CARE PRACTICES

7.5.1 Routine Newborn Care

Routine newborn care is crucial to ensure the survival of newborns. **Table 7.8** presents information regarding routine newborn care practices at facilities offering normal delivery services.

Almost all types of public facilities (93%) offering normal delivery services engage in routine newborn
practices such as routine complete (head-to-toe) examination of newborns before discharge (Table 7.8).

- Weighing the newborn immediately upon delivery is nearly universal at DHs, MCWCs, UHCs, NGOs, and private clinics/hospitals offering normal delivery services (**Table 7.8**).
- There are strong differences in the administration of vitamin K to newborns among facilities offering normal delivery services. Half of the union-level public facilities (50%) administer vitamin K, whereas this service is universal in DHs (97%), UHCs (97%), and private facilities (99%), and widely available in MCWCs (80%) and NGO clinics (88%) (**Table 7.8**).

7.5.2 Essential Medicines for Newborn Care

- Eighty-four percent of public health facilities that offer normal delivery services had amoxicillin syrup or suspension available on the day of the visit. The availability of amoxicillin is the highest among CCs (91%), UHFWCs (90%), and MCWCs (90%). Around one-third of the NGO facilities (32%) and one-fifth of the private facilities (19%) had amoxicillin available on the day of the visit (Table 7.9 and Figure 7.7.1).
- The availability of amoxicillin declined from 71% to 32% in NGO facilities and 55% to 19% in private facilities from 2014 to 2022. USC/RDs and DHs also noted sharp declines (**Figure 7.7.1**).



Figure 7.7.1 Availability of amoxicillin in health facilities, by facility type

- Injectable gentamicin is available in about one-third of public health facilities (31%), excluding CCs, in 2022—an increase from 9% in 2017 (Table 7.9 and Figure 7.7.2).
- The availability of gentamicin among facilities offering normal delivery services is highest in UHCs (78%), increasing nearly two-fold from 2017. Less than two-thirds of the DHs (58%), 45% of private facilities, and 39% of MCWCs had valid gentamicin available on the day of the visit. Gentamicin is available in only about one-fourth of the UHFWCs (23%) (**Table 7.9** and **Figure 7.7.2**).



Figure 7.7.2 Availability of gentamicin in health facilities, by facility type Percent

- Most public health facilities (excluding CCs) offering normal delivery services (83%) had 7.1% chlorhexidine available on the day of the visit. The availability is highest in UHCs (96%) and lowest in private facilities (38%) (Table 7.9 and Figure 7.7.3).
- The availability of 7.1% chlorhexidine has increased in both public health facilities (from 27% in 2014 to 77% in 2022) and NGO facilities (from 46% in 2014 to 89% in 2022) (**Figure 7.7.3**).

Figure 7.7.3 Availability of 7.1% chlorhexidine in health facilities, by facility type



7.5.3 Availability of Equipment for Newborn Care Services

Table 7.10 provides information on the availability of equipment for newborn care services among facilities that offer normal delivery services.

- Overall, the availability of various types of equipment for newborn care services among facilities that offer normal delivery services is higher in urban areas in comparison with rural areas (**Table 7.10**). This includes incubator, suction apparatus with catheter, suction bulb or penguin sucker, newborn bag and mask, infant scale, fetal stethoscope, thermometer, and thermometer for low body temperature.
- Eighty percent of public health facilities (excluding CCs) offering delivery services have functioning infant weighing scales in 2022, nearly a two-fold increase from 41% in 2017. The availability of an infant scale is almost universal among NGO clinics (98%), followed by UHCs (96%), MCWCs (94%), DHs (90%), and private facilities (89%). Nearly six in ten CCs (59%) have an infant weighing scale observed on site (Table 7.10 and Figure 7.8.1).





- More than one-quarter (28%) of public health facilities (28%) offering normal delivery services have a functioning fetal stethoscope in 2022, which has increased markedly from 6% in 2017 (Table 7.10 and Figure 7.8.2).
- The majority of NGOs (83%), UHCs (75%), and DHs (71%) offering normal delivery services had functional fetal stethoscopes on the day of visit (**Table 7.10** and **Figure 7.8.2**).



Figure 7.8.2 Availability of functioning fetal stethoscope for newborn care services, by facility type

- Most public health facilities (excluding CCs) that offer normal delivery services (83%) have a functioning newborn bag and mask available in 2022 (Table 7.10), compared to 56% in 2017 (Figure 7.8.3).
- Newborn bags and masks are almost universally available in DHs (98%), UHCs (97%), and NGO facilities (95%). The availability of newborn bags and masks for newborn care services is also reasonably high in private facilities (82%). Newborn bags and masks are available in around one-quarter of the CCs (26%) (Table 7.10 and Figure 7.8.3).



Figure 7.8.3 Availability of functioning newborn bag and mask for newborn care services, by facility type

• Four in ten (39%) public health facilities (excluding CCs) that offer normal delivery services reported having suction apparatus with catheter to conduct delivery on the day of the survey (**Table 7.10**). There is a marked increase in UHCs (from 57% in 2017 to 78% in 2022); however, the availability of functioning suction apparatus with catheter increased in all types of facilities except CCs and private facilities in the last five years (**Figure 7.8.4**).





- The availability of functioning suction bulb or penguin sucker for newborn care services in facilities that offer normal delivery services increased from 64% to 73% among public health facilities from 2017 to 2022 (**Figure 7.8.5**).
- A suction bulb or penguin sucker is universally available among UHCs (98%), NGO clinics (97%), and DHs (92%). More than four in five private facilities (85%) have functioning suction bulbs or penguin suckers (**Table 7.10** and **Figure 7.8.5**).

Figure 7.8.5 Availability of functioning suction bulb or penguin sucker for newborn care services, by facility type



7.6 NEWBORN SIGNAL FUNCTIONS

Reduction of the neonatal mortality rate stands as an important target of the sustainable development goals (SDGs). To achieve this, all health facilities should be adequately prepared for providing high-quality delivery care during childbirth and effectively limiting preventable neonatal deaths, thereby reducing the overall neonatal mortality rate. Service availability and readiness of health facilities for delivering newborn services can be measured through a set of newborn signal functions. The 2022 BHFS incorporated a comprehensive set of newborn signal functions for the assessment of health facilities in this regard. Each of facility providing normal delivery care was asked whether they had performed the newborn signal functions ever and, if so, whether they had carried those out at least once during the three months preceding the survey.

The newborn signal functions are categorized into primary, basic, comprehensive, and advanced newborn signal functions based on their necessity and importance in preventing neonatal deaths.

Primary newborn signal functions (NSF): Iron and folic acid supplementation in pregnant women; hand wash each time before touching the baby; drying of the baby immediately after birth; delayed umbilical cord clamping;

clean cord cutting sterile blade; single application of 7.1% chlorhexidine (CHX); neonatal resuscitation; perform skin-to-skin care immediately after birth; and early initiation of breast feeding.

Basic newborn signal functions: Primary newborn signal functions and immunize patients with tetanus toxoid, administer oral antibiotic, administer intramuscular antibiotic, and newborn airway suction.

Comprehensive newborn signal functions: Basic newborn signal functions and administer intravenous antibiotic; administer antibiotic for premature rupture of membrane; oxygen therapy for management of newborn infection and respiratory problems; KMC for preterm or LBW newborn; and phototherapy for term or preterm with hyperbilirubinemia.

Advanced newborn signal function: Comprehensive newborn signal function and incubator support for sick term baby, pre-term baby, or if the baby receiving KMC becomes sick; and corticosteroids for pre-term labor.

- The primary nine newborn signal functions are performed in half of the public facilities (excluding CCs) offering normal delivery services (**Table 7.10.1**).
- Most of the DHs (82%), UHCs (79%), and NGOs (76%) performed all nine primary newborn signal functions during the last three months preceding the survey. Primary newborn signal functions are performed in two-thirds of MCWCs (65%) and more than half of private hospitals (51%). However, only 15% of CCs offering normal delivery services provided all nine primary newborn signal functions in the last three months (Table 7.10.1 and Figure 7.9.1).



Figure 7.9.1 Primary newborn signal functions, by facility type, 2022 BHFS

- Nearly four in ten DHs, three in ten NGOs, and two in ten private hospitals performed all 13 basic newborn signal functions in the last three months (Table 7.10.1 and Figure 7.9.2).
- Thirteen percent of MCWCs perform the basic newborn signal functions (**Table 7.10.1** and **Figure 7.9.2**).



Figure 7.9.2 Basic newborn signal functions, by facility type, 2022 BHFS Percent

- Among all facilities offering delivery services, performance of the 18 comprehensive signal functions in the three months before the survey is very low. DHs have the highest (31%) performance of comprehensive newborn signal functions during the three months before the survey (Table 7.10.1 and Figure 7.9.3).
- About one in ten UHCs and NGOs perform the 18 comprehensive newborn signal functions (**Table 7.10.1** and **Figure 7.9.3**).

Figure 7.9.3 Comprehensive newborn signal functions, by facility type, 2022 BHFS



- Around one-fifth of the DHs (21%) reported performance of all 20 advanced newborn signal functions within the last three months before the survey (**Table 7.10.1** and **Figure 7.9.4**).
- The practice of performing all 20 advanced newborn signal functions is very low in UHCs (4%) and private (4%) facilities (**Table 7.10.1** and **Figure 7.9.4**).

Figure 7.9.4 Advanced newborn signal functions, by facility type, 2022 BHFS Percent



7.7 BASIC MANAGEMENT AND ADMINISTRATIVE SYSTEMS

7.7.1 Recent In-Service Training and Supervision

Supportive management practices such as training and supervision play a significant role in ensuring quality of care. In-service training in maternal and newborn health care not only improves the knowledge of delivery care providers but also enhances their skills. **Table 7.11** presents information on specific in-service training reported by providers of normal delivery or newborn care services within six months or 24 months of the survey.

- About one-fifth of providers in public facilities received training related to delivery care and/or newborn care during the 24 months preceding the 2022 BHFS, as compared with about one-fourth of providers at NGO facilities and one-tenth of providers at private hospitals (**Table 7.11**).
- Supportive supervision helps to sustain providers' knowledge and skills. Almost all providers in public and NGO facilities have universally reported that they received personal supervision during the six months preceding the survey, except those in private facilities (88%) (**Table 7.11**).
- More than 90% of providers at rural facilities (95%) and urban facilities (93%) report receiving personal supervision (Table 7.11).

7.7.2 In-Service Training in Newborn and Delivery Care by Topic

Table 7.12 shows the percentages of providers who reported receiving specific in-service training related to delivery and newborn care.

- Around 10% of the providers of delivery care or newborn services in public facilities received training on early and exclusive breastfeeding, thermal care, sterile cord cutting and care, and kangaroo mother care for low-birth-weight babies during the 24 months preceding the survey (Table 7.12).
- The proportion of providers who reported ever receiving training in these areas is more than three times higher than those who reported receiving training in the 24 months preceding the survey. Only 5% of providers reported receiving training on newborn infection management during the 24 months preceding the survey (**Table 7.12**).

In-service training in delivery and newborn care services not only improves the knowledge of providers but also their skills. **Table 7.13** presents information on specific in-service training reported by interviewed providers of normal delivery or newborn care services. This includes training that they reported receiving either at any prior point or within 24 months of the survey.

- More than one-third of the providers of normal delivery and newborn care services in public facilities had received training on newborn resuscitation at any time. However, only 10% reported receiving such training in the past 24 months. Thirteen percent of the providers at private hospitals and 39% at NGO clinics had ever received training on resuscitation (**Table 7.13**).
- Around one-third of providers in public facilities had ever received in-service training on application of 7.1% chlorhexidine (for umbilical cord care), but only 10% reported receiving such training in the past 24 months. One in ten providers at private hospitals and around four in ten of those in NGO facilities had training on chlorhexidine application (**Table 7.13**).
- Around 15% of providers from public facilities had received in service training on emergency triage assessment (ETAT) at any time. The percentage of providers with ETAT training increased in all types
of facilities, with the exception of CCs. Only 7% of providers in CCs and 4% of providers in private facilities have staff trained on ETAT (**Table 7.13** and **Figure 7.10.1**).

Figure 7.10.1 Training on emergency triage assessment for newborn care services, by facility type

Percent



- Only 27% of providers in public facilities had ever received training in essential newborn care, and only 19% have training in comprehensive newborn care (**Table 7.13**).
- Less than 20% of the providers had been trained in integrated management of childhood illness (IMCI) at any time, and only 5% had received such training in the past 24 months. One-fourth of the providers at NGO facilities and a small percentage of providers at private hospitals (4%) had ever been trained in IMCI (Table 7.13).
- Providers in rural areas reported receiving all trainings in immediate newborn care more than their counterparts in urban areas (**Table 7.13**).
- In-service training on different services is presented in Figure 7.10.2. These services include newborn resuscitation, essential newborn care, umbilical cord care, and comprehensive newborn care package. Thirteen percent of all providers in public health facilities have training on all the services. The percentage of providers having such training is highest in MCWCs (25%) and UHFWCs (25%); and lowest in private facilities (6%) (Table 7.13 and Figure 7.10.2).



Figure 7.10.2 In-service training on different services* for newborn care, by facility type

Percent

*Services: Newborn resuscitation, essential newborn care, umbilical cord care, comprehensive newborn care

DELIVERY AND NEWBORN CARE

LIST OF TABLES

- Table 7.1 Availability of maternal health services
- Table 7.2 Guidelines, trained staff, and equipment for delivery services
- Table 7.3 Availability of staff trained on normal delivery or newborn care
- Table 7.4 Medicines and commodities for delivery
- Table 7.5 Items for infection control during provision of delivery care
- Table 7.6 Signal functions for emergency obstetric care
- Table 7.7 Readiness of health facilities to provide normal delivery services
- Table 7.8 Newborn care practices
- Table 7.9
 Essential medicines for newborn care
- Table 7.10 Availability of equipment for newborn care services
- Table 7.10.1 Newborn signal functions
- Table 7.11 Supportive management for providers of delivery care
- Table 7.12
 Training for providers of normal delivery services: Immediate newborn care
- Table 7.13
 Training for providers of normal delivery services: Immediate newborn care
- Table 7.14
 Key indicators for delivery and newborn care services in MCWCs

Table 7.1: Availability of maternal health services

Among all facilities, the percentages offering specific maternity services and the full range of maternity services, and among facilities that offer normal delivery services, the percentages having a skilled provider available on-site or on-call 24 hours a day to conduct deliveries, with or without an observed duty schedule, by background characteristics, Bangladesh HFS 2022

| | | | | | | | | | | | Percer facilities | tage of offering | | |
|--|--|--|--|--|--|--|--|---|---|---|---|---|---|--|
| | | | Perc | centage of fa | acilities offe | ring: | | | | | normal services | delivery that have: | _ | |
| Background characteristic | Antenatal care (ANC) | Normal delivery service | Cesarean delivery | Postnatal care (PNC) | ANC and normal delivery service | ANC, normal delivery, and cesarean delivery | ANC, normal delivery, and PNC | ANC, normal delivery, PNC, and cesarean delivery | Number of facilities (weighted) | Number of facilities (un- weighted) | Provider of delivery care available on-site or on-call 24 hours/ day, with observed duty schedule | Provider of delivery care available on-site or on-call 24 hours/ day, with or without observed duty schedule | Number of facilities offering normal delivery services (weighted) | Number of facilities offering normal delivery services (un- weighted) |
| Facility type | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 99.8 100.0 99.0 100.0 | 97.4 100.0 87.1 99.5 | 55.6 100.0 40.8 52.7 | 99.6 100.0 97.9 100.0 | 97.4 100.0 87.1 99.5 | 55.6 100.0 40.8 52.7 | 97.4 100.0 87.1 99.5 | 55.6 100.0 40.8 52.7 | 41 4 7 30 | 319 62 100 157 | 86.5 82.3 80.3 88.5 | 98.8 98.4 95.5 99.5 | 40 4 6 30 | 305 62 87 156 |
| Union-level public facilities UHFWC USC/RD | 98.3 100.0 93.7 | 49.4 59.2 23.4 | - | 96.8 99.0 90.9 | 49.4 59.2 23.4 | - | 49.2 58.9 23.4 | - | 310 225 85 | 434 293 141 | 36.7 37.7 29.5 | 86.1 87.7 75.6 | 153 133 20 | 206 170 36 |
| Community clinic (CC) | 99.8 | 4.7 | - | 97.4 | 4.7 | - | 4.7 | - | 994 | 488 | 30.0 | 52.1 | 46 | 24 |
| NGO clinic/hospital | 100.0 | 33.5 | 20.6 | 98.1 | 33.5 | 20.6 | 33.5 | 20.6 | 29 | 127 | 87.6 | 100.0 | 10 | 39 |
| Private hospital | 72.4 | 89.0 | 92.6 | 75.9 | 68.9 | 67.3 | 65.9 | 64.2 | 183 | 189 | 55.2 | 96.2 | 163 | 166 |
| Location Urban Rural | 79.3 99.3 | 81.7 16.6 | 80.3 0.7 | 80.9 97.3 | 66.9 16.5 | 61.4 0.6 | 64.5 16.5 | 59.0 0.6 | 235 1,322 | 514 1,043 | 60.7 39.4 | 96.3 80.5 | 192 220 | 409 331 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 98.8 97.4 95.4 90.5 97.8 97.7 98.0 97.5 | 19.2 30.6 30.1 30.4 26.4 17.1 31.6 18.2 | 6.4 13.5 17.8 20.8 12.2 5.9 6.9 6.3 | 91.9 96.6 94.7 90.5 96.3 97.4 93.4 95.5 | 18.2 28.2 27.5 23.5 25.1 15.8 31.6 17.4 | 5.2 10.7 14.7 11.4 10.9 4.2 6.9 5.5 | 18.2 27.8 27.0 22.6 24.8 15.4 31.6 17.4 | 5.2 10.2 14.2 10.5 10.9 3.8 6.9 5.5 | 111 291 317 210 214 193 99 122 | 152 244 208 212 190 136 151 | 48.6 56.8 56.1 40.6 43.0 53.3 36.8 43.9 | 85.4 95.2 90.6 92.2 86.5 78.5 67.4 82.1 | 21 89 96 64 56 33 31 22 | 73 124 133 92 102 84 69 63 |
| Total | 96.3 | 26.5 | 12.7 | 94.8 | 24.1 | 9.8 | 23.7 | 9.4 | 1,557 | 1,557 | 49.3 | 87.8 | 412 | 740 |
| Total excluding CCs | 90.1 | 65.0 | 35.2 | 90.3 | 58.4 | 27.0 | 57.3 | 26.0 | 563 | 1,069 | 51.8 | 92.4 | 366 | 716 |
| Total public | 99.5 | 17.8 | 1.7 | 97.3 | 17.8 | 1.7 | 17.8 | 1.7 | 1,345 | 1,241 | 43.8 | 81.7 | 239 | 535 |
| Total public excluding CCs | 98.5 | 55.0 | 6.5 | 97.1 | 55.0 | 6.5 | 54.8 | 6.5 | 351 | 753 | 47.1 | 88.8 | 193 | 511 |
| Note: "-" Means cesare | an delivery | is not offer | ed at union-l | evel public f | acilities or a | at public con | nmunity clin | iics. | | | | | | |

Table 7.2: Guidelines, trained staff, and equipment for delivery services

Among facilities offering normal delivery services, the percentages having guidelines, at least one staff member recently trained in delivery care, and basic equipment for routine delivery available in the facility on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Perce offerin serv | entage of fa lg normal c vices that h | acilities Ielivery ave: | | | | | | Equipment | t | | | | | | |
|-------------------------------------|---|--|---|---|-------------------------------------|-------------------------------|---|-------------------------------|---|-----------------------------|------------------------------|---------------------|---------------------|---|---|--|
| Background characteristic | Guide- lines on BEmOC or CEmOC ¹ | Staff trained in delivery care during the past 24 months ² | Staff trained in delivery care at any time ² | Emer- gency trans- port ³ | Examina- tion light ⁴ | Delivery pack ⁵ | Suction appa- ratus (mucus extractor) | Manual vacuum extractor | Vacuum aspirator or D&C kit ⁶ | Neonatal bag and mask | Parto- graph ⁷ | Gloves ⁸ | Delivery bed | Sterili- zation equip- ment ⁹ | Number of facilities offering normal delivery services (weighted) | Number of facilities offering normal delivery services (un- weighted) |
| Facility type | | | | | | | | | | | | | | | | |
| District and upazila public | | | | | | | | | | | | | | | | |
| facilities | 17.4 24.2 | 20.7 12 9 | 59.3 54.8 | 86.0 96.8 | 89.8 93.5 | 94.4 96.8 | 77.1 83.9 | 27.3 38.7 | 40.2 43.5 | 95.7 98.4 | 87.8 90.3 | 95.0 98.4 | 89.9 91 9 | 86.6 98.4 | 40 4 | 305 62 |
| MCWC | 12.6 | 19.4 | 81.8 | 63.5 | 90.7 | 93.1 | 68.7 | 28.6 | 40.5 | 89.4 | 77.3 | 95.3 | 86.3 | 85.0 | 6 | 87 |
| UHC | 17.4 | 22.1 | 55.2 | 89.2 | 89.1 | 94.4 | 77.9 | 25.4 | 39.7 | 96.7 | 89.6 | 94.5 | 90.3 | 85.3 | 30 | 156 |
| Union-level public facilities | 11.5 | 6.6 | 38.8 | 1.7 | 67.7 | 90.6 | 29.2 | 19.7 | 16.6 | 79.1 | 40.0 | 86.5 | 78.8 | 48.4 | 153 | 206 |
| UHFWC | 10.7 | 5.9 | 38.3 | 1.5 | 68.2 | 91.8 | 29.0 | 19.8 | 16.4 | 79.8 | 39.4 | 86.9 | 79.4 | 49.3 | 133 | 170 |
| USC/RD | 16.8 | 11.1 | 42.1 | 3.6 | 64.9 | 82.8 | 30.6 | 19.1 | 17.8 | 74.3 | 44.1 | 83.7 | 75.0 | 42.4 | 20 | 36 |
| Community clinic (CC) | 0.0 | 9.6 | 56.5 | 3.3 | 63.6 | 31.8 | 0.0 | 4.7 | 0.0 | 26.3 | 5.1 | 70.5 | 54.5 | 0.0 | 46 | 24 |
| NGO clinic/ hospital | 29.0 | 21.9 | 60.8 | 42.4 | 92.9 | 94.6 | 83.4 | 45.4 | 56.3 | 94.6 | 91.5 | 100.0 | 94.5 | 85.7 | 10 | 39 |
| Private hospital | 3.5 | 7.7 | 22.9 | 32.4 | 91.2 | 86.3 | 88.0 | 32.5 | 53.8 | 81.8 | 20.7 | 92.5 | 80.6 | 89.5 | 163 | 166 |
| Location | | | | | | | | | | | | | | | | |
| Urban | 6.5 | 9.7 | 27.7 | 40.0 | 91.2 | 87.3 | 86.2 | 32.8 | 52.1 | 84.3 | 31.5 | 92.9 | 82.1 | 89.9 | 192 | 409 |
| Rural | 9.3 | 8.6 | 45.1 | 8.5 | 68.9 | 78.9 | 28.0 | 17.1 | 15.7 | 69.1 | 36.8 | 84.2 | 74.8 | 41.0 | 220 | 331 |
| Division Barishal | 7.4 | 10.4 | 44.0 | 17.5 | 79.7 | 80.4 | 45.4 | 17.0 | 27.9 | 79.6 | 36.6 | 95.0 | 62.7 | 54.5 | 21 | 73 |
| Dhaka | 16.2 | 15.2 | 45.9 33.6 | 22.5 31.3 | 74.3 80.4 | 91.8 | 62.2 61.2 | 29.3 29.2 | 30.4 47.5 | 77.9 82.4 | 33.5 45 7 | 90.5 87 1 | 89.8 71.1 | 72.0 | 89 96 | 124 |
| Khulna | 3.3 | 4.0 | 31.1 | 23.9 | 70.7 | 70.1 | 55.2 | 13.1 | 19.3 | 60.9 | 17.9 | 81.8 | 83.8 | 64.7 | 64 | 92 |
| Rajshahi | 2.1 | 8.8 | 25.3 | 17.8 | 87.8 | 73.2 | 49.7 | 25.7 | 38.0 | 78.5 | 20.1 | 89.7 | 74.5 | 58.0 | 56 | 102 |
| Rangpur | 5.1 | 5.6 | 44.5 | 18.1 | 81.6 | 96.1 | 46.9 | 31.9 | 23.9 | 86.6 | 47.9 | 88.9 | 78.0 | 51.5 | 33 | 84 |
| Mymensingh | 8.5 7.4 | 6.9 | 34.9 47.7 | 22.4 17.6 | 89.6 78.6 | 73.0 | 51.5 40.6 | 24.6 9.3 | 28.5 26.2 | 67.2 75.0 | 51.4 25.7 | 93.8 83.8 | 76.5 73.9 | 40.1 | 22 | 69 63 |
| Total | 8.0 | 0.0 | 37.0 | 23.2 | 70.3 | 82.8 | 55 1 | 24.4 | 32.7 | 76.2 | 34.3 | 88.2 | 78.2 | 63.8 | /12 | 740 |
| | 0.0 | 5.1 | 57.0 | 20.2 | 13.5 | 02.0 | 55.1 | 24.4 | 52.7 | 10.2 | 54.5 | 00.2 | 10.2 | 05.0 | 412 | 740 |
| excluding CCs | 9.0 | 9.0 | 34.6 | 25.8 | 81.3 | 89.2 | 62.1 | 26.9 | 36.8 | 82.5 | 38.0 | 90.5 | 81.2 | 71.9 | 366 | 716 |
| Total public | 10.2 | 9.5 | 45.6 | 16.2 | 70.6 | 79.9 | 31.6 | 18.1 | 17.3 | 71.7 | 41.3 | 84.8 | 76.0 | 45.5 | 239 | 535 |
| Total public excluding CCs | 12.7 | 9.5 | 43.1 | 19.3 | 72.3 | 91.4 | 39.2 | 21.3 | 21.5 | 82.5 | 49.9 | 88.3 | 81.1 | 56.4 | 193 | 511 |

Note: The indicators presented in this table comprise the staff training and equipment domains for assessing readiness to provide delivery care within the health facility assessment methodology proposed by the WHO and USAID (2012).

Basic emergency obstetric care (BEmOC) guidelines or comprehensive emergency obstetric care (CEmOC) guidelines.

² Facility has at least one interviewed staff member providing the service who reports receiving in-service training in delivery care. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision. ³ Facility has a functioning ambulance or other vehicle for emergency transport that is stationed at and operates from the facility.

⁴ A functioning flashlight is acceptable.

⁵ Either the facility has a sterile delivery pack available at the delivery site or all of the following individual equipment must be present: cord clamp/thread, episiotomy scissors, scissors (or blade) to cut cord, suture material with needle, and needle holder. ⁶ Facility has a functioning vacuum aspirator or else a dilation and curettage (D&C) kit available.

⁷ A blank partograph is available at the service site.

⁸ Disposable latex gloves or equivalent are available at the service site.

⁹ Facility reports that some instruments are processed in the facility and the facility has a functioning electric dry heat sterilizer, a functioning electric autoclave, or a non-electric autoclave with a functioning heat source available somewhere in the facility.

Table 7.3: Availability of staff trained on normal delivery or newborn care

Among facilities that offer normal delivery services, the percentages with staff trained on specific topics related to delivery and newborn care, by background characteristics, Bangladesh HFS 2022

| | Percenta | age of facil | lities offerir | ng normal c | lelivery or | newborn c in-servic | are service e training | es that repo | ort having a | at least one | e staff men | nber with | | |
|--|---|--|--|--|---|--|--|---|--|--|--|---|---|---|
| Background characteristic | Training on IMPAC during the past 24 months | Training on IMPAC at any time | Training on routine care for labor and delivery during the past 24 months | Training on routine care for labor and delivery at any time | Training on AMTSL during the past 24 months | Training on AMTSL at any time | Training on emer- gency obstetric care/life- saving skills during the past 24 months | Training on emer- gency obstetric care/life- saving skills at any time | Training on post- abortion care during the past 24 months | Training on post- abortion care at any time | Training on neonatal resusci- tation during the past 24 months | Training on neonatal resusci- tation at any time | Number of facilities offering normal delivery services (weighted) | Number of facilities offering normal delivery services (unweighted) |
| Facility type | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 11.0 8.1 9.0 11.9 | 43.3 43.5 54.0 40.9 | 17.4 19.4 13.7 18.0 | 61.2 61.3 76.1 58.0 | 17.1 14.5 13.7 18.2 | 64.1 66.1 72.5 62.0 | 12.3 12.9 11.2 12.4 | 47.2 58.1 61.9 42.5 | 13.4 16.1 13.6 12.9 | 51.9 62.9 65.6 47.3 | 51.1 46.8 34.5 55.3 | 84.7 83.9 84.1 84.9 | 40 4 6 30 | 305 62 87 156 |
| Union-level public facilities UHFWC USC/RD | 2.9 2.8 3.5 | 19.1 19.7 15.4 | 5.3 4.3 12.3 | 35.5 34.7 41.1 | 3.8 3.3 7.1 | 32.1 32.3 30.7 | 1.8 1.5 3.5 | 15.6 14.7 21.6 | 3.1 2.7 5.8 | 19.1 19.1 19.3 | 13.3 12.9 16.3 | 55.0 56.2 46.7 | 153 133 20 | 206 170 36 |
| Community clinic (CC) | 0.0 | 24.9 | 9.6 | 51.8 | 0.0 | 34.2 | 0.0 | 16.9 | 0.0 | 20.9 | 4.6 | 48.4 | 46 | 24 |
| NGO clinic/hospital | 16.4 | 40.2 | 18.6 | 57.6 | 18.6 | 55.0 | 18.6 | 33.2 | 13.1 | 29.6 | 49.9 | 79.7 | 10 | 39 |
| Private hospital | 5.7 | 16.4 | 7.6 | 26.1 | 5.9 | 22.0 | 3.4 | 17.9 | 5.5 | 19.9 | 10.3 | 27.8 | 163 | 166 |
| Location Urban Rural | 6.6 3.2 | 20.1 22.8 | 9.4 7.2 | 31.3 41.3 | 7.9 4.1 | 27.6 35.9 | 4.9 2.5 | 21.9 18.6 | 7.1 3.0 | 25.1 21.2 | 16.7 14.8 | 37.1 55.6 | 192 220 | 409 331 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 4.7 10.5 1.7 3.6 3.7 2.5 7.1 1.2 | 14.8 30.2 12.4 26.0 13.1 32.2 26.6 18.9 | 10.7 15.3 6.1 3.4 7.7 4.6 10.4 3.6 | 44.7 45.9 31.6 32.6 25.5 43.6 36.2 43.4 | 7.8 9.5 3.4 3.3 3.0 4.6 12.0 7.5 | 42.5 40.2 26.6 29.6 19.3 41.7 33.9 34.8 | 6.3 8.3 3.0 0.3 1.2 7.3 1.6 | 21.4 25.4 18.2 20.7 9.0 27.0 21.9 20.4 | 11.9 7.2 4.1 0.7 2.1 3.0 12.2 4.0 | 25.5 30.5 15.9 22.3 12.4 31.6 31.9 25.7 | 14.6 16.2 16.3 13.2 15.5 20.5 15.1 13.7 | 54.9 46.3 41.6 43.9 37.9 63.6 58.5 56.2 | 21 89 96 64 56 33 31 22 | 73 124 133 92 102 84 69 63 |
| Total | 4.8 | 21.6 | 8.2 | 36.6 | 5.8 | 32.0 | 3.6 | 20.1 | 4.9 | 23.0 | 15.7 | 47.0 | 412 | 740 |
| Total excluding CCs | 5.4 | 21.1 | 8.0 | 34.7 | 6.6 | 31.7 | 4.1 | 20.5 | 5.5 | 23.3 | 17.1 | 46.8 | 366 | 716 |
| Total public | 3.7 | 24.3 | 8.2 | 43.0 | 5.3 | 37.9 | 3.2 | 21.1 | 4.2 | 24.9 | 18.0 | 58.7 | 239 | 535 |
| Total public excluding CCs | 4.6 | 24.2 | 7.8 | 40.9 | 6.5 | 38.8 | 4.0 | 22.2 | 5.2 | 25.9 | 21.2 | 61.2 | 193 | 511 |

Note: Training refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision. IMPAC = Integrated management of pregnancy and childbirth. AMTSL = Active management of third stage of labor.

Table 7.4: Medicines and commodities for delivery

Among facilities offering normal delivery services, the percentages with essential medicines and commodities for delivery care and priority medicines for mothers observed to be available on the day of the survey, by facility type, Bangladesh HFS 2022

| | | | | | Facilit | y type | | | | | | | | |
|--|--|-------------|------|------|---|--------|--------|----------------------------------|----------------------------|---------------------|-------|---------------------------|-----------------|-------------------------------------|
| Medicines | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commu- nity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Essential medicines for delivery ¹ | | | | | | | | | | | | | | |
| Injectable uterotonic | | | | | | | | | | | | | | |
| (oxvtocin) ² | 87.9 | 83.9 | 77.1 | 90.8 | 39.2 | 38.7 | 42.4 | 0.0 | 96.1 | 82.0 | 57.8 | 65.1 | 39.8 | 49.3 |
| Injectable antibiotic ³ | 72.5 | 98.4 | 51.7 | 73.1 | 4.4 | 3.4 | 11.5 | 0.0 | 69.7 | 72.2 | 38.9 | 43.8 | 15.0 | 18.6 |
| Injectable magnesium | | | | | | | | | | | | | | |
| sulphate ² | 53 7 | 64 5 | 33.6 | 56.4 | 16.8 | 16.6 | 17.8 | 0.0 | 68.3 | 61 9 | 37 5 | 42.3 | 10.8 | 24 5 |
| Injectable diazenam | 65.1 | 72.6 | 11 3 | 60.0 | 8.6 | 87 | 8.2 | 0.0 | 56 1 | 77.0 | 11 3 | 16.5 | 16.0 | 20.4 |
| Skin disinfectant | 78.4 | 72.0 | 74.8 | 80.0 | 10.0 | 50.3 | 47.0 | 19.6 | 80.2 | 82.5 | 63.1 | 68.6 | 10.4 | 55.8 |
| Introvonous fluido with | 70.4 | 12.0 | 74.0 | 00.0 | 43.5 | 50.5 | 47.0 | 13.0 | 03.2 | 02.5 | 05.1 | 00.0 | 40.0 | 00.0 |
| intravenous nuius with | 62.0 | 66.4 | 50.2 | 62.0 | 20.4 | 07.4 | 40.6 | 47 | 62.0 | F1 0 | 20.4 | 42.0 | 20.0 | 26.0 |
| Infusion set | 62.0 | 66.1 | 50.3 | 63.9 | 29.1 | 27.4 | 40.6 | 4.7 | 63.8 | 51.9 | 39.4 | 43.8 | 29.9 | 30.0 |
| Priority medicines for mothers ⁵ | | | | | | | | | | | | | | |
| Sodium chloride | | | | | | | | | | | | | | |
| injectable solution | 88.8 | 85.5 | 59.8 | 95.5 | 40.8 | 40.8 | 40.9 | 0.0 | 73.3 | 60.4 | 49.5 | 55.7 | 41.0 | 50.8 |
| Injectable calcium | | | | | | | | | | | | | | |
| duconate | 3.4 | 97 | 44 | 23 | 5.4 | 51 | 71 | 0.0 | 29.7 | 31.2 | 15 4 | 17.3 | 40 | 50 |
| Ampicillin powder | 0.4 | 0.1 | | 2.0 | 0.4 | 0.1 | | 0.0 | 20.1 | 0112 | 10.1 | 11.0 | 1.0 | 0.0 |
| for injection | 33.0 | 11 3 | 12 | 43.1 | 0.8 | 04 | 35 | 0.0 | 34 | 93 | 73 | 82 | 61 | 75 |
| Injectable | 55.0 | 11.0 | 1.2 | 40.1 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 5.5 | 7.0 | 0.2 | 0.7 | 7.0 |
| motropidozolo | 62.4 | FG F | 20.7 | 71 / | 1 5 | 1 4 | 2.2 | 0.0 | E1 E | E6 7 | 20.4 | 24.2 | 116 | 111 |
| | 03.1 | 50.5 | 20.7 | /1.4 | 1.5 | 1.4 | 2.2 | 0.0 | 51.5 | 50.7 | 30.4 | 34.2 | 11.0 | 14.4 |
| Misoprostol capsules | | 40.4 | 70.0 | | | 70.0 | 50.0 | 40.0 | | | | <u> </u> | 07.4 | 74 5 |
| or tablets | 63.4 | 48.4 | 79.3 | 62.2 | 13.1 | 76.3 | 56.2 | 48.8 | 76.5 | 50.8 | 60.9 | 62.4 | 67.1 | 71.5 |
| Azithromycin capsules | | | | | | | | | | | | | | |
| or tablets or oral liquid | 82.5 | 95.2 | 9.3 | 96.3 | 6.0 | 1.2 | 38.1 | 3.9 | 74.9 | 49.7 | 32.1 | 35.7 | 18.4 | 21.9 |
| Cefixime capsules | | | | | | | | | | | | | | |
| or tablets | 69.3 | 87.1 | 16.5 | 78.0 | 2.0 | 0.0 | 15.2 | 0.0 | 68.8 | 55.3 | 31.0 | 34.9 | 12.9 | 16.0 |
| Benzathine benzyl | | | | | | | | | | | | | | |
| penicillin powder | | | | | | | | | | | | | | |
| for injection | 11.7 | 24.2 | 1.1 | 12.0 | 2.9 | 2.9 | 3.4 | 7.8 | 0.9 | 6.6 | 5.7 | 5.5 | 5.3 | 4.8 |
| Injectable | | | | | | | | | | | | | | |
| betamethasone/ | | | | | | | | | | | | | | |
| dexamethasone | 67.3 | 71.0 | 27.5 | 75.3 | 2.0 | 0.4 | 13.1 | 0.0 | 53.5 | 53.4 | 29.7 | 33.5 | 12.6 | 15.6 |
| Nifedipine capsules | | | | | | | | | | | | | | |
| or tablets | 4.4 | 6.5 | 3.5 | 4.4 | 0.1 | 0.0 | 1.1 | 0.0 | 11.5 | 21.1 | 9.1 | 10.3 | 0.8 | 1.0 |
| | | | | | | | | | | | | | | |
| Number of facilities offering normal delivery services | 40 | 4 | c | 20 | 450 | 100 | 20 | 46 | 40 | 462 | 44.0 | 266 | 220 | 102 |
| (weighted) | 40 | 4 | б | 30 | 153 | 133 | 20 | 40 | 10 | 163 | 412 | 300 | 239 | 193 |
| Number of facilities offering normal | | | | | | | | | | | | | | |
| (unweighted) | 305 | 62 | 87 | 156 | 206 | 170 | 36 | 24 | 39 | 166 | 740 | 716 | 535 | 511 |

Note: The essential medicines presented in this table comprise the medicines domain for assessing readiness to provide basic obstetric care within the health facility assessment methodology proposed by the WHO and USAID (2012). ¹ All essential medicines for delivery were assessed and must be available at the service delivery site. ² Injectable uterotonic (e.g., oxytocin) and injectable magnesium sulphate are also classified as essential medicines for mothers. ³ Injectable penicillin, injectable gentamicin, injectable ampicillin, or injectable ceftriaxone. ⁴ Normal saline solution, lactated Ringer's solution, or 5% dextrose solution.

⁵ The priority medicines for mothers are defined by the WHO; the list is published at https://www.who.int/publications/i/item/WHO_EMP_MAR_2011.1

Table 7.5: Items for infection control during provision of delivery care

Among facilities offering normal delivery services, the percentages with indicated items for infection control observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | | Percenta | ge of facilit | es offering r | normal delive | ry services | s that have | items for infe | ction control | | |
|--|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|--|-------------------------------------|-------------------------------------|-------------------------------------|---------------------------------|---|---|
| | | | Soap and | Alcohol- based | Soap and running water or else alcohol- based | | | | Average items | Number of facilities offering normal delivery | Number of facilities offering normal delivery |
| Background | | Running | running | hand | hand | Latex | Sharps | Waste | available | services | services |
| characteristic | Soap | water ¹ | water | disinfectant | disinfectant | gloves ² | container | receptacle3 | (out of 6) ⁴ | (weighted) | (unweighted) |
| Facility type | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 84.5 88.7 90.7 82.5 | 93.6 93.5 95.3 93.2 | 82.7 88.7 89.5 80.4 | 90.8 95.2 86.3 91.1 | 96.2 98.4 95.4 96.1 | 95.0 98.4 95.3 94.5 | 54.0 56.5 60.9 52.2 | 83.1 88.7 75.8 83.8 | 5.0 5.2 5.0 5.0 | 40 4 6 30 | 305 62 87 156 |
| Union-level public facilities UHFWC USC/RD | 73.8 74.7 67.9 | 77.6 78.2 73.4 | 70.9 71.3 67.9 | 71.3 71.3 71.8 | 86.3 87.7 77.0 | 86.5 86.9 83.7 | 60.1 62.0 47.4 | 68.2 69.5 59.5 | 4.4 4.4 4.0 | 153 133 20 | 206 170 36 |
| Community clinic (CC) | 65.9 | 69.9 | 65.9 | 74.9 | 79.6 | 70.5 | 71.6 | 62.2 | 4.1 | 46 | 24 |
| NGO clinic/hospital | 93.0 | 95.1 | 93.0 | 95.1 | 100.0 | 100.0 | 75.5 | 91.3 | 5.5 | 10 | 39 |
| Private hospital | 84.3 | 86.6 | 83.2 | 86.0 | 90.2 | 92.5 | 35.2 | 65.2 | 4.5 | 163 | 166 |
| Location | | | | | | | | | | | |
| Urban Rural | 84.5 73.3 | 88.0 77.2 | 83.6 71.0 | 87.4 73.5 | 92.0 85.2 | 92.9 84.2 | 40.3 61.0 | 68.7 68.0 | 4.6 4.4 | 192 220 | 409 331 |
| Division | 91 7 | 94.9 | 70.1 | 70.0 | 97 / | 05.0 | 42.7 | 79.0 | 4.6 | 21 | 72 |
| Chattogram Dhaka | 78.8 78.8 | 87.6 79.5 | 78.5 77.0 | 90.1 81.3 | 93.1 85.9 | 90.5 87.1 | 56.8 51.4 | 70.2 70.7 | 4.7 4.5 | 89 96 | 124 133 |
| Khulna Rajshahi Rangpur | 68.3 78.8 87.7 | 69.0 83.8 89.0 | 63.8 78.3 85.4 | 63.3 83.0 73.6 | 79.4 91.3 94.9 | 81.8 89.7 88.9 | 32.0 49.8 66.1 | 47.0 72.8 75.5 | 3.6 4.6 4.8 | 64 56 33 | 92 102 84 |
| Sylhet Mymensingh | 89.3 73.6 | 91.4 81.6 | 87.6 73.6 | 89.9 70.9 | 95.1 80.4 | 93.8 83.8 | 77.0 37.2 | 74.1 73.1 | 5.2 4.2 | 31 22 | 69 63 |
| Total | 78.6 | 82.2 | 76.9 | 80.0 | 88.4 | 88.2 | 51.3 | 68.4 | 4.5 | 412 | 740 |
| Total excluding CCs | 80.2 | 83.8 | 78.3 | 80.6 | 89.5 | 90.5 | 48.8 | 69.1 | 4.5 | 366 | 716 |
| Total public | 74.1 | 78.8 | 71.9 | 75.3 | 86.7 | 84.8 | 61.3 | 69.6 | 4.4 | 239 | 535 |
| Total public excluding CCs | 76.0 | 80.9 | 73.4 | 75.4 | 88.4 | 88.3 | 58.8 | 71.3 | 4.5 | 193 | 511 |

¹ Piped water, water in bucket with specially fitted tap, or water in pour pitcher.
 ² Non-latex equivalent gloves are acceptable.
 ³ Waste receptacle with plastic bin liner.
 ⁴ Average number of items (soap, running water, alcohol-based hand disinfectant, latex gloves, sharps container, and waste receptacle) available for infection control at the service site on the day of the survey.

Table 7.6: Signal functions for emergency obstetric care

Among facilities offering normal delivery services, the percentages reporting that they performed the signal functions for emergency obstetric care at least once during the three months before the survey, by background characteristics, Bangladesh HFS 2022

| | Perce that a | ntage of fa | cilities nteral: | | Percent | age of facilit | ies that carr | ied out: | | Perce th | entage of fa at carried o | cilities ut: | Number of facilities | Number of facilities |
|--|--|--|--|--|--|--|--|--|--|--|--|---|--|--|
| Background characteristic | Antibiotics | Oxytocin | Anticon- vulsants | Assisted vaginal delivery | Manual removal of placenta | Removal of retained products of concep- tion (MVA) | Neonatal resus- citation | Blood trans- fusion | Cesarean delivery | Three signal functions ¹ | Seven signal functions ² | All 9 signal functions ³ | offering normal delivery services (weighted) | offering normal delivery services (unweighted) |
| Facility type | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 90.6 98.4 65.4 94.9 | 96.8 98.4 89.9 98.0 | 58.8 95.2 31.2 59.3 | 53.6 56.5 57.5 52.3 | 94.5 95.2 84.9 96.4 | 82.0 98.4 72.3 81.7 | 90.4 93.5 74.2 93.4 | 60.0 93.5 22.7 63.0 | 57.0 98.4 46.8 53.0 | 55.9 93.5 27.7 56.2 | 26.7 46.8 14.8 26.3 | 13.3 43.5 4.4 10.7 | 40 4 6 30 | 305 62 87 156 |
| Union-level public facilities UHFWC USC/RD | 29.8 30.0 29.1 | 60.2 59.3 66.6 | 9.7 10.9 1.1 | 37.8 38.5 33.3 | 64.5 65.5 57.6 | 38.3 38.2 39.0 | 50.1 49.6 52.8 | - | - | 5.2 5.8 1.1 | 0.9 0.8 1.1 | - | 153 133 20 | 206 170 36 |
| Community clinic (CC) | 0.0 | 31.8 | 0.0 | 17.2 | 57.2 | 36.0 | 29.7 | - | - | 0.0 | 0.0 | - | 46 | 24 |
| NGO clinic/ hospital | 90.2 | 100.0 | 49.3 | 64.1 | 90.3 | 75.3 | 86.6 | 55.0 | 61.5 | 46.0 | 31.5 | 27.7 | 10 | 39 |
| Private hospital | 89.7 | 93.5 | 64.2 | 57.0 | 80.2 | 64.7 | 80.9 | 70.1 | 96.6 | 61.6 | 37.6 | 26.8 | 163 | 166 |
| Location Urban Rural | 90.8 28.4 | 94.0 57.9 | 63.8 11.4 | 58.2 33.9 | 82.4 65.6 | 69.0 40.2 | 82.1 49.6 | 68.5 5.5 | 92.0 4.3 | 61.1 8.1 | 37.2 2.2 | 26.3 0.6 | 192 220 | 409 331 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 69.9 57.3 64.2 61.0 50.7 52.5 48.7 44.6 | 76.5 82.6 75.0 68.4 69.4 79.9 67.7 74.1 | 41.9 38.2 50.5 20.2 32.6 27.9 36.8 21.3 | 50.8 58.7 54.2 26.5 42.4 34.1 43.5 27.3 | 93.7 78.1 70.4 56.1 75.4 82.1 81.6 69.0 | 48.1 64.9 60.6 26.8 49.0 63.7 64.0 43.1 | 78.2 67.1 67.2 54.1 62.3 50.7 73.7 77.4 | 32.6 36.0 49.7 35.3 27.6 23.2 18.6 25.8 | 32.0 44.2 55.5 57.2 46.2 32.1 21.8 33.2 | 38.6 35.2 46.6 20.0 28.1 27.9 31.3 17.5 | 21.1 26.5 7.9 20.0 12.5 20.8 4.5 | 9.8 14.2 21.9 7.7 7.8 8.3 10.9 3.2 | 21 89 96 64 56 33 31 22 | 73 124 133 92 102 84 69 63 |
| Total | 57.5 | 74.7 | 35.9 | 45.2 | 73.4 | 53.6 | 64.8 | 34.9 | 45.2 | 32.8 | 18.6 | 12.5 | 412 | 740 |
| Total excluding CCs | 64.8 | 80.1 | 40.4 | 48.8 | 75.5 | 55.9 | 69.2 | 39.3 | 50.9 | 37.0 | 20.9 | 14.1 | 366 | 716 |
| Total public | 34.3 | 60.9 | 16.1 | 36.5 | 68.1 | 45.2 | 52.9 | 10.1 | 9.6 | 12.7 | 5.0 | 2.2 | 239 | 535 |
| Total public excluding CCs | 42.5 | 67.9 | 19.9 | 41.1 | 70.8 | 47.4 | 58.5 | 12.5 | 11.9 | 15.7 | 6.2 | 2.8 | 193 | 511 |

Note: "-" Means blood transfusion and cesarean delivery services are not provided at union-level public facilities or at community clinics.

MVA = Manual vacuum aspiration.

¹ Antibiotics, oxytocin, and anticonvulsants.

 ² Antibiotics, oxytocin, anticonvulsants, assisted vaginal delivery, manual removal of placenta, removal of retained products of conception, and neonatal resuscitation.
 ³ Antibiotics, oxytocin, anticonvulsants, assisted vaginal delivery, manual removal of placenta, removal of retained products of conception, neonatal resuscitation, blood transfusion, and cesarean delivery.

Table 7.7: Readiness of health facilities to provide normal delivery services

Among facilities that offer normal delivery services, the percentages with 13 readiness items and the average readiness score, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Guide- lines on BEmOC ¹ or CEmOC ¹ | Staff trained in delivery care at any time ² | Exami- nation light ³ | Delivery pack ⁴ | Suction appa- ratus | Neonatal bag and mask | Parto- graph⁵ | Gloves ⁶ | Injectable uterotonic oxytocin | Injectable antibiotic | Mag- nesium sulphate | Skin dis- infectant | Intra- venous fluids with infusion set | Average readiness score (out of 13) ⁷ | Number of facilities offering normal delivery services (weighted) | Number of facilities offering normal delivery services (un- weighted) |
|--|--|---|--|--|--|--|--|--|--|--|--|--|--|---|---|--|
| Facility type | | | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 17.4 24.2 12.6 17.4 | 59.3 54.8 81.8 55.2 | 89.8 93.5 90.7 89.1 | 94.4 96.8 93.1 94.4 | 77.1 83.9 68.7 77.9 | 95.7 98.4 89.4 96.7 | 87.8 90.3 77.3 89.6 | 95.0 98.4 95.3 94.5 | 87.9 83.9 77.1 90.8 | 72.5 98.4 51.7 73.1 | 53.7 64.5 33.6 56.4 | 78.4 72.6 74.8 80.0 | 62.0 66.1 50.3 63.9 | 9.7 10.3 9.0 9.8 | 40 4 6 30 | 305 62 87 156 |
| Union-level public facilities UHFWC USC/RD | 11.5 10.7 16.8 | 38.8 38.3 42.1 | 67.7 68.2 64.9 | 90.6 91.8 82.8 | 29.2 29.0 30.6 | 79.1 79.8 74.3 | 40.0 39.4 44.1 | 86.5 86.9 83.7 | 39.2 38.7 42.4 | 4.4 3.4 11.5 | 16.8 16.6 17.8 | 49.9 50.3 47.0 | 29.1 27.4 40.6 | 5.8 5.8 6.0 | 153 133 20 | 206 170 36 |
| Community clinic (CC) | 0.0 | 56.5 | 63.6 | 31.8 | 0.0 | 26.3 | 5.1 | 70.5 | 0.0 | 0.0 | 0.0 | 19.6 | 4.7 | 2.8 | 46 | 24 |
| NGO clinic/ hospital | 29.0 | 60.8 | 92.9 | 94.6 | 83.4 | 94.6 | 91.5 | 100.0 | 96.1 | 69.7 | 68.3 | 89.2 | 63.8 | 10.3 | 8 | 39 |
| Private hospital | 3.5 | 22.9 | 91.2 | 86.3 | 88.0 | 81.8 | 20.7 | 92.5 | 82.0 | 72.2 | 61.9 | 82.5 | 51.9 | 8.4 | 163 | 166 |
| Location Urban Rural | 6.5 9.3 | 27.7 45.1 | 91.2 68.9 | 87.3 78.9 | 86.2 28.0 | 84.3 69.1 | 31.5 36.8 | 92.9 84.2 | 82.6 36.2 | 71.9 10.1 | 60.7 17.3 | 81.7 46.8 | 53.1 27.4 | 8.6 5.6 | 192 220 | 409 331 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 7.4 16.2 8.3 3.3 2.1 5.1 8.5 7.4 | 44.0 45.9 33.6 31.1 25.3 44.5 34.9 47.7 | 79.7 74.3 80.4 70.7 87.8 81.6 89.6 78.6 | 80.4 91.8 90.2 70.1 73.2 96.1 73.0 71.6 | 45.4 62.2 61.2 55.2 49.7 46.9 51.5 40.6 | 79.6 77.9 82.4 60.9 78.5 86.6 67.2 75.0 | 36.6 33.5 45.7 17.9 20.1 47.9 51.4 25.7 | 95.0 90.5 87.1 81.8 89.7 88.9 93.8 83.8 | 62.4 65.9 60.1 40.1 48.9 49.9 52.7 | 38.7 39.1 48.5 48.4 25.1 27.6 26.3 39.1 | 33.1 39.3 42.1 45.3 24.0 30.1 38.5 37.1 | 63.2 67.5 64.5 75.6 50.1 57.9 59.4 49.3 | 41.4 48.6 53.4 26.2 20.1 36.9 28.6 46.2 | 7.1 7.5 7.6 6.5 5.9 7.0 6.7 6.5 | 21 89 96 64 56 33 31 22 | 73 124 133 92 102 84 69 63 |
| Total | 8.0 | 37.0 | 79.3 | 82.8 | 55.1 | 76.2 | 34.3 | 88.2 | 57.8 | 38.9 | 37.5 | 63.1 | 39.4 | 7.0 | 412 | 740 |
| Total excluding CCs | 9.0 | 34.6 | 81.3 | 89.2 | 62.1 | 82.5 | 38.0 | 90.5 | 65.1 | 43.8 | 42.3 | 68.6 | 43.8 | 7.5 | 366 | 716 |
| Total public | 10.2 | 45.6 | 70.6 | 79.9 | 31.6 | 71.7 | 41.3 | 84.8 | 39.8 | 15.0 | 19.8 | 48.8 | 29.9 | 5.9 | 239 | 535 |
| Total excluding CCs | 12.7 | 43.1 | 72.3 | 91.4 | 39.2 | 82.5 | 49.9 | 88.3 | 49.3 | 18.6 | 24.5 | 55.8 | 36.0 | 6.6 | 193 | 511 |

¹ Basic emergency obstetric care (BEmOC) guidelines or comprehensive emergency obstetric care (CEmOC) guidelines.

² Facility has at least one interviewed staff member providing the service who reports receiving in-service training in delivery care. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ A functioning flashlight is acceptable.

 ³ A functioning flashlight is acceptable.
 ⁴ Either the facility has a sterile delivery pack available at the delivery site or all of the following individual equipment must be present: cord clamp, episiotomy scissors, scissors (or blade) to cut cord, suture material with needle, and needle holder.
 ⁵ A blank partograph is available at the service site.
 ⁶ Disposable latex gloves or equivalent are available at the service site.
 ⁷ Average readiness score is the average number of 13 items (guidelines on BEmOC or CEmOC, staff trained in delivery care any time, examination light, delivery pack, suction apparatus, neonatal bag and mask, partograph, gloves, oxytocin, injectable antibiotic, magnesium sulphate, skin disinfectant, and intravenous fluid with infusion set) available for providing normal delivery core of the service. providing normal delivery services.

Table 7.8: Newborn care practices

Among facilities offering normal delivery services, the percentages reporting that the indicated practice is a routine component of newborn care, by facility type, Bangladesh HFS 2022

| | | | | | Fa | acility type | | | | | | | | |
|--|--|-------|------|------|---|--------------|--------|----------------------------------|----------------------------|---------------------|-------|---------------------------|-----------------|-------------------------------------|
| Newborn care practices | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commu- nity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Routine complete (head-to-toe) examination of | | | | | | | | | | | | | | |
| newborns before discharge | 99.1 | 100.0 | 96.3 | 99.5 | 94.0 | 94.2 | 93.2 | 85.2 | 98.8 | 92.0 | 92.8 | 93.8 | 93.2 | 95.1 |
| Weighing the newborn immediately upon delivery | 99.2 | 100.0 | 97.6 | 99.5 | 81.9 | 85.4 | 58.7 | 50.5 | 100.0 | 96.1 | 86.1 | 90.6 | 78.8 | 85.5 |
| Administration of vitamin K to newborn | 94.1 | 96.8 | 79.6 | 96.8 | 50.3 | 50.3 | 50.7 | 31.6 | 88.4 | 99.0 | 72.6 | 77.8 | 54.1 | 59.5 |
| Number of facilities offering normal delivery services (weighted) Number of facilities offering normal delivery services | 40 | 4 | 6 | 30 | 153 | 133 | 20 | 46 | 10 | 163 | 412 | 366 | 239 | 193 |
| (unweighted) | 305 | 62 | 87 | 156 | 206 | 170 | 36 | 24 | 39 | 166 | 740 | 716 | 535 | 511 |
| | | | | | | | | | | | | | | |

Table 7.9: Essential medicines for newborn care

Among facilities that offer normal delivery services, the percentages with essential medicines for newborns observed to be available on the day of the survey, by facility type, Bangladesh HFS 2022

| | | | | | Facilit | y type | | | | | | | | |
|--|--|---------------|--------------|--------------|---|-------------|-------------|----------------------------------|----------------------------|---------------------|--------------|---------------------------|-----------------|-------------------------------------|
| Essential medicines for newborn care | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commu- nity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Antibiotic eye ointment for newborn | 63.3 | 64.5 | 4.8 | 75.6 | 9.7 | 6.3 | 32.6 | 88.9 | 2.8 | 15.8 | 26.1 | 18.1 | 34.0 | 20.9 |
| Injectable gentamicin Injectable ceftriaxone | 70.1 86.2 | 58.1 100.0 | 39.3 52.9 | 78.4 91.3 | 21.1 4.6 | 23.3 3.4 | 6.2 12.7 | 0.0 0.0 | 35.2 75.8 | 44.8 81.6 | 33.2 44.1 | 37.4 49.7 | 25.3 17.4 | 31.3 21.6 |
| Amoxicillin syrup/ suspension | 68.0 | 32.3 | 89.8 | 68.7 | 85.3 | 89.6 | 56.7 | 91.4 | 31.5 | 18.5 | 56.6 | 52.2 | 83.6 | 81.7 |
| Ampicillin injection | 33.0 | 11.3 | 1.2 | 43.1 | 0.8 | 0.4 | 3.5 | 0.0 | 3.4 | 9.3 | 7.3 | 8.2 | 6.1 | 7.5 |
| solution | 92.0 | 67.7 | 89.8 | 96.1 | 80.5 | 84.3 | 54.8 | 53.0 | 89.3 | 38.4 | 62.1 | 63.2 | 77.1 | 82.9 |
| Number of facilities offering normal delivery services (weighted) Number of facilities offering normal delivery services | 40 | 4 | 6 | 30 | 153 | 133 | 20 | 46 | 10 | 163 | 412 | 366 | 239 | 193 |
| (unweighted) | 305 | 62 | 87 | 156 | 206 | 170 | 36 | 24 | 39 | 166 | 740 | 716 | 535 | 511 |

Note: The essential medicines and antibiotic eye ointment for newborn measures presented in this table comprise the medicines domain for assessing readiness to provide basic obstetric care within the health facility assessment methodology proposed by the WHO and USAID (2012).

Table 7.10: Availability of equipment for newborn care services

Among facilities that offer normal delivery services, the percentages having indicated equipment, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Incubator | Suction apparatus with catheter | Suction bulb or penguin sucker | Newborn bag and mask | Timer | Infant scale | Fetal stetho- scope | Thermo- meter | Thermo- meter for low body temper- ature | Number of facilities offering normal delivery services (weighted) | Number of facilities offering normal delivery services (unweighted) |
|---|----------------------------------|--|---|--------------------------------------|--------------------------------------|------------------------------|--------------------------------------|--------------------------------------|--|---|---|
| Facility type | | | | | | | | | | | |
| District and upazila public facilities | 10.0 | 77.1 | 95.9 | 95.7 | 96.6 | 95.4 | 70.6 | 79.1 | 19.4 | 40 | 305 |
| DH MCWC UHC | 27.4 2.3 9.1 | 83.9 68.7 77.9 | 91.9 87.4 98.2 | 98.4 89.4 96.7 | 96.8 92.0 97.6 | 90.3 94.2 96.4 | 71.0 52.5 74.5 | 77.4 78.1 79.6 | 25.8 18.1 18.7 | 4 6 30 | 62 87 156 |
| Union-level public facilities UHFWC USC/RD | 1.4 1.5 1 1 | 29.2 29.0 30.6 | 76.3 78.6 | 79.1 79.8 74.3 | 83.7 83.1 87.6 | 75.4 78.8 52.9 | 22.3 22.5 21.3 | 75.4 76.4 68.5 | 18.6 18.0 22.5 | 153 133 20 | 206 170 36 |
| Community clinic (CC) | 0.0 | 0.0 | 42.7 | 26.3 | 84.0 | 59.0 | 9.6 | 65.9 | 20.7 | 46 | 24 |
| NGO clinic/hospital | 10.4 | 83.4 | 96.6 | 94.6 | 88.7 | 97.8 | 82.5 | 88.4 | 16.6 | 8 | 39 |
| Private hospital | 13.4 | 88.0 | 85.4 | 81.8 | 56.0 | 88.8 | 49.5 | 83.8 | 29.8 | 163 | 166 |
| Location Urban Rural | 13.0 1.8 | 86.2 28.0 | 86.4 71.6 | 84.3 69.1 | 62.7 84.1 | 89.5 74.2 | 54.1 23.5 | 83.1 74.2 | 28.9 18.4 | 192 220 | 409 331 |
| Division Barishal Chattogram Dhaka Khulna Pajebabi | 8.0 10.8 9.8 1.8 2.6 | 45.4 62.2 61.2 55.2 | 74.2 76.6 76.7 86.7 78.0 | 79.6 77.9 82.4 60.9 78 5 | 73.3 81.2 72.6 51.7 81 2 | 81.2 77.4 83.5 78.3 | 42.6 47.5 49.2 11.4 36 7 | 86.4 85.1 83.8 55.8 80.1 | 17.7 34.0 22.9 14.4 26.7 | 21 89 96 64 | 73 124 133 92 102 |
| Rangpur Sylhet Mymensingh | 4.7 10.5 4.2 | 46.9 51.5 40.6 | 80.6 67.1 89.5 | 86.6 67.2 75.0 | 81.9 76.4 85.6 | 82.7 79.5 69.9 | 34.0 32.0 37.5 | 83.6 84.1 64.3 | 14.1 25.0 15.4 | 33 31 22 | 84 69 63 |
| Total | 7.0 | 55.1 | 78.5 | 76.2 | 74.2 | 81.4 | 37.8 | 78.3 | 23.3 | 412 | 740 |
| Total excluding CCs | 7.9 | 62.1 | 83.1 | 82.5 | 72.9 | 84.2 | 41.3 | 79.9 | 23.6 | 366 | 716 |
| Total public | 2.6 | 31.6 | 73.1 | 71.7 | 85.9 | 75.6 | 28.0 | 74.2 | 19.2 | 239 | 535 |
| Total public excluding CCs | 3.2 | 39.2 | 80.4 | 82.5 | 86.4 | 79.6 | 32.4 | 76.2 | 18.8 | 193 | 511 |

Note: The essential equipment items for newborn care presented in this table comprise the medicines domain for assessing readiness to provide basic obstetric care within the health facility assessment methodology proposed by the WHO and USAID (2012).

Table 7.10.1: Newborn signal functions

Among facilities offering normal delivery services, the percentages reporting that they performed the newborn signal functions at least once during the three months before the survey, by background characteristics, Bangladesh HFS 2022

| | | | | | Facili | ty type | | | | | | | | |
|---|--|-------|------|------|---|---------|--------|-------------------------------|----------------------------|---------------------|-------|---------------------------|-----------------|-------------------------------------|
| Newborn signal functions | District and upazila public facilities | DH | MCWC | UHC | Union- level public facilities | UHFWC | USC/RD | Commu- nity clinic (CC) | NGO clinic/ hospital | Private hospital | Total | Total excluding CCs | Total public | Total public excluding CCs |
| Iron and folic acid supplementation in pregnant women | 96.0 | 95.2 | 94.2 | 96.5 | 88.3 | 86.9 | 98.0 | 87.0 | 92.9 | 73.7 | 83.3 | 82.8 | 89.4 | 89.9 |
| Hand wash each time before touching the baby | 98.8 | 100.0 | 97.7 | 98.8 | 83.2 | 83.9 | 78.5 | 70.1 | 96.7 | 97.6 | 89.3 | 91.7 | 83.3 | 86.5 |
| Drying of the baby immediately after birth | 98.4 | 100.0 | 96.5 | 98.6 | 84.3 | 84.2 | 84.8 | 70.1 | 96.7 | 97.8 | 89.7 | 92.2 | 83.9 | 87.2 |
| Delayed umbilical cord clamping | 97.9 | 100.0 | 97.7 | 97.6 | 83.3 | 84.6 | 75.1 | 54.6 | 96.7 | 91.3 | 85.0 | 88.8 | 80.2 | 86.4 |
| Clean cord cutting with sterile blade | 94.5 | 93.5 | 96.6 | 94.3 | 83.2 | 83.5 | 81.3 | 66.2 | 96.7 | 90.9 | 85.8 | 88.2 | 81.8 | 85.6 |
| Single application of 7.1% CHX | 98.4 | 98.4 | 94.0 | 99.3 | 86.0 | 86.3 | 83.6 | 70.1 | 100.0 | 88.5 | 86.7 | 88.8 | 85.0 | 88.6 |
| Neonatal resuscitation | 90.4 | 93.6 | 74.2 | 93.4 | 50.1 | 49.6 | 52.8 | 29.7 | 86.6 | 80.9 | 64.8 | 69.2 | 52.9 | 58.5 |
| Perform skin to skin care immediately after birth | 99.1 | 98.4 | 98.8 | 99.3 | 85.3 | 86.2 | 79.1 | 65.1 | 96.7 | 95.3 | 88.6 | 91.6 | 83.7 | 88.2 |
| Early initiation of breast feeding | 99.0 | 98.4 | 98.8 | 99.1 | 83.8 | 83.7 | 84.8 | 70.1 | 96.7 | 97.6 | 89.5 | 92.0 | 83.7 | 87.0 |
| Immunize patient with tetanus toxoid | 71.2 | 69.4 | 52.8 | 75.3 | 40.6 | 37.9 | 58.6 | 48.8 | 68.0 | 51.6 | 49.5 | 49.6 | 47.3 | 47.0 |
| Administer oral antibiotic | 58.8 | 77.4 | 55.2 | 56.7 | 28.3 | 27.3 | 34.9 | 7.8 | 66.6 | 63.7 | 43.9 | 48.4 | 29.4 | 34.6 |
| Administer intramuscular antibiotic | 54.5 | 71.0 | 29.0 | 57.5 | 8.4 | 8.2 | 9.9 | 4.6 | 45.7 | 59.7 | 33.6 | 37.3 | 15.4 | 18.0 |
| Suction of airway of newborn | 90.4 | 98.4 | 81.3 | 91.2 | 57.0 | 57.3 | 55.2 | 26.8 | 88.3 | 89.1 | 70.3 | 75.8 | 56.8 | 64.0 |
| Administer intravenous antibiotic | 64.9 | 85.5 | 28.9 | 69.4 | - | - | - | - | 42.2 | 71.7 | 69.1 | - | 64.9 | - |
| Administer antibiotic for premature rupture of membrane | 82.2 | 98.4 | 63.1 | 83.9 | - | - | - | - | 67.0 | 71.1 | 73.0 | - | 82.2 | - |
| Oxygen therapy for management of newborn infection and respiratory problem | 93.0 | 100.0 | 68.8 | 97.1 | - | - | - | - | 85.6 | 86.8 | 87.9 | - | 93.0 | - |
| KMC for preterm or LBW newborn | 64.0 | 83.9 | 33.5 | 67.5 | - | - | - | - | 41.8 | 29.5 | 36.6 | - | 64.0 | - |
| Phototherapy for term or preterm with hyperbilirubinemia | 24.0 | 80.6 | 4.4 | 19.7 | - | - | - | - | 18.6 | 25.3 | 24.7 | - | 24.0 | - |
| Incubator support for sick term baby, preterm baby or if the baby receiving KMC becomes sick | 14.6 | 56.5 | 4.5 | 10.5 | - | - | - | - | 3.3 | 16.8 | 15.8 | - | 14.6 | - |
| Corticosteroids for pre- term labor | 64.8 | 91.9 | 40.2 | 66.0 | - | - | - | - | 59.1 | 68.3 | 67.2 | - | 64.8 | - |
| Primary NSFs ¹ | 77.0 | 82.3 | 64.9 | 78.8 | 43.0 | 43.9 | 37.0 | 15.0 | 76.2 | 50.7 | 47.0 | 51.1 | 43.3 | 50.1 |
| Basic NSFs ² | 27.9 | 38.7 | 12.9 | 29.5 | 2.4 | 2.1 | 4.0 | - | 33.0 | 21.5 | 14.5 | - | 7.7 | - |
| Comprehensive NSFs ³ | 10.7 | 30.7 | 2.3 | 9.5 | - | - | - | - | 8.6 | 6.9 | 7.7 | - | 10.7 | - |
| Advanced NSFs ⁴ | 5.3 | 21.0 | 1.1 | 3.8 | - | - | - | - | 0.0 | 3.8 | 3.9 | - | 5.3 | - |
| Number of facilities offering normal delivery services (weighted) | 40 | 4 | 6 | 30 | 153 | 133 | 20 | 46 | 10 | 163 | 412 | 366 | 239 | 193 |
| offering normal delivery | 305 | 62 | 87 | 156 | 206 | 170 | 36 | 24 | 39 | 166 | 740 | 716 | 535 | 511 |

Note: "-" Means facilities do not have the provision to perform the indicated newborn signal function.

The facilities that do not have the provision to perform the indicated newborn signal function are excluded from the denominator.

NSF= newborn signal function.

¹ Primary newborn signal functions (all 9 NSFs): Iron and folic acid supplementation in pregnant women, hand wash each time before touching the baby, drying of the baby immediately after birth, delayed umbilical cord clamping, clean cord cutting sterile blade, single application of 7.5% CHX, neonatal resuscitation, perform skin-to-skin care immediately after birth, and ²Basic newborn signal functions (all 13 NSFs): Primary newborn signal functions and Immunize patient with tetanus toxoid, administer oral antibiotic, administer intramuscular antibiotic,

and suction of airway of newborn.

³ Comprehensive newborn signal function (all 18 NSFs): Basic newborn signal functions and administer intravenous antibiotic, administer antibiotic for premature rupture of membrane, oxygen therapy for management of newborn infection and respiratory problem, KMC for pre-term of LBW newborn, phototherapy for term or pre-term with hyperbilirubinemia.
⁴ Advanced newborn signal function (all 20 NSFs): Comprehensive newborn signal functions and incubator support for sick term baby, pre-term baby or if the baby receiving KMC

becomes sick, and corticosteroids for pre-term labor.

Table 7.11: Supportive management for providers of delivery care

Among interviewed providers of normal delivery or newborn care services, the percentages who report receiving training related to their work and personal supervision during the specified time periods, by background characteristics, Bangladesh HFS 2022

| | Percentage of | interviewed provider: | s who received: | | |
|--|--|--|--|--|--|
| Background characteristic | Training related to delivery and/or newborn care during the 24 months preceding the survey ¹ | Personal supervision during the 6 months preceding the survey ² | Training related to delivery and/or newborn care during the 24 months and personal supervision during the 6 months preceding the survey | Number of interviewed providers of normal delivery or newborn care services (weighted) | Number of interviewed providers of normal delivery or newborn care services (unweighted) |
| Facility type | | | | | |
| District and upazila public facilities DH MCWC UHC | 20.9 16.6 24.0 22.3 | 97.3 96.5 97.1 97.5 | 20.6 16.5 23.3 21.9 | 963 237 25 701 | 2,114 420 328 1,366 |
| Union-level public facilities UHFWC USC/RD | 18.6 18.8 17.3 | 93.6 93.1 97.0 | 17.5 17.5 17.3 | 179 154 25 | 334 267 67 |
| Community clinic (CC) | 22.4 | 100.0 | 22.4 | 60 | 30 |
| NGO clinic/hospital | 25.8 | 98.0 | 25.3 | 59 | 147 |
| Private hospital | 10.4 | 88.4 | 9.2 | 823 | 427 |
| Location Urban Rural | 15.3 20.5 | 93.1 94.6 | 14.6 19.8 | 1,514 570 | 1,993 1,059 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 12.0 20.3 12.1 10.3 17.2 23.9 31.2 18.3 | 96.6 91.4 94.6 89.1 96.8 89.6 98.3 95.9 | 12.0 20.1 11.3 9.3 15.9 23.4 30.6 17.0 | 125 403 605 234 270 215 108 124 | 352 533 542 352 376 391 243 263 |
| Total | 16.7 | 93.5 | 16.0 | 2,084 | 3,052 |
| Total excluding CCs | 16.6 | 93.3 | 15.8 | 2,024 | 3,022 |
| Total public | 20.7 | 96.9 | 20.2 | 1,202 | 2,478 |
| Total excluding CCs | 20.6 | 96.7 | 20.1 | 1,142 | 2,448 |

¹ Training here refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.
² Personal supervision refers to any form of technical support or supervision from a facility-based supervisor or from a visiting supervisor. It may include, but is not limited to, review of records and observation of work, with or without any feedback to the health worker.

Table 7.12: Training for providers of normal delivery services: Immediate newborn care

Among interviewed providers of normal delivery or newborn care services, the percentages who report receiving in-service training on topics related to delivery and newborn care during the 24 months preceding the survey and at any time, by background characteristics, Bangladesh HFS 2022

| | Percentag | Number of | | | | | | | | | | |
|------------------------------|-----------------------------------|-------------|---------------------------------|-------------|---------------------------------|-------------|---------------------------------|-------------|--|-------------|--|---|
| Background characteristic | Early and exclusive breastfeeding | | Newborn infection management | | Thermal care | | Sterile cord cutting and care | | Kangaroo mother care for low birth weight babies | | interviewed providers of normal delivery or | Number of interviewed providers of normal |
| | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | newborn care services (weighted) | delivery or newborn care services (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public | | | | | | | | | | | | |
| facilities | 9.1 | 32.9 | 4.6 | 15.2 | 8.4 | 25.8 | 8.1 | 28.9 | 11.4 | 31.0 | 963 | 2,114 |
| DH | 7.7 | 34.1 | 3.4 | 16.6 | 6.2 | 26.3 | 5.8 | 29.4 | 8.4 | 30.3 | 237 | 420 |
| MCWC | 9.5 | 58.5 | 5.1 | 23.8 | 9.8 | 46.5 | 9.5 | 52.1 | 13.9 | 53.0 | 25 | 328 |
| UHC | 9.6 | 31.6 | 4.9 | 14.5 | 9.1 | 25.0 | 8.8 | 27.8 | 12.4 | 30.5 | 701 | 1,366 |
| Union-level | | | | | | | | | | | | |
| public facilities | 10.1 | 48.9 | 6.3 | 23.4 | 10.3 | 43.9 | 11.7 | 48.3 | 10.4 | 45.8 | 179 | 334 |
| UHFWC | 10.1 | 52.0 | 6.3 | 25.1 | 10.4 | 47.8 | 11.8 | 51.6 | 11.0 | 49.8 | 154 | 267 |
| USC/RD | 9.8 | 29.6 | 6.1 | 12.9 | 9.7 | 19.4 | 11.0 | 28.2 | 6.1 | 21.1 | 25 | 67 |
| Community clinic (CC) | 17.9 | 66.7 | 0.0 | 6.0 | 15.7 | 63.0 | 2.1 | 52.5 | 13.4 | 62.2 | 60 | 30 |
| NGO clinic/hospital | 17.6 | 39.0 | 10.2 | 23.4 | 16.2 | 32.6 | 19.2 | 37.5 | 17.3 | 37.3 | 59 | 147 |
| Private hospital | 6.8 | 15.9 | 4.5 | 12.1 | 3.7 | 13.3 | 3.2 | 14.1 | 5.1 | 11.1 | 823 | 427 |
| Location | | | | | | | | | | | | |
| Urban | 8.5 | 24.6 | 4.6 | 13.8 | 6.1 | 19.8 | 5.6 | 22.0 | 7.8 | 20.9 | 1.514 | 1,993 |
| Rural | 9.6 | 39.6 | 4.9 | 17.1 | 9.8 | 34.0 | 9.3 | 35.4 | 12.5 | 37.7 | 570 | 1,059 |
| Division | | | | | | | | | | | | |
| Barishal | 4.6 | 24.5 | 24 | 10.9 | 4.0 | 20.9 | 3.6 | 21.5 | 66 | 23.7 | 125 | 352 |
| Chattogram | 12.4 | 32.2 | 5.6 | 16.8 | 11 7 | 20.0 | 10.1 | 28.7 | 12.3 | 29.6 | 403 | 533 |
| Dhaka | 8.2 | 19.9 | 5.8 | 12.1 | 4.7 | 13.7 | 5.0 | 17.7 | 5.3 | 16.0 | 605 | 542 |
| Khulna | 4.1 | 33.4 | 1.8 | 19.8 | 2.5 | 30.8 | 3.5 | 33.3 | 5.5 | 31.8 | 234 | 352 |
| Rajshahi | 7.8 | 31.0 | 2.1 | 11.8 | 6.8 | 28.5 | 5.5 | 28.4 | 10.2 | 24.9 | 270 | 376 |
| Rangpur | 8.2 | 29.9 | 5.1 | 14.0 | 10.2 | 24.8 | 8.0 | 24.8 | 13.8 | 30.8 | 215 | 391 |
| Sylhet | 14.7 | 44.4 | 7.8 | 22.8 | 10.0 | 35.0 | 11.0 | 37.8 | 14.3 | 36.5 | 108 | 243 |
| Mymensingh | 11.2 | 35.0 | 6.6 | 14.6 | 9.0 | 28.9 | 8.8 | 29.1 | 11.0 | 30.8 | 124 | 263 |
| Total | 8.8 | 28.7 | 4.7 | 14.7 | 7.1 | 23.7 | 6.6 | 25.6 | 9.1 | 25.5 | 2,084 | 3,052 |
| Total excluding CCs | 8.5 | 27.6 | 4.8 | 14.9 | 6.9 | 22.6 | 6.7 | 24.8 | 8.9 | 24.4 | 2,024 | 3,022 |
| Total public | 9.7 | 36.9 | 4.6 | 16.0 | 9.1 | 30.4 | 8.4 | 32.9 | 11.4 | 34.8 | 1,202 | 2,478 |
| Total excluding CCs | 9.3 | 35.4 | 4.9 | 16.5 | 8.7 | 28.7 | 8.7 | 31.9 | 11.3 | 33.4 | 1,142 | 2,448 |

Note: Training here refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

Table 7.13: Training for providers of normal delivery services: Immediate newborn care

Among interviewed providers of normal delivery or newborn care services, the percentages who report receiving in-service training on topics related to delivery and newborn care during the 24 months preceding the survey and at any time, by background characteristics, Bangladesh HFS 2022

| | Per | centage of | interviewed | providers | of normal de | elivery or n | ewborn car | e service | s who repo | rt receivir | ng in-servic | e training | in: | | |
|--|--|----------------|------------------------------------|----------------|---|----------------|---|----------------|------------------------------------|----------------|------------------------------------|----------------|--|---|---|
| | Newborn resuscitation using bag and mask | | Essential newborn care | | Umbilical cord care (use of 7.1% chlorhexidine) | | Emergency triage assessment training (ETAT) | | IMCI guidelines (0-59 days) | | Comprehensive newborn care | | Training on different services ¹ | Number of | Number of |
| Background | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | During the past 24 months | At any time | At any time | providers of normal delivery or newborn care services (weighted) | providers of normal delivery or newborn care services (unweighted) |
| Facility type | | | | | | | | | | | | | | | |
| District and upazila public facilities | 10.7 | 33.8 | 8.5 | 24.4 | 8.7 | 29.4 | 4.8 | 14.8 | 4.4 | 16.3 | 6.1 | 14.9 | 10.9 | 963 | 2,114 |
| DH | 8.5 | 38.3 | 7.2 | 26.9 | 5.8 | 31.6 | 6.3 | 19.2 | 5.1 | 14.1 | 6.1 | 15.2 | 12.5 | 237 | 420 |
| MCWC | 11.4 | 56.2 | 8.4 | 44.8 | 8.8 | 51.3 | 4.6 | 17.0 | 3.3 | 22.8 | 7.2 | 35.0 | 24.7 | 25 | 328 |
| UHC | 11.4 | 31.5 | 8.9 | 22.9 | 9.7 | 27.9 | 4.3 | 13.3 | 4.3 | 16.8 | 6.0 | 14.1 | 9.9 | 701 | 1,366 |
| Union-level public facilities | 10.4 | 47.8 | 10.0 | 40.6 | 12.1 | 51.2 | 4.7 | 17.1 | 3.6 | 26.5 | 9.1 | 35.9 | 23.3 | 179 | 334 |
| UHFWC | 10.3 | 50.6 | 10.3 | 43.1 | 12.2 | 54.8 | 4.6 | 17.5 | 3.5 | 28.3 | 10.0 | 38.8 | 24.8 | 154 | 267 |
| USC/RD | 11.1 | 30.9 | 8.1 | 25.0 | 11.3 | 29.0 | 5.6 | 15.0 | 4.7 | 15.7 | 3.4 | 17.7 | 14.2 | 25 | 67 |
| Community clinic (CC) | 2.1 | 37.3 | 0.0 | 17.8 | 13.9 | 62.2 | 0.0 | 7.0 | 9.4 | 25.1 | 2.2 | 29.6 | 9.5 | 60 | 30 |
| NGO clinic/ hospital | 21.3 | 38.5 | 17.7 | 36.1 | 19.5 | 36.9 | 7.6 | 16.1 | 11.6 | 25.2 | 9.1 | 19.5 | 18.1 | 59 | 147 |
| Private hospital | 4.6 | 13.0 | 3.7 | 8.7 | 3.4 | 9.9 | 1.7 | 4.3 | 1.4 | 3.8 | 1.7 | 7.0 | 5.9 | 823 | 427 |
| Location | | | | | | | | | | | | | | | |
| Urban | 7.8 | 23.7 | 6.5 | 17.6 | 6.2 | 19.8 | 3.5 | 10.2 | 3.1 | 10.0 | 4.1 | 11.4 | 9.1 | 1,514 | 1,993 |
| Rural | 9.8 | 35.7 | 7.5 | 25.5 | 10.3 | 37.9 | 3.4 | 12.0 | 4.7 | 19.9 | 5.7 | 21.4 | 13.0 | 570 | 1,059 |
| Division Barishal | 45 | 26.5 | 36 | 17.6 | 33 | 20.8 | 16 | 10.4 | 0.8 | 14.2 | 23 | 10.8 | 86 | 125 | 352 |
| Chattogram | 8.8 | 26.8 | 9.7 | 23.5 | 10.3 | 29.3 | 6.4 | 15.3 | 6.1 | 14.5 | 4.8 | 15.7 | 11.9 | 403 | 533 |
| Dhaka | 6.9 | 18.9 | 4.5 | 14.7 | 5.6 | 17.2 | 3.2 | 6.7 | 3.6 | 9.0 | 3.9 | 10.7 | 8.9 | 605 | 542 |
| Khulna | 6.4 | 34.5 | 3.8 | 21.9 | 4.1 | 30.7 | 2.6 | 13.8 | 2.3 | 16.0 | 2.7 | 17.3 | 12.9 | 234 | 352 |
| Rajshahi | 8.8 | 30.9 | 7.2 | 18.0 | 7.2 | 22.8 | 1.8 | 8.5 | 3.0 | 13.6 | 5.2 | 13.1 | 8.3 | 270 | 376 |
| Rangpur | 11.0 | 30.6 | 8.6 | 23.6 | 8.9 | 26.6 | 3.6 | 12.5 | 2.8 | 12.5 | 7.0 | 15.5 | 10.5 | 215 | 391 |
| Sylhet | 13.9 | 38.1 | 10.5 | 30.1 | 10.8 | 34.7 | 4.1 | 13.6 | 3.2 | 14.6 | 5.3 | 17.5 | 10.1 | 108 | 243 |
| wymensingn | 10.7 | 29.8 | 9.3 | 16.0 | 11.0 | 31.4 | 2.4 | 8.9 | 2.0 | 14.0 | 0.3 | 20.0 | 10.8 | 124 | 203 |
| Total | 8.3 | 27.0 | 6.7 | 19.8 | 7.3 | 24.7 | 3.5 | 10.7 | 3.5 | 12.7 | 4.6 | 14.1 | 10.2 | 2,084 | 3,052 |
| Total excluding CCs | 8.5 | 26.7 | 6.9 | 19.8 | 7.1 | 23.6 | 3.6 | 10.8 | 3.4 | 12.4 | 4.6 | 13.7 | 10.2 | 2,024 | 3,022 |
| Total public | 10.2 | 36.1 | 8.3 | 26.5 | 9.5 | 34.3 | 4.5 | 14.8 | 4.6 | 18.2 | 6.3 | 18.8 | 12.7 | 1,202 | 2,478 |
| Total excluding CCs | 10.7 | 36.0 | 8.7 | 27.0 | 9.2 | 32.8 | 4.8 | 15.2 | 4.3 | 17.9 | 6.5 | 18.2 | 12.9 | 1,142 | 2,448 |

Note: Training here refers only to in-service training. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision. ¹ Training on Newborn resuscitation, essential newborn care, umbilical cord care, and comprehensive newborn care

Table 7.14: Key indicators for delivery and newborn care services in MCWCs

Among facilities offering normal delivery services, the percentages with key indicators, Bangladesh HFS 2022

| | Facility type | | | | |
|--|---|---|--|--|--|
| Indicators | MCWC (district-level) | MCWC (upazila and union-level) | | | |
| Availability | | | | | |
| Normal delivery service | 98.5 | 68.3 | | | |
| Number of facilities (weighted) | 5 | 3 | | | |
| Number of facilities (unweighted) | 62 | 38 | | | |
| Readiness | | | | | |
| Guidelines on BEmOC ¹ or CEmOC ¹ Staff trained in delivery care at any time ² Examination light ³ Delivery pack ⁴ Suction apparatus Neonatal bag and mask Partograph ⁵ Gloves ⁶ Injectable uterotonic oxytocin Injectable uterotonic oxytocin Injectable antibiotic Magnesium sulphate Skin disinfectant Intravenous fluids with infusion set Average readiness score ⁷ | 13.2 86.9 95.0 98.4 76.8 95 90.3 95.0 85.4 63.8 39.8 78.9 49.2 9.7 | 11.4 69.8 80.4 80.5 49.6 76.2 46.5 96.0 57.5 23.3 19.0 64.9 52.9 7.3 | | | |
| Functionality BEMOC signal function CEMOC signal function Primary signal function Basic signal function Comprehensive signal function Advance signal function | 17.9 6.3 70.2 18.3 3.2 1.6 | 7.4 0.0 52.3 0.0 0.0 0.0 | | | |
| Number of facilities offering normal delivery services (weighted) | 4 | 2 | | | |
| Number of facilities offering normal delivery services (unweighted) | 61 | 26 | | | |

¹ Basic emergency obstetric care (BEmOC) guidelines or comprehensive emergency obstetric care (CEmOC) guidelines.
 ² Facility has at least one interviewed staff member providing the service who reports

² Facility has at least one interviewed staff member providing the service who reports receiving in-service training in delivery care. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ A functioning flashlight is acceptable

⁴ Either the facility has a sterile delivery pack available at the delivery site or all of the following individual equipment must be present: cord clamp, episiotomy scissors, scissors (or blade) to cut cord, suture material with needle, and needle holder. ⁵ A blank partograph available at the service site.

⁶ Disposable latex gloves or equivalent available at the service site.

⁷ Average readiness score is the average number of items (out of 13 items) available for providing normal delivery services.

NON-COMMUNICABLE DISEASES

Key Findings

Diabetes

- Forty-five percent of public health facilities offer services related to diabetes, including the diagnosis, prescription of treatment, and management of patients with diabetes (Table 8.1 and Figure 8.1).
- Diabetes services are near universal in DHs (98%) and UHCs (99%), whereas in unionlevel facilities their provision is considerably lower (38%). Approximately two-thirds of the NGO clinics (78%) and private hospitals (70%) offer diabetes services (**Table 8.1** and **Figure 8.1**).
- Among facilities offering diabetes services, more than eight in 10 public facilities (85%) have a blood pressure apparatus and functional adult weighing scale (86%) (Table 8.1 and Figure 8.2).
- Among facilities offering both diagnosis and treatment services for diabetes 21% have blood glucose diagnostic capacity, and about one-third have both urine protein and urine glucose diagnostic capacity (Table 8.2 and Figure 8.3).
- Among facilities offering both diagnosis and treatment services for diabetes, 37% of public facilities have Metformin available—an increase of 18 percentage points from 2017. A similar increase was observed for injectable insulins in public facilities, from 17% in 2017 to 37% in 2022. (Table 8.2 and Figure 8.3).

Cardiovascular Diseases

- One-fourth of public facilities offer services for cardiovascular diseases (CVDs). This service is universally available among DHs (98%) and UHCs (97%), while 26% of union-level facilities offer these services. Additionally, 59% of NGO clinics and 67% of private hospitals offer CVD services (Table 8.3 and Figure 8.4).
- Among the public facilities that offer CVD services, 12% have guidelines for the diagnosis and management of CVDs and 22% have staff with recent training in CVD services (Table 8.3 and Figure 8.5).
- Ninety-three percent of the public facilities that offer CVD services have a stethoscope, 89% have a functioning blood pressure apparatus, and 86% have a functioning adult weighing scale (Table 8.3 and Figure 8.5).
- The availability of essential medicines in public health facilities offering CVD services increased from 2017 to 2022, with aspirin rising from 13% to 46%, beta blockers from 16% to 43%, and calcium channel blockers rising from 20% to 40% (Table 8.4 and Figure 8.6).
- Among facilities offering CVD services, 52% of DHs, 54% of UHCs, and 76% of private hospitals have oxygen available (Table 8.4).

Chronic Respiratory Diseases

- Fifty-one percent of public facilities provide services for chronic respiratory diseases (**Table 8.5** and **Figure 8.7**).
- Ninety-eight percent of both DHs and UHCs offer services for chronic respiratory diseases, while 48% of union-level facilities provide these services. Additionally, 64% of

NGO clinics and 66% of private hospitals offer services related to chronic respiratory diseases (**Table 8.5** and **Figure 8.7**).

- Among the public facilities providing services for chronic respiratory diseases, 15% have guidelines for diagnosis and management of chronic respiratory diseases and 19% have staff with recent training related to chronic respiratory diseases (Table 8.5 and Figure 8.8).
- Ninety percent of public facilities that offer services for chronic respiratory diseases have a stethoscope. The availability of peak flow meters and spacers for inhalers is low at district and upazila-level public facilities (6% and 10%, respectively); however, it is much higher among private facilities (24%) (Table 8.5 and Figure 8.8).
- Among the essential medicines for treatment of chronic respiratory diseases, salbutamol inhalers/tablets are widely available (92%) in public facilities that provide services for chronic respiratory diseases (Table 8.6 and Figure 8.9).
- Forty-nine percent of DHs, 54% of UHCs, and 76% of private hospitals have oxygen available (Table 8.6).

Hypertension

- Fifty-four percent of public facilities offer hypertension-related services (Table 8.7 and Figure 8.10).
- Ninety-seven percent of DHs and 99% of UHCs are offering services related to hypertension, while only 50% of union-level facilities offer these services. Additionally, 75% of NGO clinics and 70% of private hospitals offer hypertension-related services (Table 8.7 and Figure 8.10).
- Seventeen percent of public facilities that provide services related to hypertension have guidelines for hypertension, and only 20% have staff with recent training in hypertension (Table 8.7 and Figure 8.11).
- Ninety-one percent of public facilities that offer hypertension-related services have a stethoscope, 88% have a blood pressure apparatus, and only 84% have an adult weighing scale (Table 8.7 and Figure 8.11).
- Angiotensin receptor blockers and calcium channel blockers are widely available in DHs (88% and 70%), UHCs (91% and 71%), and private hospitals (48% and 54%) (Table 8.8).

Cervical Cancer

- Overall, 4% of public facilities in Bangladesh offer diagnosis/screening services for cervical cancer (Table 8.9 and Figure 8.13).
- Eighty-six percent of DHs and 89% of UHCs offer diagnosis/screening services for cervical cancer, while only 1% of union-level facilities offer these services. Additionally, 21% of NGO clinics and 10% of private hospitals offer diagnosis/screening services for cervical cancer (Table 8.9 and Figure 8.13).
- Among the public facilities that offer cervical cancer services, 26% have guidelines available for screening and 49% have staff with recent training in cervical cancer diagnosis/screening services (Table 8.9 and Figure 8.14).

8.1 BACKGROUND

oncommunicable diseases (NCDs), also known as chronic diseases, encompass a cluster of diseases or health conditions that tend to be of long duration. They arise from a combination of genetic, physiological, environmental, and behavioral factors. The main types of NCDs are cardiovascular diseases (such as heart attack and stroke), cancers, chronic respiratory diseases (such as chronic obstructive pulmonary disease and asthma), and diabetes (WHO 2023).

NCDs have a disproportionate impact on people living in low- and middle-income countries, such as Bangladesh, where more than three-quarters of global annual NCD deaths (31.4 million) occur. There is a common misconception that NCDs affect only older populations. However, evidence suggests that each year, 17 million NCD-related deaths occur globally before the age of 70 years (IHME n.d.).

Risk factors contributing to NCDs, such as unhealthy diets, physical inactivity, exposure to tobacco smoke, alcohol abuse, or air pollution, affect people from all walks of life, making them particularly vulnerable to NCD-related conditions. Forces such as rapid unplanned urbanization, the globalization of unhealthy lifestyles, and an aging population are all factors driving growth of these diseases. Elevated blood pressure and blood lipids, increased blood glucose, and obesity may occur in individuals as a result of unhealthy diets and inadequate physical activity. These are referred to as metabolic risk factors and can contribute to the development of cardiovascular disease, ultimately resulting in NCDs and leading to premature deaths (WHO 2023).

Each year, Bangladesh experiences 580,000 deaths from NCDs, which accounts for 67% of the total number of deaths in the country. This is a significant social and economic development issue. NCDs can result in the loss of the primary income earner within families, potentially resulting in both poverty and the withdrawal of family members from employment and educational opportunities to care for affected loved ones. The continuing care, and often expensive treatment interventions, required for these diseases can result in immense pressure on the health system and potentially catastrophic costs to families as a result of out-of-pocket expenditures. From 2011–2025, total global economic losses among low- and middle-income countries stemming from the four major NCDs (heart disease, cancer, chronic respiratory disease, and diabetes) are estimated at US\$7 trillion. It is vital to prevent and reduce morbidity and mortality from NCDs to attain sustainable development (DGHS 2018).

NCDs (inclusive of injuries) account for 61% of the total disease burden in the country, as compared with 39% from communicable diseases, maternal and child health, and nutrition combined (World Bank 2011). However, the health system is not prepared to tackle this challenge. For example, Bangladesh had an estimated 7.7 qualified health care providers (Ahmed 2011) per 10,000 population, whereas the WHO recommends 25.0 per 10,000 population. In addition, physicians and nurses account for only 5% of all health care providers, as most of these individuals in these roles receive only informal training and serve the needs of the majority, particularly the economically disadvantaged (MOHFW and WHO Bangladesh 2021). Until recently, few health workers received training in NCD prevention and management, though the number has increased since 2013 (Kabir 2022).

The growing burden of NCDs compounds increases the burden on already stressed health systems, disproportionately affecting the most impoverished populations and hindering the social and economic development of countries (Islam et al. 2014). Addressing NCDs and improving health outcomes are essential for interventions to sustainably impact development and to achieve the targets outlined in the Sustainable Development Goals (SDGs). Failure to address these issues could compound existing health challenges and affect the overall development of countries, including those with emerging economies like Bangladesh (Pullar et al. 2018). According to a new report by the WHO, the world's most economically challenged countries have the potential to gain US\$350 billion by 2030 through scaled up investments in the prevention and treatment of

chronic diseases, such as heart disease and cancer, with an additional cost of only US\$1.27 per person annually. Such actions would result in more than 8 million lives saved over the same period (WHO 2018).

Using the information in the 2022 BHFS, this chapter addresses questions about the availability of services for selected NCDs. The chapter is organized as follows:

- **Diabetes:** Section 8.2, including **Tables 8.1–8.2** and **Figures 8.1–8.3**, presents information on the availability of diabetes diagnostic and treatment services.
- **Cardiovascular diseases:** Section 8.3, including **Tables 8.3–8.4** and **Figures 8.4–8.6** presents information on the availability of cardiovascular disease diagnostic and/or treatment services.
- **Chronic respiratory diseases:** Section 8.4, including **Tables 8.5–8.6** and **Figures 8.7–8.9** presents information on the availability of diagnostic and/or treatment services for chronic respiratory diseases.
- **Hypertension:** Section 8.5, including **Tables 8.7–8.8** and **Figures 8.10–8.12** presents information on the availability of hypertension diagnostic and/or treatment services.
- Cervical cancer: Section 8.6, including Table 8.9 and Figures 8.13–8.14 presents information on the availability of cervical cancer screening services.

8.2 AVAILABILITY OF SERVICES FOR DIABETES

8.2.1 Service Provision

Integrating diabetes diagnosis and treatment into relevant health services increases the opportunities for case detection and treatment follow-up. In health facilities in Bangladesh, clients who seek health care specifically for symptoms of diabetes are usually seen in a general outpatient department (OPD). However, there are also specific diabetic clinics or service areas in some health facilities.

Availability of Services for Diabetes

- Forty-five percent of public facilities in Bangladesh (44% excluding CCs) offer diabetes services. Ninety-eight percent of DHs, 99% of UHCs, 78% of NGOs, and 70% private facilities offer diabetes services (**Table 8.1** and **Figure 8.1**).
- Ninety-five percent of DHs and 91% of UHCs offer both diagnosis and treatment services for diabetes. However, only 63% of private facilities offer these services (Table 8.1).
- Forty five percent of CCs offer any services for diabetes (**Table 8.1** and **Figure 8.1**).
- Seventy-two percent of urban facilities offer services for diabetes, compared to 44% of rural facilities (**Table 8.1**).
- The availability of diabetes services ranges from a high of 55% of facilities in Chattogram division to a low of 35% of facilities in Barishal (**Table 8.1**).



Figure 8.1 Availability of diabetes services in health facilities, by facility type

Service Guidelines

 Among the public facilities that offered diabetes services, only 8% (6% excluding CCs) had guidelines for the diagnosis and management of diabetes on the day of survey. This was a decrease from 16% in 2017 (Table 8.1 and Figure 8.2).

Trained Staff

- Twenty-four percent of public facilities (26% excluding CCs) offering diabetes services have at least one staff member who received in-service training in diabetes services during the 24 months before the survey (**Table 8.1** and **Figure 8.2**), an increase from 13% in 2017.
- Forty-three percent of DHs, 79% of UHCs, and 23% of CCs (which mainly provide diabetes screening services), have staff members recently trained in diabetes services (**Table 8.1**).
- One-fifth of urban facilities (20%) and rural facilities (22%) have staff trained in the last 24 months (**Table 8.1**).
- Facilities in Sylhet are most likely to have staff with recent training in diabetes services (36%), while facilities in Rajshahi are least likely to have recently trained staff (13%) (**Table 8.1**).
- The percentage of all public facilities with staff who had ever received any relevant in-service training (54%) is higher than the proportion of facilities with staff who had received recent training (24%) (Table 8.1).

Equipment

 Availability of equipment for diabetes services has slightly declined since 2017 among public facilities offering diabetes services—e.g., blood pressure (BP) apparatus availability declined from 87% to 85%, availability of an adult weighing scale declined from 89% to 86%, and access to a height board declined from 79% to 70% (**Table 8.1** and **Figure 8.2**).





(N=717)

8.2.2 Availability of Diagnostic Capacity and Medicines for Diabetes

(N=212)

Diagnostic Capacity

Blood glucose diagnostic capacity among the public facilities offering diabetes diagnosis and treatment services has decreased since 2017, from 26% in 2017 to 21% in 2022. However, an incremental increase was observed in the capacity to test for urine protein (from 20% in 2017 to 30% in 2022) and urine glucose (from 18% in 2017 to 30% in 2022) (Table 8.2 and Figure 8.3).

(N=600)

- Private hospitals and NGO facilities are more likely than district and upazila public facilities to have the capacity to conduct all three tests (blood glucose, urine protein, and urine glucose) (**Table 8.2**).
- Urban facilities are much more likely than rural facilities to offer these diagnostic tests. Division-level results have shown that, overall, Dhaka and Chattagram division has the highest proportion of facilities providing the three diagnostic services (**Table 8.2**).

Medicines

- Since 2017, the availability of medicines among the public facilities offering diabetes diagnosis and treatment services has increased. Specifically, metformin availability rose from 19% to 37%, glibenclamide rose from 3% to 5%, injectable insulin rose from 17% to 37%, and availability of injectable glucose solution rose from 6% to 8% on the day of the survey (Table 8.2 and Figure 8.3).
- Medicines for diabetes require a prescription from a physician, and thus MCWCs, UHFWCs, and CCs
 generally have very limited capacity to offer these medicines. Among other facility types, DHs, UHCs,

and private hospitals are more likely to have medicines for the management of diabetes than USC/RDs and NGO facilities (**Table 8.2**).

• Urban facilities are more likely to be equipped with essential medicines than rural facilities (Table 8.2).

Figure 8.3 Diagnostic capacity and medicines to provide diabetes services in public facilities

Percent



8.3 AVAILABILITY OF SERVICES FOR CARDIOVASCULAR DISEASE

8.3.1 Service Provision

Table 8.3 and **Figures 8.4** & **8.5** provide information on the availability of services related to cardiovascular disease. **Table 8.3** and **Figure 8.5** also present information on the availability of service guidelines, trained staff, and equipment that support the provision of quality cardiovascular disease services.

Availability of Services for Cardiovascular Disease

- Twenty-five percent (33% excluding CCs) of public facilities in Bangladesh offer cardiovascular disease services, a decrease from 40% in 2017 (**Table 8.3** and **Figure 8.4**).
- Cardiovascular disease services are universally available in DHs (98%) and UHCs (97%). While the
 majority of NGOs (59%) and private hospitals (67%) offer these services, cardiovascular disease
 services are available in only 23% of CCs in 2022.
- Ninety percent of DHs and 87% of UHCs offer both diagnostic and treatment services for cardiovascular disease in 2022. However, greater than half of the private facilities (57%) offer both diagnostic and treatment services in 2022 (Table 8.3).

- Sixty-eight percent of urban facilities provide any services related to cardiovascular disease, compared to 24% of rural facilities (**Table 8.3**).
- Facilities in Dhaka (39%) are most likely to offer services related to cardiovascular disease, whereas facilities in Mymensingh (24%) are least likely to offer such services (**Table 8.3**).



Figure 8.4 Availability of cardiovascular disease services in health facilities, by facility type

Service Guidelines

• Among public facilities offering cardiovascular disease services, 12% have guidelines available for the diagnosis and management of cardiovascular diseases (**Table 8.3** and **Figure 8.5**).

Trained Staff

- Twenty-two percent (31% excluding CCs) of public facilities offering cardiovascular disease services have at least one staff member who received in-service training in cardiovascular disease-related services during the 24 months before the survey (Table 8.3). In 2017, this proportion was 14% (Figure 8.5).
- Forty-three percent of DHs, 79% of UHCs, and 17% of CCs (which mainly provide cardiovascular disease screening services) had staff with recent training on the day of survey (**Table 8.3**).
- Less than one-fifth of urban (16%) and rural (19%) facilities that offer CVD services have recently trained staff (**Table 8.3**).
- Facilities in Barishal are most likely to have staff with recent in-service training (31%), while facilities in Mymensingh and Rajshahi (both 8%) are least likely to have staff with recent training (**Table 8.3**).
- The proportion of facilities with staff who had ever received any relevant in-service training (44%) is higher than that of staff who had received recent training (22%) (**Table 8.3**).

Equipment

Among public facilities that offer CVD services, 93% had a stethoscope and 89% had a blood pressure apparatus available on the day of survey, a slight decrease from 2017. The availability of adult weighing scales among facilities offering CVD services has remained unchanged (85% in 2017 to 86% in 2022) (Table 8.3 and Figure 8.5).

Figure 8.5 Availability of trained staff, guidelines, and basic equipment to provide cardiovascular disease services in public facilities



8.3.2 Availability of Medicines and Commodities Related to Cardiovascular Disease

The 2022 BHFS collected data on the availability of essential medicines and commodities to support treatment of patients with cardiovascular disease. **Table 8.4** and **Figure 8.6** show that aspirin, beta blockers, and calcium channel blockers (amlodipine/nifedipine) are the most available medicines in public facilities.

- Among public facilities offering cardiovascular disease-related services, 43% (50% excluding CCs) had beta blockers available on the day of the survey visit and 40% (47% excluding CCs) had calcium channel blockers. By comparison, 46% (53% excluding CCs) of public facilities had aspirin, 14% (16% excluding CCs) had thiazide diuretics, and only 1% (the same percentage excluding CCs) had angiotensin-converting enzyme (ACE) inhibitors (enalapril) available.
- Thirty-five percent (40% excluding CCs) of public facilities had oxygen, in either cylinders or concentrators, or an oxygen distribution system on the day of survey.
- With the exception of ACE inhibitors, district and upazila level public facilities were more likely than other facility types to have all essential medicines available. Among public facilities, UHCs were most likely to have the essential medicines and commodities. Because a prescription is required from a physician, CCs are not able to offer these necessary medications; however, some union-level facilities had essential medicines and commodities available on the day of survey (**Table 8.4**).





8.4 AVAILABILITY OF SERVICES FOR CHRONIC RESPIRATORY DISEASES

8.4.1 Service Provision

Percent

Table 8.5 and **Figures 8.7** & **8.8** provide information from the 2022 BHFS on the availability of services for chronic respiratory diseases. **Figures 8.7** & **8.8** also present information on the availability of service guidelines, trained staff, and equipment that support the provision of quality services related to respiratory diseases.

Availability of Services for Chronic Respiratory Diseases

- Fifty-one percent (52% excluding CCs) of all public facilities can diagnose, prescribe treatment, or manage patients with chronic respiratory diseases (**Table 8.5** and **Figure 8.7**).
- DHs (98%), UHCs (98%), USC/RDs (70%), and private hospitals (66%) are more likely to provide services for chronic respiratory diseases than other facilities.
- Ninety-two percent of DHs and 91% of UHCs offer both diagnosis and treatment services for chronic respiratory diseases in 2022. By contrast, only 60% of private facilities offer both of these services (Table 8.5).
- Sixty-eight percent of urban facilities provide any services related to chronic respiratory diseases, compared to 50% of rural facilities (**Table 8.5**).
- Facilities in Rangpur (63%) are most likely to offer services related to chronic respiratory diseases, while facilities in Barishal (35%) are least likely to offer such services (**Table 8.5**).



Figure 8.7 Availability of chronic respiratory disease services in health facilities, by facility type

Service Guidelines

- Fifteen percent (8% excluding CCs) of public facilities offering services for chronic respiratory diseases had guidelines for the diagnosis and management of these diseases available at the service site on the day of the survey (**Table 8.5** and **Figure 8.8**).
- UHCs (26%) and CCs (17%) are more likely to have guidelines available at the service site than other facility types (**Table 8.5**).

Trained Staff

- Among public facilities that offer services related to chronic respiratory diseases, 19% (20% excluding CCs) have at least one staff member who had received in-service training in chronic respiratory diseases during the 24 months before the survey in 2022 (Table 8.5 and Figure 8.8).
- UHCs (67%) and DHs (36%) are more likely than CCs (19%), NGO facilities (17%), and private hospitals (11%) to have at least one staff member with recent training (**Table 8.5**).
- Facilities in Sylhet (33%) are most likely to have staff with recent in-service training, while facilities in Rajshahi (12%) are least likely to have staff with recent training in chronic respiratory diseases (Table 8.5).
- The proportion of public facilities with staff who had ever received any relevant in-service training (48%) is higher than that of staff who had received recent training (19%).

Equipment

- Overall, 90% (98% excluding CCs) of public facilities offering services related to respiratory diseases have a stethoscope and 1% (4% excluding CCs) have spacers for inhalers available at the relevant service sites (**Table 8.5** and **Figure 8.8**).
- In 2022, 11% of NGOs and 24% of private facilities have functioning peak flow meters. Ten percent of NGOs and 25% of private facilities have spacers for inhalers available in 2022. The percentages for both of these equipment's availability in DHs and UHCs are relatively low (Table 8.5).

Figure 8.8 Availability of trained staff, guidelines, and basic equipment to provide chronic respiratory disease-related services among public facilities



8.4.2 Availability of Medicines and Commodities for Chronic Respiratory Diseases

Table 8.6 and **Figure 8.9** show that salbutamol inhalers or tablets are the most widely available (92% in public facilities) of the medicines considered essential for the treatment of chronic respiratory diseases.

- Among public facilities that offer both diagnosis and treatment services for chronic respiratory diseases, 1% (2% excluding CCs) had beclomethasone inhalers, 3% (7% excluding CCs) had prednisolone tablets, 8% (25% excluding CCs) provided hydrocortisone injections, and 1% (3% excluding CCs) had injectable epinephrine available on the day of survey (Table 8.6 and Figure 8.9).
- Eleven percent (30% excluding CCs) of public facilities that offer both diagnosis and treatment services for chronic respiratory diseases have oxygen in either cylinders or concentrators or an oxygen distribution system (Table 8.6 and Figure 8.9).
- Overall, private hospitals are more likely to have the essential medicines and commodities than other types of facilities that offer both diagnosis and treatment for chronic respiratory diseases. Among public facilities, DHs are generally most likely to have these medicines and commodities (**Table 8.6**).

Figure 8.9 Availability of commodities and medicines to provide chronic respiratory disease services among public facilities



8.5 AVAILABILITY OF SERVICES FOR HYPERTENSION

8.5.1 Service Provision

Table 8.7 and **Figures 8.10** & **8.11** provide information from the 2022 BHFS on the availability of services for hypertension. **Figures 8.10** & **8.11** also present information on the availability of service guidelines, trained staff, and equipment that support the provision of quality services for hypertension.

Availability of Services for Hypertension

- Fifty-four percent (55% excluding CCs) of public health facilities can diagnose, prescribe treatment, or manage patients with hypertension (**Table 8.7** and **Figure 8.10**).
- DHs (97%), UHCs (99%), NGO clinics (75%), and private hospitals (70%) are more likely to provide services for hypertension than other facilities (Table 8.7 and Figure 8.10).
- Ninety-two percent of DHs and UHCs offer both diagnosis and treatment services for hypertension in 2022. However, only 65% of private hospitals offer both of these services together (Table 8.7).
- Seventy-two percent of urban facilities provide services for hypertension, compared to 53% of rural facilities (**Table 8.7**).
- Mymensingh (64%) has the highest percentage of facilities providing hypertension services, while Barishal (39%) has the lowest proportion (Table 8.7).



Figure 8.10 Availability of hypertension services in health facilities, by facility type

Service Guidelines

- Seventeen percent (11% excluding CCs) of public facilities offering services for hypertension had guidelines for the diagnosis and management of hypertension available at the service site on the day of the survey (**Table 8.7** and **Figure 8.11**).
- UHCs (39%), CCs (20%), NGO facilities (12%), and DHs (13%), were most likely to have guidelines available on the day of survey.

Trained Staff

- Among public facilities that offer services for hypertension, 20% (22% excluding CCs) have at least one staff member who had received in-service training in hypertension during the 24 months before the survey (**Table 8.7** and **Figure 8.11**).
- UHCs (79%), DHs (43%), and CCs (20%) are more likely than other facilities to have at least one staff member with recent training (**Table 8.7**).
- Facilities in Sylhet (37%) are most likely and those in Rajshahi (9%) are least likely to have staff with recent training compared to facilities in other divisions (**Table 8.7**).
- The percentage of public facilities with staff who had ever received any relevant in-service training (45%) is higher than that of staff who received recent training (20%).

Equipment

• Overall, 91% (98% excluding CCs) of public facilities that offer services for hypertension have a stethoscope available at the relevant service sites (**Table 8.7** and **Figure 8.11**).

• The proportion of public facilities having blood pressure apparatus is high across all facility types (**Table 8.7**).





8.5.2 Availability of Medicines and Commodities for Hypertension

- Among facilities offering services for hypertension, 34% of public facilities (46% excluding CCs) had angiotensin receptor blockers (ARBs) and 28% (38% excluding CCs) had calcium channel blockers available on the day of the survey visit (**Table 8.8** and **Figure 8.12**). However, only 8% (10% excluding CCs) had thiazide available on the day of the survey visit.
- ARBs and calcium channel blockers are more available in DHs (88% and 70%, respectively), UHCs (91% and 71%, respectively), and private hospitals (48% and 54%, respectively) (**Table 8.8**).
- Urban facilities are more likely to have ARBs and calcium channel blockers (49% and 54%, respectively) than their rural counterparts (27% and 21%, respectively) (**Table 8.8**).

Figure 8.12 Availability of medicines to provide services for hypertension among public facilities

Percent



8.6 AVAILABILITY OF SERVICES FOR CERVICAL CANCER

Table 8.9 and **Figures 8.13 & 8.14** provide information from the 2022 BHFS on the availability of services for cervical cancer. **Figures 8.13 & 8.14** also present information on the availability of service guidelines, trained staff, and equipment that support the provision of quality services for cervical cancer.

8.6.1 Availability of Screening Services for Cervical Cancer

- Four percent (10% excluding CCs) of public facilities provide diagnosis/screening services for cervical cancer (**Table 8.9** and **Figure 8.13**).
- DHs (86%), UHCs (89%), MCWCs (30%), and NGO clinics (21%) are more likely to provide diagnosis/screening services for cervical cancer than other facilities (**Table 8.9**).
- Twenty-one percent of NGOs offer diagnosis/screening services for cervical cancer. However, only 10% of private hospitals offer these services.
- Eighteen percent of urban facilities provide diagnosis/screening services for cervical cancer, compared to only 3% of rural facilities (**Table 8.9**).
- Overall, the availability of cervical cancer diagnosis/screening services is lower than 10% across all divisions (**Table 8.9**).

Figure 8.13 Availability of cervical cancer screening/diagnosis services in health facilities, by facility type



8.6.2 Service Guidelines

- Twenty-six percent (39% excluding CCs) of public facilities that offer diagnosis/screening services for cervical cancer had guidelines for screening available at the service site on the day of the survey (Table 8.9 and Figure 8.14).
- Among facilities offering diagnosis/screening services for cervical cancer, DHs (47%), UHCs (41%), and NGOs (30%) were most likely to have guidelines for screening cervical cancer available on the day of survey (**Table 8.9**).

8.6.3 Trained Staff

- Among public facilities offering diagnosis/screening services for cervical cancer, 49% (46% excluding CCs) have at least one staff member who had received any in-service training related to observation of acetic acid, pap smear, or Human Papilloma Virus (HPV) test for screening of cervical cancer during the 24 months before the survey (Table 8.9 and Figure 8.14).
- Among facilities offering diagnosis/screening services for cervical cancer, UHCs (52%), DHs (45%), and CCs (56%) are more likely than other facilities to have at least one staff member with recent training available on the day of survey (**Table 8.9**).
- Facilities offering diagnosis/screening services in Mymensingh (60%) and Rangpur (51%) are more likely than facilities in other divisions to have staff with recent cervical cancer training (**Table 8.9**).

8.6.4 Equipment

• Overall, 67% (97% excluding CCs) of public facilities that offer screening services for cervical cancer have a speculum available at the relevant service site (**Table 8.9** and **Figure 8.14.**).
- Speculum availability is nearly universal in DHs, MCWCs, UHCs, union-level facilities, and private hospitals offering diagnosis/screening services for cervical cancer (**Table 8.9**).
- Urban facilities offering diagnosis/screening services for cervical cancer are nearly two times more likely to have a speculum than rural facilities (94% and 50%) (Table 8.9).
- All facilities offering diagnosis/screening services for cervical cancer in Barishal have a speculum, and availability is high in most other divisions with the exception of Mymensingh (**Table 8.9**).

Figure 8.14 Availability of trained staff, guidelines, and basic equipment to diagnose/screen cervical cancer in public facilities

Percent



| 2017 BHFS | ■2022 Bł |
|-----------|----------|
| (N=24) | (N=55) |

NON-COMMUNICABLE DISEASES

LIST OF TABLES

- Table 8.1 Availability of diabetes services and guidelines, trained staff, and equipment
- Table 8.2 Diagnostic capacity and essential medicines for diabetes
- Table 8.3 Availability of cardiovascular disease services and guidelines, trained staff, and equipment
- Table 8.4 Availability of essential medicines and commodities for cardiovascular disease
- Table 8.5 Availability of chronic respiratory disease services and guidelines, trained staff, and equipment
- **Table 8.6** Availability of essential medicines and commodities for chronic respiratory diseases
- Table 8.7 Availability of hypertension services and guidelines, trained staff, and equipment
- Table 8.8 Availability of essential medicines and commodities for hypertension
- Table 8.9 Availability of cervical cancer diagnosis/screening services and guidelines, trained staff, and equipment

Table 8.1: Availability of diabetes services and guidelines, trained staff, and equipment

Among all facilities, the percentages that offer services for diabetes, and among facilities that offer services for diabetes, the percentages with guidelines, at least one staff member recently trained on diabetes, at least one staff member ever trained on diabetes, and the indicated equipment observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | | Percentage of for dia | facilities offe abetes that h | ering services nave: | | Equipment | Number of | Number of | | | | |
|--|---|--|---|--|---|--|--|--|--|--|--|--|
| Background characteristic | offering both diagnosis and treatment services for diabetes | Percentage of facilities offering any services for diabetes ¹ | Number of facilities (weighted) | Number of facilities (unweighted) | Guidelines for the diagnosis and management of diabetes | Trained staff (last 24 months) ² | Trained staff (ever) ³ | Blood pressure apparatus ⁴ | Adult weighing scale | Height board or stadiometer | facilities offering services for diabetes (weighted) | facilities offering services for diabetes (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 79.1 95.2 22.7 90.5 | 89.3 98.4 44.9 98.8 | 41 4 7 30 | 319 62 100 157 | 32.2 16.4 6.6 37.4 | 68.3 42.6 4.4 79.2 | 84.2 88.5 34.6 89.0 | 95.9 91.8 97.8 96.3 | 80.7 75.4 88.8 80.6 | 64.9 52.5 66.4 66.6 | 36 4 3 29 | 261 61 45 155 |
| Union-level public facilities UHFWC USC/RD | 15.0 9.5 29.7 | 37.9 29.7 59.8 | 310 225 85 | 434 293 141 | 6.9 7.0 6.8 | 13.0 1.3 28.4 | 29.5 13.4 50.6 | 97.0 99.2 94.2 | 79.1 90.8 63.6 | 53.8 59.8 45.8 | 118 67 51 | 177 85 92 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 2.2 61.7 62.9 | 44.9 78.4 70.3 | 994 29 183 | 488 127 189 | 20.5 14.6 9.4 | 23.1 12.9 12.9 | 57.5 26.8 32.9 | 81.3 90.8 98.3 | 88.2 98.0 94.6 | 74.1 67.7 31.7 | 446 23 129 | 217 103 139 |
| Location Urban Rural | 62.6 6.6 | 72.4 44.0 | 235 1,322 | 514 1,043 | 13.2 18.0 | 19.5 22.3 | 40.4 51.9 | 97.1 84.9 | 92.9 86.3 | 40.1 69.7 | 170 582 | 402 495 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 13.2 20.5 17.3 12.4 13.3 9.9 16.2 13.2 | 35.1 55.2 51.8 48.6 50.7 39.0 40.7 51.2 | 111 291 317 210 214 193 99 122 | 152 244 264 208 212 190 136 151 | 11.3 19.8 16.7 16.8 12.7 7.0 20.8 30.6 | 33.1 26.8 17.6 20.1 13.3 18.8 35.9 23.6 | 54.5 55.7 40.0 50.4 43.7 48.9 53.4 59.6 | 72.9 86.5 89.0 88.1 88.5 85.5 99.7 89.0 | 80.7 87.1 88.5 88.2 89.5 91.3 83.0 87.4 | 60.4 66.6 47.0 64.4 60.9 75.0 77.7 75.1 | 39 161 164 102 109 75 40 62 | 79 155 153 120 123 102 76 89 |
| Total | 15.1 | 48.3 | 1,557 | 1,557 | 16.9 | 21.7 | 49.3 | 87.7 | 87.8 | 63.0 | 752 | 897 |
| Total excluding CCs | 37.7 | 54.3 | 563 | 1,069 | 11.6 | 19.6 | 37.3 | 97.0 | 87.2 | 46.9 | 306 | 680 |
| Total public | 7.6 | 44.7 | 1.345 | 1.241 | 8.3 | 23.9 | 53.6 | 85.2 | 86.0 | 69.6 | 600 | 655 |
| Total public excluding CCs | 22.6 | 44.0 | 351 | 753 | 5.7 | 26.2 | 42.5 | 96.8 | 79.4 | 56.4 | 154 | 438 |

Note: The indicators presented in this table comprise the staff and training and equipment domains for assessing readiness to provide services for diabetes within the health facility assessment methodology proposed by the WHO and USAID (2012).

¹ Providers in the facility diagnose, prescribe treatment for, or manage patients with diabetes.

² At least one interviewed provider of diabetes services reported receiving in-service training in diabetes services during the 24 months preceding the survey. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ At least one interviewed provider of diabetes services reported ever receiving in-service training in diabetes services. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

⁴ Functioning digital blood pressure machine or manual sphygmomanometer with stethoscope.

Table 8.2: Diagnostic capacity and essential medicines for diabetes

Among facilities offering both diagnosis and treatment services for diabetes, the percentages having the indicated diagnostic capacity and essential medicines observed at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | | | | | | | | Number of facilities offering both | Number of facilities offering both |
|---|-------------------------------------|-------------------------------------|-------------------------------------|------------------------------------|------------------------------------|------------------------------------|------------------------------------|---|---|
| | Dia | agnostic capa | city | | Med | licines | | and | and |
| Background characteristic | Blood glucose ¹ | Urine protein ² | Urine glucose ³ | Metformin | Gliben- clamide | Injectable insulin | Injectable glucose solution | treatment services for diabetes (weighted) | treatment services for diabetes (unweighted) |
| Facility type | | | | | | | | | |
| District and upazila public facilities DH MCWC UHC | 37.6 20.3 38.8 40.2 | 66.7 78.0 43.1 66.4 | 67.2 78.0 43.1 67.0 | 83.1 81.4 4.1 88.1 | 14.3 11.9 0.0 15.6 | 84.6 83.1 0.0 90.0 | 23.8 30.5 0.0 24.2 | 33 4 2 27 | 225 59 23 143 |
| Union-level public facilities UHFWC USC/RD | 19.4 28.1 12.1 | 19.5 34.8 6.5 | 18.0 34.8 3.7 | 23.4 4.1 39.8 | 1.5 0.0 2.8 | 21.5 0.0 39.8 | 1.5 0.0 2.8 | 47 22 25 | 72 26 46 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 0.0 77.7 53.6 | 0.0 79.1 83.9 | 0.0 76.6 87.0 | 0.0 13.2 46.1 | 0.0 4.7 11.6 | 0.0 12.1 52.3 | 0.0 4.8 27.7 | 22 18 115 | 12 80 126 |
| Location | 50 5 | 00 F | | 40.0 | 44.0 | 50.0 | 04.0 | 140 | 0.40 |
| Rural | 52.5 22.5 | 80.5 26.4 | 83.0 25.2 | 46.2 29.1 | 2.6 | 52.2 26.5 | 24.8 5.5 | 87 | 343 172 |
| Division | | | | | | | | | |
| Barishal Chattogram Dhaka Khulna | 30.3 59.4 53.5 27.8 | 45.2 68.9 72.4 46.4 | 44.6 70.9 72.4 46.4 | 36.8 47.2 50.3 27.7 | 5.1 7.7 10.2 11.1 | 35.2 51.9 55.7 27.3 | 13.6 27.6 22.3 12.8 | 15 60 55 26 | 55 96 82 62 |
| Rajshahi Rangpur Sylhet Mymonsingh | 32.9 28.7 14.8 21.2 | 65.5 49.8 40.7 47.0 | 68.0 49.1 43.8 | 30.0 27.4 40.2 21.2 | 12.0 4.1 8.5 | 31.9 29.2 38.5 | 12.6 10.5 5.0 | 28 19 16 | 63 63 43 |
| | 21.5 | 47.5 | 43.5 | 51.2 | 1.5 | 34.0 | 0.1 | 10 | 51 |
| Total | 41.3 | 60.4 | 61.5 | 39.9 | 8.4 | 42.6 | 17.6 | 235 | 515 |
| Total excluding CCS | 45.7 21.0 | 00.1 30.1 | 67.9 20.0 | 44.1 37 <i>1</i> | 9.Z | 47.1 37.1 | 19.4 8.4 | 213 | 300 |
| Total public excluding CCs | 26.9 | 39.0 | 38.3 | 48.0 | 6.8 | 47.5 | 10.7 | 80 | 297 |

Note: The indicators presented in this table comprise the diagnostics, medicines, and commodities domains for assessing readiness to provide services for diabetes within the health facility assessment methodology proposed by the WHO and USAID (2012). ¹ Facility had a functioning glucometer and unexpired glucose test strips in the facility on the day of the survey. ² Facility had unexpired urine dipsticks for testing for urine protein available in the facility on the day of the survey. ³ Facility had unexpired urine dipsticks for testing for urine glucose available in the facility on the day of the survey.

Table 8.3: Availability of cardiovascular disease services and guidelines, trained staff, and equipment

Among all facilities, the percentages offering services for cardiovascular diseases, and among facilities offering services for cardiovascular diseases, the percentages having guidelines, at least one staff member recently trained on cardiovascular disease, and the indicated equipment observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Dereentage of | | | | Percentage of fa cardiovascul | cilities offering lar diseases tha | services for at have: | | Equipment | Number of facilities | Number of facilities | |
|--|---|---|---------------------------------------|---|--|---|--------------------------------------|--------------|---------------------------------|----------------------------|---|--|
| Background characteristic | facilities offering both diagnosis and treatment services for cardiovascular diseases | Percentage of facilities offering any services for cardiovascular diseases ¹ | Number of facilities (weighted) | Number of facilities (unweighted) | Guidelines for diagnosis and management of cardiovascular diseases | Trained staff (last 24 months) ² | Trained staff (ever) ³ | Stethoscope | Blood pressure apparatus⁴ | Adult weighing scale | services for cardio- vascular diseases (weighted) | offering services for cardiovascular diseases (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public facilities | 75.0 | 85.1 | 41 | 319 | 25.1 | 69.9 | 84.3 | 94.9 | 95.7 | 81.0 75 4 | 35 | 243 |
| | 90.3 | 90.4 30.1 | 4 | 02 100 | 6.4 | 42.0 | 86.9 26.4 | 96.7 | 91.8 | 75.4 | 4 | 30 |
| | 86.6 | 96.6 | 30 | 157 | 28.4 | 70.1 | 20.4 | 90.7 | 90.7 | 93.3 80 Q | 20 | 152 |
| | 00.0 | 30.0 | 50 | 157 | 20.4 | 75.1 | 00.5 | 34.4 | 30.2 | 00.9 | 23 | 152 |
| public facilities | 10.9 | 26.1 | 310 | 434 | 6.8 | 14.4 | 29.8 | 98.5 | 96.7 | 81.4 | 81 | 123 |
| UHFWC USC/RD | 5.2 25.9 | 19.7 43.2 | 225 85 | 293 141 | 6.1 7.7 | 4.5 26.3 | 16.0 46.3 | 98.8 98.2 | 98.8 94.2 | 94.5 65.6 | 44 37 | 57 66 |
| Community clinic (CC) | 1.0 | 22.7 | 994 | 488 | 11.6 | 16.8 | 43.2 | 90.1 | 85.2 | 87.9 | 226 | 109 |
| NGO clinic/hospital | 43.6 | 59.2 | 29 | 127 | 10.8 | 9.6 | 17.3 | 96.1 | 89.6 | 97.3 | 17 | 76 |
| Private hospital | 57.3 | 67.3 | 183 | 189 | 4.9 | 8.5 | 24.5 | 98.5 | 98.3 | 93.2 | 123 | 134 |
| Location | 50.0 | 07.0 | 005 | 544 | | 10.0 | 00.0 | | | 04.7 | 100 | 005 |
| Urban | 56.6 | 67.8 | 235 | 514 | 8.4 | 16.2 | 32.0 | 98.0 | 96.9 | 91.7 | 160 | 365 |
| Rural | 4.5 | 24.4 | 1,322 | 1,043 | 10.8 | 18.8 | 41.4 | 92.4 | 88.7 | 86.2 | 322 | 320 |
| Division Barishal | 11.7 | 30.2 | 111 | 152 | 11.8 | 31.0 | 45.7 | 88.4 | 78.7 | 88.7 | 34 | 74 |
| Chattogram | 16.5 | 34.5 | 291 | 244 | 9.4 | 25.5 | 44.7 | 97.3 | 92.6 | 88.8 | 100 | 115 |
| Dhaka | 13.8 | 39.1 | 318 | 264 | 11.7 | 13.4 | 25.5 | 93.9 | 91.1 | 91.9 | 124 | 126 |
| Knuina Raichabi | 11.1 | 24.9 | 210 | 208 | 13.9 | 20.3 | 40.3 | 96.3 | 95.4 | 88.3 | 52 | (/ |
| Rangour | 8.3 | 29.3 | 193 | 190 | 0.1 | 16.8 | 52.0 | 92.0 | 90.2 | 87.1 | 53 | 79 |
| Svlhet | 9.6 | 26.8 | 99 | 136 | 10.7 | 24.5 | 35.9 | 92.5 | 99.5 | 74.1 | 27 | 58 |
| Mymensingh | 9.0 | 24.0 | 122 | 151 | 20.8 | 8.3 | 34.6 | 91.6 | 90.5 | 82.5 | 29 | 63 |
| Total | 12.3 | 31.0 | 1.557 | 1,557 | 10.0 | 17.9 | 38.3 | 94.2 | 91.4 | 88.0 | 482 | 685 |
| Total excluding CCs | 32.3 | 45.5 | 563 | 1.069 | 8.7 | 18.9 | 33.9 | 97.9 | 96.8 | 88.1 | 256 | 576 |
| Total public | 5.5 | 25.4 | 1,345 | 1,241 | 11.8 | 21.7 | 44.3 | 92.6 | 89.0 | 85.7. | 342 | 475 |
| Total public excluding CCs | 18.4 | 33.2 | 351 | 753 | 12.4 | 31.2 | 46.3 | 97.4 | 96.4 | 81.3 | 116 | 366 |

Note: The indicators presented in this table comprise the staff and training and equipment domains for assessing readiness to provide services for cardiovascular diseases within the health facility assessment methodology proposed by the WHO and USAID (2012).

¹ Providers in the facility diagnose, prescribe treatment for, or manage patients with cardiovascular diseases.

² At least one interviewed provider of cardiovascular disease services reported receiving in-service training in cardiovascular diseases during the 24 months preceding the survey. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ At least one interviewed provider of cardiovascular disease services reported ever receiving in-service training in cardiovascular diseases. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

⁴ Functioning digital blood pressure machine or manual sphygmomanometer with stethoscope.

Table 8.4: Availability of essential medicines and commodities for cardiovascular disease

Among facilities offering both diagnosis and treatment services for cardiovascular disease, the percentages having the indicated essential medicines and commodities observed at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percenta | Number of facilities offering both | Number of facilities offering both | | | | | |
|------------------------------|-------------------------------|--|--|--|------|------|---|---|
| Background characteristic | ACE inhibitors (enalapril) | Thiazide | Beta blockers (atenolol) | Calcium channel blockers a blockers (amlodipine/ atenolol) nifedipine) Aspirin | | | diagnosis and treatment services for cardiovascular disease (weighted) | diagnosis and treatment services for cardiovascular disease (unweighted) |
| Facility type | | | | | | | | |
| District and upazila | | | | | | | | |
| public facilities | 2.1 | 26.8 | 69.5 | 67.9 | 81.4 | 54.1 | 31 | 211 |
| DH | 3.6 | 10.7 | 35.7 | 66.1 | 67.9 | 51.8 | 4 | 56 |
| MCWC | 0.0 | 0.0 | 5.2 | 0.0 | 0.0 | 60.6 | 1 | 18 |
| UHC | 2.0 | 30.7 | 78.0 | 71.6 | 87.5 | 54.2 | 26 | 137 |
| Union-level | | | | | | | | |
| public facilities | 0.0 | 5.4 | 32.1 | 27.0 | 27.2 | 27.4 | 34 | 56 |
| UHFWC | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.5 | 12 | 15 |
| USC/RD | 0.0 | 8.3 | 49.2 | 41.3 | 41.7 | 26.7 | 22 | 41 |
| Community clinic | | | | | | | | |
| (CC) | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10 | 5 |
| NGO clinic/hospital | 0.8 | 4.4 | 16.6 | 37.5 | 11.9 | 35.6 | 12 | 56 |
| Private hospital | 6.2 | 16.3 | 36.3 | 52.2 | 39.1 | 75.8 | 105 | 116 |
| Location | | | | | | | | |
| Urban | 5.0 | 15.7 | 38.8 | 53.4 | 41.9 | 69.1 | 133 | 309 |
| Rural | 0.9 | 11.7 | 35.6 | 31.5 | 35.7 | 30.6 | 59 | 135 |
| Division | | | | | | | | |
| Barishal | 5.2 | 17.8 | 37.8 | 46.7 | 31.2 | 56.4 | 13 | 54 |
| Chattogram | 0.1 | 11.7 | 39.5 | 53.0 | 45.0 | 67.0 | 48 | 76 |
| Dhaka | 7.4 | 23.9 | 49.4 | 61.9 | 45.8 | 68.2 | 44 | 71 |
| Khulna | 0.0 | 10.4 | 32.0 | 35.2 | 31.1 | 41.9 | 23 | 54 |
| Raishahi | 7.5 | 10.5 | 29.6 | 33.2 | 29.0 | 45.0 | 27 | 61 |
| Rangpur | 2.7 | 5.4 | 23.0 | 22.5 | 27.9 | 53.5 | 16 | 51 |
| Svlhet | 3.3 | 21.2 | 28.6 | 37.4 | 42.2 | 41.5 | 10 | 32 |
| Mymensingh | 3.9 | 11.4 | 46.1 | 59.1 | 67.7 | 54.4 | 11 | 45 |
| Total | 3.8 | 14.5 | 37.8 | 46.7 | 40.0 | 57.3 | 192 | 444 |
| Total excluding CCs | 4.0 | 15.3 | 39.8 | 49.2 | 42.2 | 60.4 | 182 | 439 |
| Total public | 0.9 | 13.6 | 43.4 | 40.4 | 46.2 | 34.9 | 75 | 472 |
| Total public excluding CCs | 1.0 | 15.7 | 50.1 | 46.6 | 53.2 | 40.2 | 65 | 267 |

Note: The indicators presented in this table comprise the medicines and commodities domain for assessing readiness to provide services for cardiovascular diseases within the health facility assessment methodology proposed by the WHO and USAID (2012). ¹ In cylinders or concentrators or an oxygen distribution system.

Table 8.5: Availability of chronic respiratory disease services and guidelines, trained staff, and equipment

Among all facilities, the percentages offering services for chronic respiratory diseases, and among facilities offering services for chronic respiratory diseases, the percentages having guidelines, at least one staff member recently trained on chronic respiratory diseases, and the indicated equipment observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of | | | | Percentage of for chronic r | facilities offerin espiratory disea have: | ig services ases that | | Equipment | | Number of | Number of |
|--|---|---|---------------------------------------|---|--|---|---|--------------------------|---------------------------------|-------------------------|--|--|
| Background characteristic | facilities offering both diagnosis and treatment services for chronic respiratory diseases | Percentage of facilities offering any services for chronic respiratory diseases ¹ | Number of facilities (weighted) | Number of facilities (unweighted) | Guidelines for diagnosis and management of chronic respiratory diseases | Trained staff (last 24 months) ² | Trained staff (ever) ³ | Stethoscope ⁴ | Peak flow meter ⁵ | Spacers for inhalers | offering services for chronic respiratory diseases (weighted) | offering services for chronic respiratory diseases (unweighted) |
| Facility type | | | | | | | | | | | | |
| District and upazila public facilities | 80.7 | 88.7 | 41 | 319 | 22.1 | 57.2 | 75.9 | 95.7 | 6.0 | 9.5 | 36 | 260 |
| DH | 91.9 | 98.4 | 4 | 62 | 9.8 | 36.1 | 78.7 | 96.7 | 4.9 | 9.8 | 4 | 61 |
| | 33.7 | 45.9 | 7 30 | 100 | 6.4 25.7 | 0.0 | 23.8 | 95.7 | 2.1 | 2.1 | 3 20 | 40 |
| | 30.5 | 51.1 | 50 | 157 | 20.7 | 00.5 | 01.4 | 33.0 | 0.0 | 10.4 | 23 | 100 |
| public facilities | 29.8 | 47.6 | 310 | 434 | 5.0 | 10.4 | 32.7 | 97.9 | 0.4 | 2.0 | 147 | 220 |
| UHFWC | 22.1 | 39.2 | 225 | 293 | 4.8 | 4.9 | 22.8 | 99.4 | 0.6 | 0.0 | 88 | 115 |
| USC/RD | 50.3 | 70.0 | 85 | 141 | 5.4 | 18.6 | 47.5 | 95.6 | 0.0 | 5.1 | 59 | 105 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 24.5 45.6 60.4 | 50.0 63.9 66.0 | 994 29 183 | 488 127 189 | 17.0 9.6 2.5 | 19.0 17.2 10.8 | 49.9 23.5 26.7 | 87.4 94.7 98.5 | 0.0 11.4 23.9 | 0.0 10.4 24.5 | 497 19 121 | 241 83 134 |
| Location | •••• | | | | | | | | -0.0 | | | |
| Urban Rural | 60.3 26.6 | 68.0 49.9 | 235 1,322 | 514 1,043 | 5.8 14.5 | 17.5 18.0 | 33.6 46.5 | 98.2 89.9 | 20.3 0.2 | 21.3 0.6 | 160 660 | 380 558 |
| Division | | | | | | | | | | | | |
| Barishal Chattagram | 18.8 | 34.8 | 112 | 152 | 14.3 | 26.3 | 49.5 | 90.3 | 2.0 | 1.1 | 39 157 | 80 150 |
| Dhaka | 24.8 | 52 1 | 318 | 244 | 7.0 | 14.1 | 32.8 | 93.2 | 7.0 | 73 | 165 | 156 |
| Khulna | 23.2 | 46.1 | 210 | 204 | 11.3 | 22.8 | 55.5 | 96.1 | 1.9 | 0.3 | 97 | 120 |
| Rajshahi | 41 | 57.6 | 214 | 212 | 14.3 | 11.7 | 45.7 | 90.7 | 3.2 | 2.9 | 123 | 132 |
| Rangpur | 43.8 | 62.9 | 193 | 190 | 13.4 | 15.7 | 42.6 | 89.3 | 1.7 | 1.4 | 122 | 126 |
| Sylhet | 31.8 | 44.3 | 99 | 136 | 12.0 | 33.1 | 53.2 | 85.6 | 5.0 | 5.4 | 44 | 71 |
| Mymensingh | 37.3 | 60.1 | 122 | 151 | 23.0 | 16.9 | 46.6 | 86.2 | 2.5 | 1.8 | 73 | 103 |
| Total | 31.7 | 52.7 | 1,557 | 1.557 | 12.8 | 17.9 | 43.9 | 91.5 | 4.1 | 4.6 | 820 | 938 |
| Total excluding CCs | 44.3 | 57.4 | 563 | 1,069 | 6.3 | 16.2 | 34.9 | 97.7 | 10.4 | 11.8 | 323 | 697 |
| Total public | 27.5 | 50.6 | 1,345 | 1,241 | 14.7 | 19.2 | 47.6 | 90.1 | 0.4 | 1.0 | 680 | 721 |
| Total public excluding | 35.8 | 52.4 | 351 | 753 | 8.4 | 19.7 | 41.3 | 97.5 | 1.5 | 3.5 | 183 | 480 |

¹ Providers in the facility diagnose, prescribe treatment for, or manage patients with respiratory disease.

² At least one interviewed provider of services for chronic respiratory diseases reported receiving in-service training in chronic respiratory diseases during the 24 months preceding the survey. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ At least one interviewed provider of services for chronic respiratory diseases reported ever receiving in-service training in chronic respiratory diseases. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

⁴ Functioning stethoscope available at the service site on the day of the survey.

⁵ Functioning peak flow meter available at the service site on the day of the survey.

Table 8.6: Availability of essential medicines and commodities for chronic respiratory diseases

Among facilities offering both diagnosis and treatment services for chronic respiratory diseases, the percentages having the indicated essential medicines and commodities observed at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percent | age of facilities o the in | Number of facilities offering both diagnosis and treatment services for | Number of facilities offering both diagnosis and treatment services for | | | | |
|---|-------------------------------------|-------------------------------|--|--|---------------------------|---------------------|--|--|
| Background characteristic | Salbutamol inhaler or tablets | Beclometh- asone inhaler | Prednisolone tablets | Hydro- cortisone injection | Injectable epinephrine | Oxygen ¹ | chronic respiratory diseases (weighted) | chronic respiratory diseases (unweighted) |
| Facility type | | | | | | | | |
| District and upazila public facilities DH | 89.6 71.9 | 7.7 10.5 | 19.3 29.8 | 78.3 73.7 | 9.6 7.0 | 53.2 49.1 | 33 4 | 233 57 |
| UHC | 87.8 92.5 | 0.0 8.0 | 0.0 19.4 | 14.9 84.8 | 2.9 10.5 | 49.9 54.1 | 2 27 | 34 142 |
| Union-level nublic | | | | | | | | |
| facilities | 81.6 | 0.0 | 19 | 53 | 0.0 | 21.0 | 92 | 142 |
| UHFWC | 88.1 | 0.0 | 2.8 | 4.0 | 0.0 | 27.2 | 50 | 68 |
| USC/RD | 74.0 | 0.0 | 0.9 | 6.8 | 0.0 | 13.8 | 42 | 74 |
| Community clinic | | | | | | | | |
| (CC) | 95.8 | 0.0 | 0.7 | 0.0 | 0.0 | 0.9 | 243 | 119 |
| NGO clinic/hospital | 21.1 | 2.8 | 7.2 | 29.6 | 3.4 | 34.1 | 13 | 63 |
| Private hospital | 42.7 | 18.8 | 40.1 | 55.3 | 13.5 | 75.5 | 112 | 125 |
| Location | 40.0 | 45.0 | 22.4 | FF 0 | 10.0 | 07.5 | 4.40 | 004 |
| Urban | 48.3 | 15.0 | 33.4 | 55.2 | 12.2 | 67.5 | 142 | 334 |
| Ruidi | 91.2 | 0.7 | 2.2 | 5.1 | 0.4 | 9.0 | 331 | 340 |
| Division | | | | | | | | |
| Barishal | 79.6 | 5.7 | 21.5 | 31.0 | 3.7 | 33.7 | 21 | 63 |
| Chattogram | 84.7 | 4.8 | 17.2 | 27.2 | 3.8 | 37.0 | 96 | 113 |
| Dhaka | 76.9 | 13.6 | 24.1 | 32.9 | 11.8 | 41.0 | 79 | 90 |
| Rhuina Raisbabi | 37.4 79.5 | 3.7 | 9.1 | 10.0 | 0.4 | 20.2 | 49 | 105 |
| Rajslidili Rangour | 70.3 | 1.7 | 4.7 | 0.3 | 2.5 | 19.3 | 07 85 | 105 |
| Sylhet | 84.9 | 5.8 | 12.5 | 9.1 | 33 | 16.8 | 31 | 56 |
| Mymensingh | 87.9 | 2.3 | 4.4 | 11.9 | 1.4 | 15.0 | 45 | 78 |
| Total | 78.8 | 4.8 | 11.2 | 19.5 | 3.8 | 25.8 | 493 | 682 |
| Total excluding CCs | 62.3 | 9.5 | 21.5 | 38.5 | 7.4 | 50.1 | 250 | 563 |
| Total public | 91.7 | 0.7 | 2.7 | 8.4 | 0.9 | 10.6 | 368 | 494 |
| Total public excluding CCs | 83.7 | 2.1 | 6.5 | 24.7 | 2.5 | 29.5 | 125 | 375 |

Note: The indicators presented in this table comprise the medicines and commodities domain for assessing readiness to provide services for cardiovascular diseases within the health facility assessment methodology proposed by the WHO and USAID (2012).

¹ In cylinders or concentrators or an oxygen distribution system.

Table 8.7: Availability of hypertension services and guidelines, trained staff, and equipment for hypertension services

Among all facilities, the percentages offering services for hypertension, and among facilities offering services for hypertension, the percentages having guidelines, at least one staff member recently trained on hypertension, at least one staff member ever trained on hypertension, and the indicated equipment observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| facilities offering both diagnosis and treatment characteristic Percentage of and treatment hypertension Number of facilities (augusted) Number of facilities (unweighted) Number of facilities (unweighted) Trained management of (last 24 Trained table Blood (last 24 Ac Facility type Facilities (unweighted) Image: State of table Stethoscope ⁴ Apparatus ⁵ District and upazila public facilities 81.3 90.7 41 319 32.8 67.1 81.9 95.0 96.4 8 | Number of facilities offering services for hypertension (weighted)Number of facilities offering services for hypertension (unweighted)11.03726925.046010.745410.629155 |
|---|--|
| Facility type District and upazila public facilities 81.3 90.7 41 319 32.8 67.1 81.9 95.0 96.4 8 DH 01.0 06.8 4 62 13.3 43.3 96.7 04.7 7 | 31.0 37 269 '5.0 4 60)0.7 4 54 10.6 29 155 |
| District and upazila public facilities 81.3 90.7 41 319 32.8 67.1 81.9 95.0 96.4 8 DH 04.0 06.8 4 62 13.3 43.3 86.7 06.7 04.7 7 | 31.0 37 269 75.0 4 60 10.7 4 54 10.6 29 155 |
| | 75.0 4 60 30.7 4 54 30.6 29 155 |
| יום פו.ש אינע אינע אינע אינע אינע אינע אינע אינע | 30.745430.629155 |
| MCWC 30.7 54.0 7 100 5.5 3.7 27.2 96.4 98.2 9 | 30.6 29 155 |
| UHC 92.0 98.8 30 157 39.3 78.9 88.5 94.6 96.8 8 | |
| Union-level public facilities 24.1 50.2 310 434 5.4 10.5 28.2 98.4 95.6 8 | 30.8 156 230 |
| UHFWC 11.5 42.6 225 293 4.4 3.8 17.3 99.5 97.2 9 | 91.7 96 124 |
| USC/RD 57.4 70.4 85 141 7.0 21.3 45.8 96.8 93.1 6 | 60 106 |
| Community clinic (CC) 3.8 53.4 994 488 19.6 19.5 47.1 88.7 85.4 8 NGO clinic (base) table 65.4 75.4 20 137 11.6 10.3 16.3 95.5 92.8 6 | 15.7 531 258 |
| Private hospital 64.6 69.5 183 189 3.1 8.2 24.4 98.6 99.8 9 |)3.4 127 139 |
| | |
| Urban 65.5 72.1 235 514 7.8 15.7 31.7 98.1 98.2 9 | 92.1 170 404 |
| Rural 9.8 53.2 1,322 1,043 16.8 18.6 43.5 90.9 88.0 8 | 34.6 703 590 |
| Division | |
| Barishal 13.5 38.6 111 152 13.6 24.8 44.6 82.9 70.6 8 | 30.3 43 84 |
| Chattogram 24.6 60.1 291 244 15.5 21.7 45.4 92.9 91.8 8 | 34.3 175 165 |
| Dhaka 20.8 57.6 318 264 11.5 15.5 31.6 92.8 89.2 8 | 39.3 183 170 |
| Khulna 14.5 50.7 210 208 12.5 23.4 43.0 96.4 90.0 8 | 37.0 106 127 |
| Rajshahi 15.9 59.9 214 212 17.7 9.2 43.8 94.4 93.5 8 | 37.2 128 138 |
| Rangpur 13.6 57.8 193 190 8.4 15.7 43.7 92.2 89.5 8 | 39.2 112 123 |
| Sylhet 19.6 48.1 99 136 22.7 36.5 53.4 86.4 96.6 7 | '6.2 48 80 |
| Mymensingh 17.0 64.2 122 151 26.9 11.1 34.5 89.4 88.9 8 | 34.1 78 107 |
| Total 18.2 56.0 1,557 15.0 18 41.2 92.3 89.9 8 | 36.1 873 994 |
| Total excluding CCs 43.6 60.7 563 1.069 8.0 15.9 31.9 97.9 97.0 8 | 36.6 342 736 |
| Total public 10.9 53.8 1.345 1.241 17.2 20.0 44.9 91.1 88.1 E | 34.4 724 757 |
| Total public excluding CCs 30.8 55.0 351 753 10.8 21.5 38.7 97.8 95.7 6 | 30.8 193 499 |

¹ Providers in the facility diagnose, prescribe treatment for, or manage patients with hypertension.

² At least one interviewed provider of hypertension services reported receiving in-service training in hypertension services during the 24 months preceding the survey. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ At least one interviewed provider of hypertension services reported ever receiving in-service training in hypertension services. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

⁴ Functioning stethoscope available at the service site on the day of the survey.

⁵ Functioning digital blood pressure machine or manual sphygmomanometer with stethoscope available at the service site on the day of the survey.

Table 8.8: Availability of essential medicines and commodities for hypertension

Among facilities offering both diagnosis and treatment services for hypertension, the percentages having the indicated essential medicines and commodities observed at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of fa | cilities offering the indicat | nsion that have | Number of facilities offering | Number of facilities offering | |
|---------------------------|-------------------------------|----------------------------------|--|---|---|---|
| Background characteristic | ACE inhibitors (enalapril) | Thiazide | Angiotensin receptor blockers (losartan) | Calcium channe blockers (amlodipine/ nifedipine) | both diagnosis and treatment services for hypertension (weighted) | both diagnosis and treatment services for hypertension (unweighted) |
| Facility type | | | | | | |
| District and upazila | | | | | | |
| public facilities | 1.9 | 25.8 | 84.4 | 66.3 | 33 | 233 |
| DH | 3.5 | 10.5 | 87.7 | 70.2 | 4 | 57 |
| MCWC | 0.0 | 0.0 | 3.0 | 0.0 | 2 | 31 |
| UHC | 1.9 | 30.1 | 90.5 | 71.1 | 27 | 145 |
| Union-level public | | | | | | |
| facilities | 0.0 | 3.4 | 29.0 | 24.5 | 75 | 120 |
| UHFWC | 0.0 | 0.0 | 0.0 | 0.0 | 26 | 35 |
| USC/RD | 0.0 | 5.3 | 44.5 | 37.6 | 49 | 85 |
| Community clinic | | | | | | |
| (CC) | 0.0 | 0.0 | 0.0 | 0.0 | 38 | 19 |
| NGO clinic/hospital | 0.5 | 2.9 | 20.1 | 37.9 | 19 | 85 |
| Private hospital | 5.5 | 14.5 | 48.3 | 53.7 | 118 | 130 |
| Location | | | | | | |
| Urban | 4.4 | 13.8 | 49.4 | 54.4 | 154 | 362 |
| Rural | 0.4 | 5.9 | 26.9 | 21.2 | 129 | 225 |
| Division | | | | | | |
| Barishal | 4.5 | 15.4 | 45.6 | 39.4 | 15 | 56 |
| Chattogram | 0.1 | 8.8 | 47.7 | 46.1 | 72 | 108 |
| Dhaka | 4.9 | 15.9 | 43.7 | 54.1 | 66 | 91 |
| Khulna | 0.0 | 7.2 | 30.9 | 28.7 | 30 | 73 |
| Rajshahi | 6.0 | 8.3 | 36.4 | 27.1 | 34 | 75 |
| Rangpur | 1.7 | 4.0 | 22.3 | 21.7 | 26 | 70 |
| Sylhet | 1.6 | 12.6 | 32.7 | 29.5 | 19 | 50 |
| Mymensingh | 2.1 | 6.1 | 34.4 | 35.1 | 21 | 64 |
| Total | 2.5 | 10.2 | 39.2 | 39.2 | 283 | 587 |
| Total excluding CCs | 2.9 | 11.8 | 45.2 | 45.3 | 245 | 568 |
| Total public | 0.4 | 7.7 | 34.2 | 27.8 | 146 | 372 |
| Total public excluding | 0.6 | 10 1 | 46.2 | 37.5 | 108 | 353 |
| 000 | 0.0 | 10.4 | 70.2 | 07.0 | 100 | 000 |

Table 8.9: Availability of cervical cancer diagnosis/screening services and guidelines, trained staff, and equipment

Among all facilities, the percentages offering diagnosis/screening services for cervical cancer, and among facilities offering diagnosis/screening services for cervical cancer, the percentages having guidelines, at least one staff member recently trained on cervical cancer screening, and the indicated equipment observed to be available at the service site on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | Percentage of facilities offering | | | Percentage of fac diagnosis/screening s cancer tha | cilities offering ervices for cervical it have: | Equipment | Number of facilities | Number of facilities offering diagnosis/ |
|-----------------------------|--------------------------------------|------------|--------------|--|---|-----------|----------------------|--|
| Desturyed | diagnosis/ screening | | | Guidelines for | | | screening services | screening services |
| characteristic | cancer ¹ | (weighted) | (unweighted) | cancer | Trained staff ² | Speculum | (weighted) | (unweighted) |
| Facility type | | | | | | | | |
| District and upazila public | | | | | | | | |
| facilities | 78.4 | 41 | 319 | 41.3 | 49.2 | 97.1 | 33 | 223 |
| DH | 85.5 | 4 | 62 | 47.2 | 45.3 | 98.1 | 4 | 53 |
| MCWC | 29.5 | 7 | 100 | 29.9 | 16.5 | 100.0 | 2 | 30 |
| UHC | 89.3 | 30 | 157 | 41.4 | 52.4 | 96.7 | 27 | 140 |
| Union-level public | | | | | | | | |
| facilities | 1.2 | 310 | 434 | 18.3 | 18.3 | 100.0 | 3 | 6 |
| UHFWC | 0.6 | 225 | 293 | 0.0 | 0.0 | 100.0 | 2 | 2 |
| USC/RD | 2.9 | 85 | 141 | 28.9 | 28.9 | 100.0 | 1 | 4 |
| Community clinic (CC) | 1.9 | 994 | 488 | 0.0 | 55.7 | 8.0 | 19 | 9 |
| NGO clinic/hospital | 20.6 | 29 | 127 | 30.3 | 37.2 | 67.7 | 6 | 26 |
| Private hospital | 9.7 | 183 | 189 | 20.3 | 26.2 | 97.2 | 18 | 18 |
| Location | | | | | | | | |
| Urban | 18.1 | 239 | 514 | 30.1 | 38.0 | 94.2 | 43 | 195 |
| Rural | 2.7 | 1,318 | 1,043 | 18.4 | 49.2 | 49.6 | 36 | 87 |
| Division | | | | | | | | |
| Barishal | 3.5 | 111 | 152 | 25.0 | 38.5 | 100.0 | 4 | 25 |
| Chattogram | 6.0 | 291 | 244 | 30.5 | 44.3 | 69.4 | 17 | 52 |
| Dhaka | 7.6 | 318 | 264 | 26.5 | 37.7 | 78.0 | 24 | 55 |
| Khulna | 2.9 | 210 | 208 | 16.0 | 48.5 | 58.1 | 6 | 31 |
| Rajshahi | 3.3 | 122 | 212 | 22.5 | 39.7 | 97.1 | 7 | 34 |
| Rangpur | 4.9 | 214 | 190 | 23.6 | 50.8 | 53.0 | 10 | 39 |
| Sylhet | 6.9 | 193 | 136 | 12.9 | 40.8 | 88.6 | 7 | 27 |
| Mymensingh | 3.4 | 99 | 151 | 28.7 | 59.5 | 47.3 | 4 | 19 |
| Total | 5.1 | 1,557 | 1,557 | 24.7 | 43.2 | 73.7 | 79 | 282 |
| Total excluding CCs | 10.6 | 563 | 1,069 | 32.5 | 39.2 | 94.4 | 60 | 273 |
| Total public | 4.1 | 1,345 | 1,241 | 25.6 | 49.3 | 66.8 | 55 | 238 |
| Total public excluding CCs | 10.3 | 351 | 753 | 38.8 | 45.9 | 97.4 | 36 | 229 |

¹ Providers in the facility diagnose/screen for cervical cancer.

² At least one interviewed provider of cervical cancer diagnosis/screening disease services reported receiving in-service training in cervical cancer during the 24 months preceding the survey. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

Key Findings

- One in four of the public facilities (24%) provides tuberculosis (TB) diagnostic, treatment, and/or treatment follow-up services (Table 9.1 and Figure 9.1).
- TB services are available in almost all DHs (92%) and UHCs (99%). About a quarter of private hospitals (23%) and a small percentage of NGO facilities (3%) offer TB services (Table 9.1 and Figure 9.1).
- The availability of TB services in public facilities has notably increased (from 7% to 24%) between 2017 and 2022. This can be due to the sharp increment in the availability of TB service in CCs (surging from 4% to 25%) and USC/RDs (rising from 8% to 29%) (Table 9.1 and Figure 9.1).
- Among facilities offering any TB services, the majority of DHs (79%) and UHCs (77%) have any guidelines for TB services. However, only 1% of private hospitals have guidelines for TB services (Table 9.3 and Figure 9.2).
- Three out of five public facilities (61%) offering any TB service have at least one staff member who has ever received in-service training on TB. Nearly half of the private hospitals (44%) and a quarter of the NGOs (26%) offering TB services have staff ever trained on TB (Table 9.3 and Figure 9.3).
- About six in ten DHs (58%) and UHCs (57%) that provide any TB services have the capacity to perform TB diagnostic smear microscopy on the day of the survey. However, among public facilities offering the service, only 6% have TB smear microscopy diagnostic capacity (Table 9.3 and Figure 9.4).
- Among facilities offering any TB service, four in five DHs (83%), private hospitals (78%), and around half (46%) of the UHCs have TB X-ray machines (**Table 9.2**).
- The majority of DHs (72%), UHCs (95%), and almost half of USC/RDs (43%) that offer any TB services have first-line medicines for treating TB. However, first-line TB medications are available in 2% of private facilities (Table 9.3 and Figure 9.5).
- The availability of first-line TB treatment in public facilities has dropped by more than half from 41% in 2017 to 16% in 2022 (**Table 9.3** and **Figure 9.5**).
- The average readiness score of total public health facilities (excluding CCs) for providing TB services is 1.8 out of 4. This score is relatively higher at DHs (3.0) and UHCs (3.1) (**Table 9.3** and **Figure 9.6**).

9.1 BACKGROUND

B angladesh is identified as one of the 30 high-burden countries for tuberculosis (TB) according to the World Health Organization (WHO). The estimated incidence rate for all forms of TB in Bangladesh is 221 per 100,000 population, as per the WHO Global Tuberculosis Report 2022, with a case notification rate of 181 per 100,000 population. In 2021, the estimated number of incident cases of multidrug-resistant TB (MDR-TB)/rifampicin-resistant TB (RR-TB) in Bangladesh was 4.5 thousand per year. The trend in the estimated absolute number (in thousands) of TB deaths (including both HIV-positive and HIV-negative TB) in Bangladesh has shown a decline, decreasing from 92,000 deaths in 2000 to 43,000 in 2021.

The effort to combat TB was integrated into the Millennium Development Goals (MDGs), and in the Sustainable Development Goals (SDGs), it has been positioned within health-related targets to mitigate the TB epidemic (United Nations n.d.). The target set is to reduce TB incidence by 90% and TB-related deaths by 95%, by the year 2035 compared to the levels in 2015 (WHO Tuberculosis Fact Sheet 2016).

The Mycobacterium Disease Control (MBDC) unit, under the DGHS and a part of the Bangladesh MOHFW, is responsible for overseeing the National Tuberculosis Control Program (NTP). Collaborating with 44 national and international NGOs, the MBDC implements the NTP. The primary development partners in TB initiatives in Bangladesh are the Global Fund, USAID, and WHO. These partners provide support to the Bangladesh government through various implementing agencies, including the Bangladesh Rural Development Committee (BRAC); icddr,b; the Damien Foundation; and Management Science for Health (MSH). Each organization fulfills distinct roles in the overall process—BRAC, for instance, supports TB service delivery through directly observed treatment, short-course (DOTS) centers.

Through the NTP, TB diagnostic and treatment services are available free of charge in public and private facilities throughout the country. Diagnosis and care for TB are provided through all UHCs and DHs, 44 chest disease clinics, 8 chest disease hospitals linked to these clinics, 4 divisional chest disease hospitals, DOTS centers within public and private medical college hospitals, workplaces, prisons, combined military hospitals, and other defense hospitals run by Border Guard Bangladesh (BDR), the Bangladesh Police, and the Bangladesh Ansar (Bangladesh Bureau of Statistics et al. 2016). The goal is to reduce morbidity, mortality, and transmission of TB until the disease is no longer a public health problem. Union-level public health facilities or CCs are not mandated to provide TB services; these facilities primarily screen and refer clients to UHCs or other facilities. The country has a laboratory network at the national, regional, district, and upazila levels. Smear microscopy is the primary TB diagnosis tool in the country. There are 1,145 microscopy centers and 230 GeneXpert machines in operation under NTP to detect and confirm TB cases (National Strategic Plan for TB Control 2021–2025).

The DOTS strategy was introduced by the NTP in 1993, and in 2014, the MOHFW mandated the notification of TB. The NTP has been actively implementing the Stop TB Strategy since 2006 (Annual TB Report of 2017). DOTS services have been accessible in all UHCs since 1998, and a 100% DOTS coverage milestone was achieved in 2007 (Annual TB Report 2017). The DOTS strategy encompasses the following:

- Secure political commitment with adequate and sustained financing.
- Ensure early case detection and diagnosis through quality-assured bacteriology.
- Provide standardized treatment with supervision and patient support.
- Ensure effective drug supply and management.
- Monitor and evaluate performance and impact.

This chapter provides an overview of TB services in secondary and primary health care facilities in Bangladesh and highlights the key aspects of TB-related client services, including the availability of diagnostic capacity, trained staff, and medicines. The chapter is organized as follows:

- Availability of TB services: Section 9.2, including Tables 9.1–9.2 and Figures 9.1–9.5 presents information on the availability of TB diagnostic and treatment services in Bangladesh.
- **Readiness of health facilities:** Section 9.3, including **Table 9.3** and **Figure 9.6** provides information on availability of the WHO-recommended items/tracer indicators necessary to offer quality TB services.

9.2 AVAILABILITY OF TB SERVICES

9.2.1 Service Provision

Achieving effective control of TB demands collaborative endeavors across all tiers. **Table 9.1** and **Figure 9.1** present data sourced from the 2022 BHFS, highlighting various essential indicators assessing the readiness of primary and secondary health facilities providing TB diagnostic, treatment, and follow-up services.

Availability of TB Services

- Nearly one-fourth (24%) of all public health facilities offer TB diagnostic, treatment, and follow-up services (**Table 9.1** and **Figure 9.1**).
- Among the public sectors, DHs and UHCs have higher availability than other facility types to provide TB services. Almost all DHs (92%) and UHCs (99%) offer TB diagnostic, treatment, and/or treatment follow-up services (**Table 9.1** and **Figure 9.1**).
- Only one-fourth of the CCs (25%), 23% of private hospitals, and a small proportion of NGO facilities (3%) are offering TB services (**Table 9.1** and **Figure 9.1**).
- Despite not being specifically assigned for TB services, 14% of union-level facilities (9% of UHFWCs and 29% of USC/RDs) offer TB services (**Table 9.1** and **Figure 9.1**).



Figure 9.1 Availability of TB services (TB diagnostic, treatment, and/or treatment followup services), by facility type

- Mymensingh (34%), Dhaka (28%), and Rajshahi (25%) divisions have higher percentages of facilities providing TB services compared to other divisions (Table 9.1).
- Fourteen percent of all public health facilities (7% when excluding CCs) provide referrals for TB diagnosis (**Table 9.1**).
- TB diagnostic services are available in 21% of public health facilities (excluding CCs). DHs (90%) and UHCs (98%) are the main providers of diagnostic services. Some of the USC/RDs (27%) and private hospitals (23%) also provide diagnostic services (**Table 9.1**).
- TB treatment and/or treatment follow-up services are offered by 13% of public health facilities (excluding CCs). The availability is higher in DHs (77%) and UHCs (93%) compared to the other facilities. Only 2% of private hospitals provide these services. TB treatment or treatment follow-up services are not available at all in NGOs (**Table 9.1**).

Guidelines

Guidelines for diagnosis and treatment of TB:

National guidelines for TB diagnosis and treatment are expected at all TB service sites. TB diagnosis and treatment guidelines are available in 41% of health facilities offering any TB services (excluding CCs). The percentage of facilities with TB diagnosis and treatment guidelines available is 77% for DHs and 75% for UHCs in 2022. However, only 13% of NGO facilities and 1% of private facilities have TB diagnosis and treatment guidelines (Table 9.1).

Guidelines for multidrug-resistant TB (MDR-TB):

MDR-TB diagnosis and treatment guidelines are available in 27% of public facilities offering TB services (excluding CCs). Less than half of DHs (44%) and six in ten UHCs (58%) have MDR-TB diagnosis and treatment guidelines in 2022. However, the availability of MDR-TB guidelines is lower in NGOs (13%) and private (1%) facilities (Table 9.1).

Guidelines for HIV and TB co-infection management:

In general, HIV and TB co-infection management guidelines are available in only about one-fifth (21%) of the public facilities (excluding CCs) offering TB services. These guidelines are present in nearly four in ten DHs and UHCs; however, only 13% of the NGO clinics and 1% of the private hospitals have guidelines for managing HIV and TB co-infection (Table 9.1).

Guidelines for TB infection control:

- Overall, 10% public facilities (excluding CCs) offering any TB services had TB infection control guidelines available at the time of the survey visit.
- Fourteen percent of the DHs, 16% of UHCs, 39% of NGO facilities, and 8% of private facilities have TB infection control guidelines in 2022 (**Table 9.1**).

Availability of any guidelines for TB services:

- Nearly half of the public facilities excluding CCs (44%) have any guidelines for TB services in 2022. Most of the DHs (79%) and UHCs (77%) and around one-third of the USC/RDs (32%) offering TB services have any guidelines on TB services (Table 9.3 and Figure 9.2).
- The availability of TB guidelines is lower in UHFWCs (4%), CCs (9%), NGOs (13%), and private facilities (1%) (Figure 9.2).



Figure 9.2 Availability of any guideline for TB services, by facility type

Percent

Trained Staff

- Nearly one-third of public facilities (31%) offering TB services have at least one provider trained in TB services within the previous 24 months (**Table 9.1**). However, 6 in 10 public facilities (61%) offering TB services have staff trained at any time (Table 9.3 and Figure 9.3).
- The majority of the DHs (61%) and UHCs (54%) have staff trained recently (within 24 months) in TB services (Table 9.1).
- Recently trained staff (within 24 months preceding the survey) are present in over one-quarter of the NGOs (26%) and one in five private facilities (19%) (Table 9.1).
- Around nine in ten DHs (91%) and UHCs (87%) have staff trained at any time in TB services. Moreover, 36% percent of the union-level facilities, 62% of CCs, and 26% of the NGO facilities have ever-trained staff (Table 9.3 and Figure 9.3).
- The availability of staff trained at any time in TB services in private hospitals has sharply increased from 22% in 2017 to 44% in 2022 (Table 9.3 and Figure 9.3).





Percent

9.2.2 Availability of Diagnostic Capacity and Medicines for TB Treatment

There are two types of TB: pulmonary TB (impacting the lungs) and extrapulmonary TB (affecting other body parts alongside the lungs). The primary focus of TB control services lies in identifying and effectively treating individuals with pulmonary TB. The most efficient and cost-effective method for screening pulmonary TB is through sputum microscopy using the Ziehl-Neelsen method. Additionally, sputum examination stands out as the most dependable means of diagnosing pulmonary TB. The 2022 BHFS evaluated the availability of TB diagnostic capabilities and medications for TB treatment in facilities providing any TB services. The findings from these evaluations are presented in Table 9.2.

Diagnostic Capacity

Early case detection is critical for TB control. The National Tuberculosis Control Program (NTP) has increased the number of TB service facilities with adequate diagnostic tools and equipment in recent years. Different types of diagnostic tools for TB include: microscopy examination of a patient's sputum (the most common and cost-effective), chest X-ray (radiological examination of the lungs), Gene Xpert, tuberculin skin tests (Mantoux tuberculin skin test), cultures of TB bacilli, molecular tests, fine needle aspiration cytology (FNAC), and biopsies.

In the 2022 BHFS, the assessment of TB diagnosis capacity or diagnosis of smear microscopy services included an observed functioning microscope, slides, and stains for Ziehl-Neelson tests or a fluorescence-LED microscope with all required stains.

- Considering the overall availability of TB diagnostic services, diagnostic smear microscopy is available in 25% of all public facilities (excluding CCs) offering TB services (**Table 9.2** and **Figure 9.4**).
- Between 2017 and 2022, the availability of TB diagnostic smear microscopies increased in DHs (41% to 58%) and UHCs (43% to 57%). Conversely, availability declined in NGO facilities and private hospitals (**Table 9.2** and **Figure 9.4**).

Figure 9.4 Availability of TB microscopy in health facilities, by facility type

Percent



- Among facilities offering any TB services, TB rapid diagnostic test kits are available in 19% of the public facilities (excluding CCs). The kits are observed in 32% of DHs, 45% of UHCs, and 7% of private hospitals in 2022 (**Table 9.2**).
- TB X-ray equipment is available in 22% of the public facilities (excluding CCs) that offer any TB services. Eighty-three percent of DHs, 46% of UHCs and 78% of private hospitals have the capacity to conduct TB X-rays, among facilities that offer any TB services (**Table 9.2**).

 Union and lower-level health facilities that offer any TB services lack the capacity to provide most of the TB diagnostic services (Table 9.2).

Medicine

- More than half (54%) of the public facilities (excluding CCs) offering TB services had first-line treatment medications available on the day of survey (**Table 9.2** and **Figure 9.5**).
- First-line TB medications were observed in 72% of DHs and 95% of UHCs that offer any TB services on the day of survey. However, only 3% of CCs and less than 1% of the NGO facilities have first-line TB medication available in 2022 (**Table 9.2** and **Figure 9.5**).
- Among facilities offering any TB services, injectable streptomycin is available in some of the DHs (4%), UHCs (3%), and private hospitals/clinics (1%) (**Table 9.2**).





Percent

9.3 READINESS OF HEALTH FACILITIES TO PROVIDE TB SERVICES

The WHO recommends assessing readiness to provide TB services based on the availability of specific items/tracer indicators. In this section of the report, data from the 2022 BHFS are used to construct a slightly less restrictive and Bangladesh-context-appropriate version of the WHO measures. The measures include the following four items/tracer indicators:

Guidelines:

- Any guidelines for TB (National guidelines for the diagnosis and treatment of TB, guidelines for the management of HIV and TB co-infection, or guidelines related to MDR-TB treatment)

- Trained staff:
 - At least one provider ever receiving in-service training on TB
- Diagnostic capacity:
 - TB microscopy
- Medicines:
 - First-line TB medicines

Average Readiness Score of the Health Facilities

The readiness scores are presented on a scale of 0–4, indicating the facility's preparedness to provide TB services. The lower scores in 2022 suggest a potential decrease in readiness compared to previous years.

- Among public health facilities (excluding CCs), the average readiness score for providing TB services is 1.8 in 2022 (**Table 9.2** and **Figure 9.6**).
- DHs (3.0) and UHCs (3.1) maintained a comparatively higher level of readiness than the other facilities. However, the readiness scores declined in UHFWCs, USC/RDs, NGOs, and private facilities. The average readiness score of CCs has remained unchanged (<1.0) since 2017 (Figure 9.6).

Figure 9.6 Average readiness score (out of 4) of health facilities for providing TB services



Average Score

TUBERCULOSIS

LIST OF TABLES

- Table 9.1 Availability of tuberculosis services, guidelines, and trained staff for tuberculosis services
- Table 9.2 Diagnostic capacity and availability of medicines for tuberculosis treatment
- Table 9.3 Readiness of health facilities to provide TB services

Table 9.1: Availability of tuberculosis services, guidelines, and trained staff for tuberculosis services

Among all facilities, the percentages offering any tuberculosis (TB) referral, diagnostic, treatment, and/or treatment follow-up services, and among facilities offering any TB services, the percentages having TB guidelines and at least one staff member recently trained in TB services, by background characteristics, Bangladesh HFS 2022

| | Percentage of all facilities offering: | | | | | Percentage of facilities offering any TB services that have guidelines for: | | | | | | Number of facilities | Number of facilities | |
|--|---|--|--|---|--|--|--|--|---|---|--|--|---|---|
| Background characteristic | Any referral for TB diagnosis ¹ | Any TB diagnostic services ² | Any TB diagnostic or referral services | Any TB treatment and/or treatment follow- up services ³ | Any TB diagnostic, treatment, and/or treatment follow- up services | Number of facilities (weighted) | Number of facilities (unweighted) | Diagnosis and treatment of TB | Diagnosis and treatment of MDR-TB | Manage- ment of HIV and TB co- infection | TB infection control | Trained staff ⁴ | offering any TB diagnostic, treatment, and/or treatment follow- up services (weighted) | offering any TB diagnostic, treatment, and/or treatment follow- up services (unweighted) |
| Facility type | | | | | | | | | | | | | | |
| District and upazila public facilities DH MCWC | 32.3 45.2 0.0 | 81.5 90.3 7.9 | 82.0 91.9 7.9 | 75.3 77.4 0.0 | 82.2 91.9 7.9 | 41 4 7 | 319 62 100 | 74.0 77.2 | 55.5 43.9 | 39.1 40.4 | 16.3 14.0 | 54.2 61.4 | 34 4 | 220 57 |
| UHC | 38.2 | 98.1 | 98.6 | 93.3 | 98.8 | 30 | 157 | 75.0 | 58.2 | 39.7 | 16.4 | 54.0 | 29 | 155 |
| Union-level public facilities UHFWC USC/RD | 3.7 1.6 9.3 | 13.3 8.1 26.8 | 14.2 9.0 27.9 | 4.3 1.2 12.6 | 14.2 8.5 29.3 | 310 225 85 | 434 293 141 | 16.4 4.0 25.9 | 5.1 0.0 9.0 | 6.5 4.0 8.4 | 4.5 10.4 0.0 | 4.0 0.0 7.0 | 44 19 25 | 65 23 42 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 17.0 0.8 2.7 | 22.0 2.7 23.4 | 31.4 3.1 23.4 | 4.7 0.0 1.6 | 24.5 2.7 23.4 | 994 29 183 | 488 127 189 | 6.6 13.2 0.9 | 1.5 13.2 0.9 | 2.3 13.2 0.9 | 9.5 39.3 8.4 | 32.4 26.1 19.2 | 244 1 43 | 119 4 48 |
| Location Urban Rural | 6.2 14.0 | 28.0 20.5 | 28.1 27.8 | 9.0 5.5 | 28.0 22.7 | 235 1,322 | 514 1,043 | 26.8 10.4 | 17.9 4.5 | 12.7 4.6 | 11.5 9.0 | 28.6 29.6 | 66 300 | 197 259 |
| Division Barishal Chattogram Dhaka Khulna Rajshahi Rangpur Sylhet Mymensingh | 8.0 9.3 14.0 10.7 18.3 9.4 12.0 22.4 | 16.4 21.1 24.0 20.7 24.4 17.8 17.5 28.2 | 20.3 25.3 30.2 26.9 34.4 21.2 25.7 37.1 | 6.9 4.8 7.5 2.9 5.3 4.9 2.9 15.2 | 16.4 21.8 28.0 21.6 25.4 19.0 17.5 34.1 | 111 291 317 210 214 193 99 122 | 152 244 264 208 212 190 136 151 | 30.0 12.6 11.5 6.9 19.3 13.2 23.7 6.3 | 8.3 6.3 4.7 8.6 5.9 7.2 11.3 9.4 | 5.2 8.1 4.0 8.0 3.1 7.2 15.4 5.0 | 4.8 18.2 11.0 1.5 1.1 7.5 13.6 14.3 | 25.8 25.1 32.2 13.3 34.0 29.8 20.8 46.7 | 18 64 89 45 54 37 17 42 | 40 71 89 59 59 52 36 50 |
| Total | 12.8 | 21.7 | 27.8 | 6.0 | 23.5 | 1,557 | 1,557 | 13.3 | 6.9 | 6.1 | 9.5 | 29.4 | 366 | 456 |
| Total excluding CCs | 5.3 | 21.0 | 21.6 | 8.4 | 21.6 | 563 | 1,069 | 27.0 | 17.7 | 13.6 | 9.4 | 23.5 | 122 | 337 |
| Total public | 14.4 | 21.8 | 29.0 | 6.7 | 23.9 | 1,345 | 1,241 | 15.0 | 7.7 | 6.8 | 9.5 | 30.8 | 322 | 404 |
| Total public excluding CCs | 7.1 | 21.3 | 22.2 | 12.7 | 22.2 | 351 | 753 | 41.1 | 27.0 | 20.7 | 9.7 | 25.8 | 78 | 285 |

Note: Guidelines and trained staff indicators presented in this table comprise the staff and training domain for assessing readiness to provide TB services within the health facility assessment methodology proposed by the WHO and USAID (2012); MDR-TB = multidrug-resistant tuberculosis; "-" means facilities do not have provision of TB services.

¹ Facility reports that it refers clients outside the facility for TB diagnosis, and a register was observed indicating clients who had been referred for TB diagnosis.

² Facility reports that providers in the facility make a diagnosis of TB by using any of the following methods: sputum smear only, X-ray only, either sputum or X-ray, or both sputum and X-ray.

³ Facility reports that it adheres to one of the following TB treatment regimens or approaches: Directly observe for two months and follow up for four months; directly observe for six months; follow up clients only after the first two months of

direct observation elsewhere; diagnose and treat clients while in the facility as inpatients and then discharge elsewhere for follow up; provide clients with the full treatment with no routine direct observation phase; diagnose, prescribe, or provide medicines with no follow up.

⁴ At least one interviewed provider of any of the following TB services reported receiving in-service training relevant to the particular TB service during the 24 months preceding the survey: TB diagnosis based on sputum tests using AFB smear microscopy; TB diagnosis based on clinical symptoms or TB diagnostic algorithm; treatment prescription for TB; treatment follow-up services for TB; direct observation treatment short-course (DOTS) strategy; management of HIV and TB co-infection; MDR-TB treatment and identification of need for referral. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

Table 9.2: Diagnostic capacity and availability of medicines for tuberculosis treatment

Among facilities offering any tuberculosis (TB) diagnostic, treatment, and/or treatment follow-up services, the percentages that had TB and HIV diagnostic capacity and medicines for TB treatment available in the facility on the day of the survey, by background characteristics, Bangladesh HFS 2022

| | F | Percentage of fac TB dia | cilities that ha agnostic capa | ve the following acity | | Percentage of | facilities that have: | Percentage of have the follow for treat | of facilities that wing medicines ating TB | Number of facilities offering any TB diagnostic, treatment, | Number of facilities offering any TB diagnostic, treatment, |
|--|-------------------------|-------------------------------------|-----------------------------------|-------------------------------------|--------------------|---|---|--|--|---|---|
| Background characteristic | Microscope ¹ | TB smear microscopy ² | Culture medium ³ | TB rapid diagnostic test kits | TB X-ray | HIV diagnostic capacity ⁴ | System for diagnosing HIV among TB clients⁵ | First-line treatment for TB ⁶ | Injectable streptomycin | and/or treatment follow-up services (weighted) | and/or treatment follow-up services (unweighted) |
| Facility type | | | | | | | | | | | |
| District and upazila public facilities | 87.0 | 55.7 | 2.2 | 42.7 | 49.9 | 32.0 | 52.6 | 90.5 | 2.6 | 34 | 220 |
| DH | 96.5 | 57.9 | 1.8 | 31.6 | 82.5 | 36.8 | 36.8 | 71.9 | 3.5 | 4 | 57 |
| MCWC UHC | - 87.2 | - 56.5 | - 2.3 | - 45.1 | 46.4 | - 31.9 | - 55.8 | 94.8 | - 2.5 | - 29 | - 155 |
| Union-level public facilities | 2.5 | 1.6 | 0.0 | 0.0 | 0.0 | 1.8 | 11.2 | 26.3 | 0.0 | 44 | 65 |
| UHFWC USC/RD | 0.0 4.4 | 0.0 2.8 | 0.0 0.0 | 0.0 0.0 | 0.0 0.0 | 0.0 3.2 | 8.5 13.3 | 4.5 43.0 | 0.0 0.0 | 19 25 | 23 42 |
| Community clinic (CC) NGO clinic/hospital Private hospital | 0.0 59.5 95.3 | 0.0 0.0 16.2 | 0.0 0.0 3.2 | 0.0 0.0 7.0 | 0.0 0.0 78.3 | 0.0 13.2 34.4 | 1.0 13.2 9.0 | 3.1 0.0 1.6 | 0.0 0.0 0.9 | 244 1 43 | 119 4 48 |
| Location | | | | | | | | | | | |
| Urban Rural | 90.2 4.2 | 27.4 2.8 | 2.9 0.1 | 17.5 2.0 | 68.4 1.8 | 32.1 1.8 | 21.2 5.0 | 28.1 10.7 | 1.4 0.1 | 66 300 | 197 259 |
| Division Barishal | 20.6 | 10.0 | 0.4 | 0 0 | 10.7 | 27 | 2.4 | 25.4 | 1 2 | 19 | 40 |
| Chattogram Dhaka | 28.4 23.2 | 11.5 6.0 | 2.5 0.5 | 5.6 3.4 | 22.6 15.9 | 10.0 11.1 | 15.0 7.2 | 13.5 13.1 | 0.4 0.1 | 64 89 | 40 71 89 |
| Khulna Rajshahi Rangpur | 12.2 16.6 14.9 | 4.8 6.9 7.7 | 0.0 0.0 0.0 | 7.2 3.0 4.8 | 8.7 12.5 8.5 | 3.1 5.1 6.8 | 5.1 5.7 9.3 | 9.2 8.8 18.0 | 0.0 0.0 1.0 | 45 54 37 | 59 59 52 |
| Sylhet Mymensingh | 29.3 11.2 | 11.2 3.6 | 0.0 0.0 | 8.8 2.9 | 21.4 6.2 | 14.3 1.7 | 12.3 4.0 | 16.0 18.1 | 1.8 0.0 | 17 42 | 36 50 |
| Total | 19.7 | 7.3 | 0.6 | 4.8 | 13.8 | 7.3 | 8.0 | 13.9 | 0.3 | 366 | 456 |
| Total excluding CCs | 59.2 | 21.8 | 1.7 | 14.4 | 41.5 | 21.8 | 22.0 | 35.3 | 1.0 | 122 | 337 |
| Total public | 9.5 | 6.1 | 0.2 | 4.5 | 5.3 | 3.6 | 7.8 | 15.5 | 0.3 | 322 | 404 |
| Total public excluding CCs | 39.3 | 25.2 | 1.0 | 18.6 | 21.7 | 14.9 | 29.2 | 54.2 | 1.1 | 78 | 285 |

Note: The indicators presented in this table comprise the diagnostics and medicines and commodities domains for assessing readiness to provide services for TB within the health facility assessment methodology proposed by WHO and USAID (2012); "-" means facilities do not have provision of TB services. ¹ Functioning light microscope or fluorescence-LED microscope available on the day of visit.

² Functioning microscopes, slides, and all stains for Ziehl-Neelsen test (carbol-fuchsin, sulfuric acid, methyl blue) or fluorescence-LED microscope with all stains (sulfuric acid, methyl blue, auramine stain) were available in the facility on the

day of the survey visit. ³ Solid or liquid culture medium (e.g., MGIT 960).

⁴ HIV rapid diagnostic test kits available in TB service area.
⁵ Record or register indicating TB clients who had been tested for HIV.

⁶ Four-drug fixed-dose combination (4FDC) available, or else isoniazid, pyrazinamide, rifampicin, and ethambutol are all available, or a combination of these medicines, to provide first-line treatment.

Table 9.3: Readiness of health facilities to provide TB services

Among facilities that offer any tuberculosis (TB) diagnostic, treatment, and/or treatment follow-up services, the percentages that had the indicated items available in the facility on the day of the survey, and the average readiness score, by background characteristics, Bangladesh HFS 2022

| Background characteristic | Any guidelines for TB ¹ | Trained staff at any time ² | TB microscopy ³ | First-line TB medications ⁴ | Average readiness score (out of 4) ⁵ | Number of facilities offering any TB diagnostic, treatment, and/or treatment follow-up services (weighted) | Number of facilities offering any TB diagnostic, treatment, and/or treatment follow-up services (unweighted) |
|--|---------------------------------------|--|-------------------------------|--|---|--|--|
| Facility type | | | | | | | |
| District and upazila public facilities | 75.7 | 86.2 | 55.7 | 90.5 | 3.1 | 34 | 220 |
| DH | 78.9 | 91.2 | 57.9 | 71.9 | 3.0 | 4 | 57 |
| MCWC | - | - | - | - | - | - | - |
| UHC | 76.7 | 86.7 | 56.5 | 94.8 | 3.1 | 29 | 155 |
| Union-level | | | | | | | |
| public facilities | 19.9 | 36.2 | 1.6 | 26.3 | 0.8 | 44 | 65 |
| UHFWC | 4.0 | 28.7 | 0.0 | 4.5 | 0.4 | 19 | 23 |
| USC/RD | 32.1 | 42.0 | 2.8 | 43.0 | 1.2 | 25 | 42 |
| Community clinic (CC) | 9.0 | 62.1 | 0.0 | 3.1 | 0.7 | 244 | 119 |
| NGO clinic/hospital | 13.2 | 26.1 | 0.0 | 0.0 | 0.4 | 1 | 4 |
| Private hospital | 0.9 | 43.5 | 16.2 | 1.6 | 0.6 | 43 | 48 |
| Location | | | | | | | |
| Urban | 27.7 | 55.0 | 27.4 | 28.1 | 1.4 | 66 | 197 |
| Rural | 12.9 | 59.8 | 2.8 | 10.7 | 0.9 | 300 | 259 |
| Division | | | | | | | |
| Barishal | 30.9 | 64.1 | 10.0 | 25.4 | 1.3 | 18 | 40 |
| Chattogram | 17.4 | 47.2 | 11.5 | 13.5 | 0.9 | 64 | 71 |
| Dhaka | 11.6 | 65.2 | 6.0 | 13.1 | 1.0 | 89 | 89 |
| Khulna | 12.6 | 48.1 | 4.8 | 9.2 | 0.7 | 45 | 59 |
| Rajshahi | 19.3 | 66.8 | 6.9 | 8.8 | 1.0 | 54 | 59 |
| Rangpur | 13.2 | 57.4 | 1.1 | 18.0 | 1.0 | 37 | 52 |
| Sylnet | 23.7 | 45.0 | 11.2 | 16.0 | 1.0 | 17 | 36 |
| Mymensingn | 11.7 | 70.0 | 3.6 | 18.1 | 1.0 | 42 | 50 |
| Total | 15.6 | 59.0 | 7.3 | 13.9 | 1.0 | 366 | 456 |
| Total excluding CCs | 28.7 | 52.7 | 21.8 | 35.3 | 1.4 | 122 | 337 |
| Total public | 17.5 | 61.1 | 6.1 | 15.5 | 1.0 | 322 | 404 |
| Total public excluding CCs | 44.2 | 58.0 | 25.2 | 54.2 | 1.8 | 78 | 285 |

Note: "-" Means facilities do not have provision of TB services.

¹ National guidelines for the diagnosis and treatment of TB, guidelines for the management of HIV and TB co-infection, or guidelines related to MDR-TB treatment.

² At least one interviewed provider of any of the following TB services reported receiving in-service training relevant to the particular TB service at any time: TB diagnosis based on sputum tests using AFB smear microscopy; TB diagnosis based on clinical symptoms or TB diagnostic algorithm; treatment prescription for TB; treatment follow-up services for TB; direct observation treatment short-course (DOTS) strategy; management of HIV and TB co-infection; MDR-TB treatment and identification of need for referral. The training must have involved structured sessions; it does not include individual instruction that a provider might have received during routine supervision.

³ Functioning microscopes, slides, and all stains for Ziehl-Neelsen test (carbol-fuchsin, sulfuric acid, methyl blue) or fluorescence-LED microscope with all stains (sulfuric acid, methyl blue, auramine stain) were available in the facility on the day of the survey visit.

⁴ Four-drug fixed-dose combination (4FDC) available, or else isoniazid, pyrazinamide, rifampicin, and ethambutol are all available, or a combination of these medicines, to provide first-line treatment.

⁵Average readiness score is the average number of items (out of 4 items) available for providing TB diagnostic, treatment, and/or treatment follow-up services.

REFERENCES

Ahmed, S. M., Hossain, A., Raja Chowdhury, A. M., & Bhuiya, A. 2011. *The health workforce crisis in Bangladesh: shortage, inappropriate skill-mix and inequitable distribution.* Human Resources for Health, 9(1). Retrieved from https://doi.org/10.1186/1478-4491-9-3

Bangladesh Bureau of Statistics (BBS), Centers for Disease Control and Prevention (CDC), Directorate General of Health Services, Ministry of Health and Family Welfare (Bangladesh), Global Fund to Fight Aids Tuberculosis and Malaria (GFATM), Institute of Tropical Medicine (Belgium), National Tuberculosis Control Program (NTP) (Bangladesh), Research Institute of Tuberculosis/Japan Anti-Tuberculosis Association (RIT/JATA), United States Agency for International Development (USAID). 2016. *Bangladesh National Tuberculosis Prevalence Survey 2015-2016*. Retrieved from https://ghdx.healthdata.org/record/bangladesh-national-tuberculosis-prevalence-survey-2015-2016

Bangladesh Bureau of Statistics (BBS), Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic Bangladesh. December 2023. Report on *Bangladesh Sample Vital Statistics (SVRS)* 2022. Dhaka, Bangladesh: Reproduction, Documentation & Publication Section, BBS.

Bangladesh Bureau of Statistics (BBS), Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic Bangladesh. June 2018. Report on *Bangladesh Sample Vital Statistics (SVRS) 2017*, p 63. Dhaka, Bangladesh: Reproduction, Documentation & Publication Section, BBS.

Directorate General of Health Services. 2018. *Multi-sectoral action plan for prevention and control of noncommunicable diseases 2018-2025*. Dhaka, Bangladesh: Directorate General of Health Services.

Directorate General of Health Services. 2022. *Health Bulletin 2020*. Dhaka, Bangladesh: Directorate General of Health Services.

 $https://dghs.portal.gov.bd/sites/default/files/files/dghs.portal.gov.bd/page/8983ee81_3668_4bc3_887e_c99645\\bbfce4/2022-09-20-12-31-58c5a0b12e3aad087eaa26c3ce0f1a7b.pdf$

Directorate General of Family Planning. 2023. *Renovation and Reconstruction Plan*. Dhaka, Bangladesh: Directorate General of Health Services. Accessed on 12th October, 2023.

Government of the People's Republic of Bangladesh. 2017. *Programme Implementation Plan, 4th Health, Population and Nutrition Sector Programme* (Vol. I). Dhaka, Bangladesh: Government of the People's Republic of Bangladesh.

Institute for Health Metrics and Evaluation: IHME. (n.d.). *Global Burden of Disease: GBD Collaborative Network, Global Burden of Disease Study 2019.* Retrieved from https://vizhub.healthdata.org/gbd-results

Islam, S.M., Purnat, T.D., Phuong, N.T., Mwingira, U., Schacht, K., Fröschl, G. 2014. *Non-Communicable Diseases (NCDs) in developing countries: a symposium report.* Globalization and Health. 2014;10(1). doi: 10.1186/s12992-014-0081-9

Kabir, A., Karim, M.N., Billah, B. 2022. *Health system challenges and opportunities in organizing noncommunicable diseases services delivery at primary healthcare level in Bangladesh: A qualitative study.* Front Public Health. Nov 9;10:1015245. doi: 10.3389/fpubh.2022.1015245. PMID: 36438215; PMCID: PMC9682236. Ministry of Health and Family Welfare, Government of the People's Republic of Bangladesh. 2021. *National strategic plan for tuberculosis control* 2021–2025. https://www.example.gov.bd/nsp-tb-control-2021-2025.pdf

Ministry of Health and Family Welfare (MOHFW) [Bangladesh]. 2017. 4th Health, Population and Nutrition Sector Programmme (4th HPNSP): Operational Plan for Maternal, Neonatal, Child, & Adolescent Health (MNC&AH), (January 2017-June 2024). Dhaka: Bangladesh; MOHFW.

Ministry of Health and Family Welfare (MOHFW) [Bangladesh]. 2012. *Bangladesh Population Policy 2012*. Dhaka, Bangladesh: MOHFW, Government of the People's Republic of Bangladesh.

Ministry of Health and Family Welfare (MOHFW) [Bangladesh]. *Public Health Expenditure Review 1997-2020*. Dhaka, Bangladesh: Health Economics Unit, Health Services Division, Ministry of Health and Family Welfare (MOHFW), Government of the People's Republic of Bangladesh.

Ministry of Health and Family Welfare (MOHFW) [Bangladesh]. 2012. *Health Care Financing Strategy* 2012-2032. Dhaka, Bangladesh: Health Economics Unit, Health Services Division, Ministry of Health and Family Welfare (MOHFW), Government of the People's Republic of Bangladesh.

Ministry of Health and Family Welfare Bangladesh (MOHFW) and World Health Organization (WHO) Bangladesh. 2021. *Assessment of Healthcare Providers in Bangladesh 2021*. Dhaka, Bangladesh: MOHFW, Human Resources Branch, Health Services Division and WHO Bangladesh.

National Institute of Population Research and Training (NIPORT), and icddr,b (International Centre for Diarrhoeal Disease Research, Bangladesh). 2023. *Bangladesh Health Facility Survey 2022*: Preliminary report. Dhaka, Bangladesh: NIPORT and icddr,b.

National Institute of Population Research and Training (NIPORT) and ICF. 2019. *Bangladesh Health Facility Survey 2017*. Dhaka, Bangladesh: NIPORT, ACPR, and ICF.

National Institute of Population Research and Training (NIPORT), Associates for Community and Population Research (ACPR), and ICF. 2018. *Bangladesh Health Facility Survey 2017: Preliminary Report*. Dhaka, Bangladesh: NIPORT, ACPR, and ICF.

National Institute of Population Research and Training (NIPORT), Associates for Community and Population Research (ACPR), and ICF International. 2016. *Bangladesh Health Facility Survey 2014*. Dhaka, Bangladesh: NIPORT, ACPR, and ICF International.

National Institute of Population Research and Training (NIPORT) and ICF. 2023. *Bangladesh Demographic and Health Survey 2022: Key Indicators Report*. Dhaka, Bangladesh, and Rockville, Maryland, USA: NIPORT and ICF.

National Institute of Population Research and Training (NIPORT), and ICF. 2020. *Bangladesh Demographic and Health Survey 2017-18*. Dhaka, Bangladesh, and Rockville, Maryland, USA: NIPORT and ICF.

National Institute of Population Research and Training (NIPORT), Mitra and Associates, and ICF International. 2017. *Bangladesh Demographic and Health Survey 2017: Preliminary Report*. Dhaka, Bangladesh, and Calverton, Maryland, USA.

National Institute of Population Research and Training (NIPORT), Mitra and Associates, and ICF International. 2016. *Bangladesh Demographic and Health Survey 2014*. Dhaka, Bangladesh, and Rockville, Maryland, USA: NIPORT, Mitra and Associates, and ICF International.

National Institute of Population Research and Training (NIPORT), Mitra and Associates, and ICF International. 2015. *Bangladesh Demographic and Health Survey 2014: Key Indicators Report*. Dhaka, Bangladesh and Rockville, Maryland, USA: NIPORT, Mitra and Associates, and ICF International.

National Institute of Population Research and Training (NIPORT), International Centre for Diarrhoeal Disease Research Bangladesh (icddr,b), and MEASURE Evaluation. 2017. *Bangladesh Maternal Mortality and Health Care Survey 2016: Preliminary Report.* Dhaka, Bangladesh, and Chapel Hill, NC, USA

National Institute of Population Research and Training (NIPORT), Mitra and Associates, & ORC Macro. 2003. *Bangladesh Maternal Health Services and Maternal Mortality Survey 2001*. Dhaka, Bangladesh: NIPORT, Mitra and Associates, and ORC Macro.

Pullar, J., Allen, L., Townsend, N., Williams, J., Foster, C., et al. 2018. *The impact of poverty reduction and development interventions on non-communicable diseases and their behavioral risk factors in low and lower-middle income countries: A systematic review.* PLOS ONE, 13(2): e0193378. https://doi.org/10.1371/journal.pone.0193378

Directorate General of Health Services, National Tuberculosis Control Program. 2017. *Tuberculosis Control in Bangladesh: Annual Report, 2017.* Dhaka, Bangladesh: Directorate General of Health Services (DGHS). http://ntp.gov.bd/ntp_dashboard/magazines_image/Annual%20Report-2017.pdf.

United Nations (n.d.). *Sustainable Development Goals: 17 Goals to Transform our World*. Retrieved from: https://www.un.org/en/exhibits/page/sdgs-17-goals-transform-world

World Health Organization (WHO). 2013. Service Availability and Readiness Assessment (SARA): An Annual Monitoring System for Service Delivery, Reference Manual. Version 2.1. http://www.who.int/ healthinfo/systems/SARA_Reference_Manual_Full.pdf.

World Health Organization (WHO). 2023. *Noncommunicable diseases*. Geneva, Switzerland: World Health Organization. Retrieved from: https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases

World Health Organization (WHO). 2023. *Tuberculosis fact sheet*. Geneva, Switzerland: World Health Organization. Retrieved from: https://www.who.int/news-room/fact-sheets/detail/tuberculosis

World Health Organization (WHO). 2022. *Global Tuberculosis Report 2022*. Geneva, Switzerland: World Health Organization, 2022. https://www.who.int/teams/global-tuberculosis-programme/tb-reports/global-tuberculosis-report-2022

World Health Organization (WHO). 2018. *Saving lives, spending less: a strategic response to noncommunicable diseases.* Geneva, Switzerland: World Health Organization; 2018 (WHO/NMH/NVI/18.8). Licence: CC BY-NC-SA 3.0 IGO. Retrieved from https://www.who.int/publications/i/item/WHO-NMH-NVI-18.8

World Health Organization (WHO). 2001. *Health-Care Waste Management: Rapid Assessment Tool for Country Level*. Geneva, Switzerland: World Health Organization.

World Health Organization (WHO), Division of Child Health and Development and Family and Reproductive Health. 1998. CHD 1996–1997 Report. Retrieved from http://apps.who.int/iris/bitstream/10665/66705/1/WHO_CHD_98.5.pdf.

World Bank. 2011. Non-Communicable Diseases (NCDs) – Bangladesh's Next Major Health Challenge. Washington, DC, USA: World Bank. Retrieved from https://documents1.worldbank.org/curated/en/687721505973301661/pdf/119403-BRI-P114171-PUBLIC-NCD-BD.pdf PERSONS INVOLVED IN THE SURVEY



NIPORT

Md. Shafiqul Islam Mr. Mohammed Ahsanul Alam Mr. Giyasuddin Ahamed Mr. Zahirul Islam

USAID/Bangladesh

Kanta Jamil

Dr. Mohammad Shahrear Farid

icddr,b

Dr. Shams El Arifeen

Dr. Ahmed Ehsanur Rahman Mr. Md. Abu Bakkar Siddique Dr. Shafiqul Ameen Dr. Twaha Mansurun Haque Dr. Md. Rakibul Hasan Dr. Sabit Saad Shafiq Dr. Sadman Sowmik Sarkar Dr. Ashiquzzaman Ms. Anindita Saha

D4I

Dr. Ahmed Al-Sabir Ms. Shusmita Khan Director General Director (Research) Ex. Sr. Research Associate Statistician

Senior Research, Monitoring and Evaluation Advisor, OPHN Project Management Specialist (Monitoring, Evaluation and Learning), Office of Population, Health and Nutrition

Senior Director, Maternal and Child Health Division Scientist Assistant Scientist Assistant Scientist Program Manager Study Physician Study Physician Study Physician Project Research Physician Statistical Officer

Consultant Knowledge Management and Communication Specialist

ACPR Survey Implementation Team

Prof. Dr. Sekander Hayat Khan Mr. A P M Shafiur Rahman Dr. K. M. Rezaul Haque Ms. Tauhida Nasrin Ms. Ilmul Jahan Tani Mr. S M Salamat Ullah Mr. Kh. Khairul Bashar Team Leader Co-team Leader Public Health Specialist Operations/ Logistics Manager IT Specialist Survey Specialist Data Processing Specialist

DATA PROCESSING TEAM

icddr,b

Mr. Qazi Sadeq-ur Rahman Mr. A.K.M. Tanvir Hossain Md. Toriqul Anam Mr. Md. Khairul Islam Bhuiyan

TRAINING TEAM

icddr,b

ACPR

Mr. Khandaker Khairul Bashar

ACPR

Dr. Shafiqul Ameen Dr. Twaha Mansurun Haque Dr. Md. Rakibul Hasan Dr. Sabit Saad Shafiq Dr. Sadman Sowmik Sarkar Dr. Sabrina Jabeen Dr. A K M Mahmudul Hassan Dr. Snigdho Protim Sikder Dr. Md. Ishtiak Anam Nobel Dr. Ashiquzzaman Dr. Tawsif Ismam Mr. Md. Abu Bakkar Siddique Dr. Md Abu Huzaifa Dr. Regan Barua Dr. Samia Tasnim Tamanna Dr. Md Moshiuzzaman Shakil Dr. Newmoon Rayeen Rahman Dr. Anika Nazia

QUALITY CONTROL OFFICERS

icddr,b

Dr. Ashiquzzaman Dr. Snigdho Protim Sikder Dr. Md. Ishtiak Anam Nobel Dr. Tawsif Ismam

ACPR

Dr. Md Abu Huzaifa Dr. Regan Barua Dr. Samia Tasnim Tamanna Dr. Md Moshiuzzaman Shakil Dr. Newmoon Rayeen Rahman Dr. Anika Nazia

MEDICAL OFFICER-CUM-ENUMERATOR

Dr. Abdullah Al Mamun Dr. Abdullah Al Noman Dr. Ahanaf Hossain Akif Dr. Amlan Rakshit Dr. Anika Naziat Majumder Dr. Anwara Begum Mitu Dr. Ayinuddin Ahmed Bhuivan Dr. Aysha Akther Dr. Debashish Biswas Dr. Imranul Hossain Dr. Jakia Sultana Jolly Dr. Kazi Adnan Mosharaf Dr. Lubna Rahman Dr. Mahbuba Mehjabin Dr. Maliha Kashem Chowdhury Dr. Mantasha Kifayat Mohona Dr. Md. Abir Hossain Dr. Md. Ahnaf Tahamid Noman Dr. Md. Faisal Ahmed Dr. Md. Humayun Kabir Dr. Md. Mahmud Hassan Khan Shovon Dr. Md. Shafaet ul Islam Dr. Md. Shakawat Hossain Dr. Md. Tauhidul Islam Dr. Meherab Hossain Dr. Mst. Mousume Aktar Dr. Nawshin Rubaiyat Dr. Sabrina Binte Shahadut Dr. Saif Muhammad Rifayee Dr. Samina Ferdousi Sweety Dr. Shahida Akter Nishat Dr. Shaikat Dey Dr. Shulakkhona Haque Dr. Sumiya Sharmin Dr. Tarigul Alam Dr. Zinia Afroz Dr. Zunayed Ahmad

MEDICAL ASSISTANT-CUM-ENUMERATOR

Mr. Abdul Hasan Mr. Ariful Islam Mr. Ibrahin Khalil Mr. Md. Abu Bakkar Siddik Mr. Md. Abu Hanif Mr. Md. Alomgir Hossain Sumon Mr. Md. Imram Hossain Mr. Md. Imranul Islam Mr. Md. Mahbub Hasan Mr. Md. Nazmul Haq Nannu Mr. Md. Nazmul Hossen Mr. Md. Nur Nobi Miah Mr. Md. Raihan Sheikh Mr. Md. Raisul Islam Refat Mr. Md. Saidur Rahman Mr. Md. Sajid Hossain Mr. Md. Sazzad Islam Mr. Md. Shahanur Ali Mr. Md. Shakil Ahmed Mr. Md. Shamim Sohag Mr. Md. Soleman Hossin Mr. Md. Suma Hasan Mr. Md. Zahidul Islam Munna Mr. Md. Zakir Hossain Mr. Md. Zamin Hossain Mr. Milon Paramanik Mr. Monirul Islam Tamim Mr. Mostafijur Rahman Rana Mr. Rasel Hossan Mr. Sheikh Eakub Ali Mr. Sk. Moin Hossen Mr. Subrata Chandra Paul Mr. Tanvir Ahammad Showmik Mr. Younus Mia Ms. Avesha Akter Ms. Sanjida Sultana Ripa

Ms. Shrabony Khan Jui

BANGLADESH HEALTH FACILITY SURVEY (BHFS) SUMMARY INDICATORS Appendix **B**

| AVAILABILITY OF BASIC CLIENT SERVICES % % % % Child curvative care (AI public facilities) 94 99 95 Child growth monitoring (AI public facilities) ¹ 80 90 93 Any modern method of family planing (AII public facilities) ¹ 80 90 83 Antenatial care service (AI public facilities) ¹ 80 99 100 Norral delivery (AII public facilities) 16 21 18 Antenatial care service (AI public facilities) 16 21 18 Auticability of all basic client services ² 1 30 100 All public facilities excluding CCs 20 41 30 NGO clinichospital 14 15 17 Private hospital 48 60 56 All public facilities excluding CCs 46 58 56 NGO clinichospital 49 60 51 Private hospital 08 04 01 Availability of electricity from national electricity grid line 41 45 A | Indicator | BHFS 2014 | BHFS 2017 | BHFS 2022 |
|---|--|------------------------|------------------|------------------------|
| Child curative care (All public facilities) 94 99 95 Child growth monitoring (All public facilities) 63 87 88 Child growth monitoring (All public facilities) ² 83 90 83 Any modern method of family planning (All public facilities) ² 83 99 100 Normal delivery (All public facilities) 16 21 18 Availability of all basic client services ³ 16 13 All public facilities excluding CCs 20 41 39 NGO clinic/hospital 14 15 17 7 7 Private hospital 08 04 01 14 15 All public facilities excluding CCs 46 58 66 NGO clinic/hospital 48 60 51 Private hospital 08 04 01 14 14 15 Olicin/hospital 7 57 85 51 14 14 Malbitis facilities excluding CCs 75 84 97 15 14 14 14 | AVAILABILITY OF BASIC CLIENT SERVICES | % | % | % |
| Child growth monitoring (All public facilities)638788Child vaccination (All public facilities)809093Any modern method of family public facilities) ¹² 8390100Any modern method of family public facilities) ¹² 8899100Any modern method of family public facilities) ¹² 8899100Any modern method of family public facilities)9899100Any modern method of family public facilities)181313Any public facilities excluding CCs204139100All public facilities excluding CCs204139100All public facilities excluding CCs46585656NGO clinichospital486051100Private hospital08746731All public facilities excluding CCs758688NGO clinichospital97969898Union-ever public facilities72829796District and upazita public facilities72817481NGO clinichospital969298979698Union-ever public facilities728339577586979698979698979698979698979698979698979698979698979698979698 | Child curative care (All public facilities) | 94 | 99 | 95 |
| Child vaccination (All public facilities)809083Any modern method of family planning (All public facilities)839083Any modern method of family planning (All public facilities)839083Normal delivery (All public facilities)162118All public facilities excluding CCs081613All public facilities excluding CCs080401All public facilities excluding normal delivery care7467All public facilities excluding CCs468866Co Colinch-hospital486061Private hospital080401All public facilities excluding CCs468866Co Colinch-hospital877467All public facilities excluding CCs758497Private hospital080401Avaite facilities excluding CCs758497Private hospital909097All public facilities728297Community Clinic214781Union-level public facilities769898Union-level public facilities769990Union-level public facilities779090Union-level public facilities769397Community Clinic981009790Union-level public facilities768286Community Clinic768286 <trr< td=""><td>Child growth monitoring (All public facilities)</td><td>63</td><td>87</td><td>88</td></trr<> | Child growth monitoring (All public facilities) | 63 | 87 | 88 |
| Any modern method of family planning (All public facilities)"839093Antenatal care service (All public facilities)9899100Normal delivery (All public facilities)161313Antenatal care services (All public facilities)081613All public facilities excluding CCs204139NGO clinich-sepital141517Private hospital080401All public facilities excluding normal delivery care486051All public facilities excluding CCs465856NGO clinich-sepital086051Private hospital086051All public facilities excluding CCs758497Private hospital979696All public facilities excluding CCs758497District and upazita public facilities722296Private hospital90979696Ordinch-sepital96929696Union-word public facilities72353957Community Clinic28323335District and upazita public facilities76799090Union-word public facilities76799090Union-word public facilities76828677Private hospital981009771Private hospital98909336 <tr< td=""><td>Child vaccination (All public facilities)¹</td><td>80</td><td>90</td><td>93</td></tr<> | Child vaccination (All public facilities) ¹ | 80 | 90 | 93 |
| Artenatal care service (All public facilities) 98 99 100 Normal delivery (All public facilities) 16 21 18 Availability of all basis client services? 30 13 31 All public facilities excluding CCs 20 41 33 NGO clinichospital 14 15 17 Private hospital 04 01 14 All basic client services excluding normal delivery care 45 74 67 All basic client services excluding CCs 46 80 051 OG clinichospital 48 00 51 Nicol facilities excluding CCs 75 85 74 67 All public facilities excluding CCs 75 84 97 56 All public facilities excluding CCs 72 82 97 57 Community Clinic 21 47 81 91 91 91 91 91 91 91 91 91 91 91 91 91 91 91 | Any modern method of family planning (All public facilities) ² | 83 | 90 | 83 |
| Normal delivery (All public facilities) 16 21 18 Availability of all basic client services ³ - - - All public facilities excluding CCs 20 41 39 NGO clinichospital 14 15 17 Private hospital 08 04 01 All public facilities excluding cCs 20 41 39 All public facilities excluding cCs 20 41 67 All public facilities excluding CCs 46 58 56 NGO clinichospital 08 04 01 Availability of electricity from national electricity grid line - - - All public facilities excluding CCs 75 84 97 District and upzila public facilities 72 82 97 Orinichospital 96 92 98 98 Private hospital 96 92 98 97 So clinichospital 96 92 98 97 NGO clinichospital 96 9 | Antenatal care service (All public facilities) | 98 | 99 | 100 |
| Availability of all basic client services ³ Interpret services of all public facilities excluding CCs 20 41 39 NGO clinic/hospital 14 15 17 Private hospital 08 04 01 All public facilities excluding CCs 08 04 01 All public facilities excluding normal delivery care H H 16 58 56 NGO clinic/hospital 48 60 51 57 68 56 NGO clinic/hospital 48 60 51 57 86 56 NGO clinic/hospital 75 87 86 56 | Normal delivery (All public facilities) | 16 | 21 | 18 |
| Ind Ind Ind Ind NGC clinic/nospital 14 15 17 Private hospital 08 04 01 All basic client services excluding normal delivery care I I 14 15 17 All basic client services excluding CCs 46 58 56 14 NGC clinic/nospital 48 60 51 17 NGC clinic/nospital 68 56 51 54 97 NGC clinic/nospital 68 56 51 54 97 All public facilities excluding CCs 75 85 51 53 All public facilities excluding CCs 75 84 97 57 85 Union-level public facilities 72 82 97 57 85 Community Clinic 21 47 81 56 56 56 57 57 85 57 57 85 57 57 85 57 57 57 57 | Availability of all basic client services ³ All public facilities | 08 | 16 | 13 |
| Ang Oc dinic/hospital 1 1 1 Private hospital 08 04 01 Private hospital 08 04 01 All public facilities 41 05 74 67 All public facilities 45 74 67 All public facilities 68 60 51 ORO clinic/hospital 68 60 51 Private hospital 08 04 01 Availability of electricity from national electricity grid line 75 84 97 All public facilities 72 82 97 District and upazila public facilities 72 82 97 Community Clinic 21 47 81 NGO clinic/hospital 90 90 97 Private hospital 90 90 97 All public facilities 74 40 48 All public facilities 76 79 90 Optionely public facilities 76 73 71< | All public facilities excluding CCs | 20 | 41 | 39 |
| Private hospital 0 0 0 All public facilities 45 74 67 All public facilities excluding CCs 46 58 56 NGO clinic/hospital 48 60 51 Private hospital 08 04 01 All public facilities excluding CCs 46 58 56 NGO clinic/hospital 08 04 01 Availability of electricity from national electricity grid line 77 85 85 All public facilities excluding CCs 75 84 97 District and upazila public facilities 72 82 97 Community Clinic 21 47 81 NGO clinic/hospital 96 92 98 Private hospital 90 97 96 98 Individi facilities excluding CCs 35 39 97 All public facilities 76 79 90 Union-level public facilities 76 79 90 Union-level public faci | NGO clinic/hospital | | 15 | 17 |
| All basic client services excluding normal delivery care H All basic client services excluding CCs 45 74 67 All public facilities excluding CCs 46 58 56 NGC clinic/hospital 08 04 01 All public facilities excluding CCs 66 57 84 97 All public facilities excluding CCs 75 84 97 96 98 All public facilities excluding CCs 75 84 97 96 98 Union-level public facilities 72 82 97 26 97 Community Clinic 21 47 81 96 92 98 Private hospital 90 90 90 97 40 48 All public facilities excluding CCs 35 39 57 10 10 40 45 NGO clinic/hospital 98 100 97 44 14 45 NGO clinic/hospital 98 100 97 57 53 | Private hospital | 08 | 04 | 01 |
| All public facilities excluding CCs 46 58 56 NGO clinic/hospital 48 60 51 Private hospital 08 04 01 Availability of electricity from national electricity grid line 37 57 85 All public facilities excluding CCs 75 84 97 District and upazila public facilities 72 82 97 Community Clinic 21 47 81 NGO clinic/hospital 90 90 97 Private hospital 90 90 97 Private hospital 90 90 97 All public facilities 74 40 48 All public facilities 76 79 90 All public facilities 76 79 90 Union-level public facilities 76 78 77 Private hospital 81 78 77 NGO clinic/hospital 81 78 77 NGO clinic/hospital 81 78 | All basic client services excluding normal delivery care | 45 | 74 | 67 |
| Air potic facilities excluding CCS 46 58 56 NGO clinic/hospital 08 04 01 Availability of electricity from national electricity grid line All public facilities 37 57 85 All public facilities 75 84 97 Optic facilities 97 96 98 Onion-level public facilities 97 96 98 Orion-level public facilities 72 82 97 Orion-level public facilities 72 83 Orion-level public facilities 73 57 84 NGO clinic/hospital 96 92 98 NGO clinic/hospital 96 92 98 | All public facilities | 45 | <i>(</i> 4 | 0/ |
| NGC clinic/nospital 48 60 51 Private hospital 04 01 Availability of electricity from national electricity grid line - - All public facilities excluding CCs 75 84 97 District and upazila public facilities 97 96 98 Union-level public facilities 72 82 97 Community Clinic 21 47 81 NGO clinic/hospital 90 90 97 Private hospital 90 90 97 Availability of regular electricity ⁴ 81 14 86 All public facilities excluding CCs 35 339 57 District and upazila public facilities 76 79 90 Union-level public facilities 76 79 90 Union-level public facilities 76 78 77 Private hospital 98 100 97 All public facilities excluding CCs 76 82 86 District and upazila public facilities </td <td>All public facilities excluding CCs</td> <td>46</td> <td>58</td> <td>56</td> | All public facilities excluding CCs | 46 | 58 | 56 |
| Private nospital 08 04 01 Availability of electricity from national electricity grid line 37 57 85 All public facilities excluding CCs 75 84 97 District and upazila public facilities 97 96 98 Union-level public facilities 72 82 97 Community Clinic 21 47 81 NGO clinic/hospital 96 92 98 Vivate hospital 96 92 98 NGO clinic/hospital 96 92 98 Naviability of regular electricity* 72 40 48 Availability of regular electricity* 73 96 90 Union-level public facilities excluding CCs 35 39 97 District and upazila public facilities 76 79 90 Union-level public facilities 81 78 77 NGO clinic/hospital 98 100 97 All public facilities excluding CCs 76 82 86 <t< td=""><td>NGO clinic/hospital</td><td>48</td><td>60</td><td>51</td></t<> | NGO clinic/hospital | 48 | 60 | 51 |
| Availability of electricity from national electricity grid line 37 57 85 All public facilities excluding CCs 75 84 97 District and upazila public facilities 97 96 98 Union-level public facilities 97 96 98 Community Clinic 21 47 81 NGO clinichospital 96 92 98 Private hospital 90 90 97 Aul public facilities 147 40 48 All public facilities excluding CCs 35 39 57 District and upazila public facilities 28 32 53 Community Clinic 98 100 97 NGO clinic/hospital 81 78 77 Private hospital 98 100 97 All public facilities 76 82 86 NGO clinic/hospital 81 78 75 All public facilities 74 81 85 Orisoti facilities excluding CCs < | Private hospital | 08 | 04 | 01 |
| All public facilities excluding CCs758497District and upazila public facilities979698Union-level public facilities728297Cormunity Clinic104781NGO clinic/hospital969298Private hospital909097Autility of regular electricity47048All public facilities excluding CCs353957District and upazila public facilities767990Union-level public facilities283253Ordmunity Clinic094145NGO clinic/hospital817877Private hospital817875All public facilities768286NGO clinic/hospital817875All public facilities768286District and upazila public facilities768286District and upazila public facilities768286District and upazila public facilities748185Community Clinic68777171NGO clinic/hospital88937575Nitch and upazila public facilities889375Nitch and upazila public facilities273.43.0All public facilities889375Nitch and upazila public facilities2.83.23.4Nitch and upazila public facilities2.83.2< | Availability of electricity from national electricity grid line All public facilities | 37 | 57 | 85 |
| District and upazila public facilities979698Union-level public facilities728297Community Clinic214781NGO clinic/hospital909097Private hospital909097Availability of regular electricity*7282All public facilities excluding CCs353957District and upazila public facilities767990Union-level public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital81707875NGO clinic/hospital707875All public facilities707875All public facilities748185Community Clinic687771NGO clinic/hospital869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities*203234All public facilities excluding CCs2.83.23.4Community Clinic2.83.23.4NGO clinic/hospital2.73.4303.4All public facilities excluding CCs2.83.23.4All public facilities excluding CCs2.8< | All public facilities excluding CCs | 75 | 84 | 97 |
| Union-level public facilities728297Community Clinic214781NGO clinic/hospital969298Private hospital909097Availability of regular electricity*174048All public facilities excluding CCs353957District and upazila public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital817877Private hospital817875So clinic/hospital707875All public facilities707875All public facilities707875Oltinic-hospital869094Union-level public facilities869375All public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities*2.73.430All public facilities excluding CCs2.83.23.4NGO clinic/hospital2.73.43031All public facilities excluding CCs2.83.23.4All public facilities excluding CCs2.83.23.4All public facilities excluding CCs2.83.23.4All public facilities excluding CCs2 | District and upazila public facilities | 97 | 96 | 98 |
| Community Clinic214781NGO clinic/hospital969298Private hospital909097Availability of regular electricity4174048All public facilities excluding CCs353957District and upazila public facilities767990Union-level public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital9810097Availability of client latrine5707875All public facilities excluding CCs768286District and upazila public facilities748185Community Clinic68909494Union-level public facilities748185Community Clinic68777171NGO clinic/hospital889375Private hospital94909393Union-level public facilities2.73.43.0Community Clinic6877713.0NGO clinic/hospital2.6Score (out of 6)Score (out of 6)All public facilities excluding CCs2.83.23.4NGO clinic/hospital2.73.43.03.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital2.73.43.03.0All public facilities excludi | Union-level public facilities | 72 | 82 | 97 |
| NGO clinic/hospital969298Private hospital909097Availability of regular electricity474048All public facilities excluding CCs353957District and upazila public facilities767990Outnon-level public facilities787771Optimization dupication817877Private hospital817877Private hospital707875All public facilities excluding CCs768286Octinic/hospital707875All public facilities excluding CCs768286District and upazila public facilities748185Original latrine586909393All public facilities excluding CCs687771NGO clinic/hospital748185Community Clinic889375Private hospital949093Private hospital949093Oscinic/hospital828375Inducting2.73.43.0All public facilities82.63.23.4Oscinic/hospital949093Private hospital949093All public facilities83.63.23.4All public facilities82.63.23.4All public facilities93.63.23.4All public | Community Clinic | 21 | 47 | 81 |
| Private hospital909097Availability of regular electricity*174048All public facilities excluding CCs353957District and upazila public facilities767990Union-level public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital9810097Private hospital707875All public facilities707875All public facilities768286District and upazila public facilities748185Community Clinic68909410Union-level public facilities748185Community Clinic68937571NGO clinic/hospital849375Private hospital94909332Private hospital94909330All public facilities2.73.43030All public facilities excluding CCs2.83.23.4NGO clinic/hospital6.47.04.330All public facilities excluding CCs2.83.23.4All public facilities excluding CCs2.83.23.4NGO clinic/hospital6.45.04.33.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital6.45.0< | NGO clinic/hospital | 96 | 92 | 98 |
| Availability of regular electricity ⁴ 17 40 48 All public facilities 35 39 57 District and upazila public facilities 76 79 90 Union-level public facilities 76 79 90 Community Clinic 09 41 45 NGO clinic/hospital 81 78 77 Private hospital 98 100 97 All public facilities 70 78 75 All public facilities excluding CCs 76 82 86 District and upazila public facilities 70 78 75 All public facilities 76 82 86 District and upazila public facilities 76 82 86 District and upazila public facilities 74 81 85 Community Clinic 88 93 75 NGO clinic/hospital 94 90 93 Private hospital 94 90 93 Aulpublic facilities 2.7 | Private hospital | 90 | 90 | 97 |
| All public facilities excluding CCs353957District and upazila public facilities767990Union-level public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital817877Private hospital707875All public facilities excluding CCs768286District and upazila public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Union-level public facilities748185Community Clinic687771NGO clinic/hospital949093Private hospital949093Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital5.45.65.2Private hospital5.45.65.2Private hospital5.45.65.2Private hospital5.45.65.2Private hospital5.45.65.2Private hospital5.45.65.2 | Availability of regular electricity ⁴ All public facilities | 17 | 40 | 48 |
| District and upazila public facilities767990Union-level public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital9810097Availability of client latrine ⁵ 707875All public facilities excluding CCs768286District and upazila public facilities748185Community Clinic74818566District and upazila public facilities748185Community Clinic68777171NGO clinic/hospital94909333Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)30All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3All public facilities excluding CCs2.83.23.4NGO clinic/hospital5.45.65.2Transport for emergencies ⁷ 5.45.65.2All public facilities030303 | All public facilities excluding CCs | 35 | 39 | 57 |
| Union-level public facilities283253Community Clinic094145NGO clinic/hospital817877Private hospital9810097Availability of client latrine ⁵ 707875All public facilities excluding CCs768286District and upazila public facilities869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities2.73.430All public facilities2.83.23.4NGO clinic/hospital4.75.04.3Private hospital4.75.04.3All public facilities5.45.65.2Transport for emergencies ⁷ 030303 | - District and upazila public facilities | 76 | 79 | 90 |
| Community Clinic 09 41 45 NGO clinic/hospital 81 78 77 Private hospital 98 100 97 Availability of client latrine ⁵ 70 78 75 All public facilities excluding CCs 76 82 86 District and upazila public facilities 86 90 94 Union-level public facilities 86 90 94 Oscore (out of not inc/hospital) 74 81 85 Community Clinic 68 77 71 NGO clinic/hospital 84 93 75 Private hospital 94 90 93 Aul public facilities 2.7 3.4 3.0 All public facilities excluding CCs 2.8 3.2 3.4 NGO clinic/hospital 4.7 5.0 4.3 NGO clinic/hospital 5.4 5.6 5.2 Private hospital 5.4 5.6 5.2 | Union-level public facilities | 28 | 32 | 53 |
| NGO clinic/hospital817877Private hospital9810097Availability of client latrine ⁵ 707875All public facilities excluding CCs768286District and upazila public facilities869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Auilability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Private hospital5.45.65.2All public facilities excluding CCs5.45.65.2All public facilities5.45.65.2Private hospital5.45.65.2All public facilities636363 | Community Clinic | 09 | 41 | 45 |
| Private hospital9810097Availability of client latrine5707875All public facilities excluding CCs768286District and upazila public facilities869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Analability of average basic amenities62.73.43.0All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Private hospital5.45.65.2All public facilities5.45.65.2Transport for emergencies7030303 | NGO clinic/hospital | 81 | 78 | 77 |
| Availability of client latrine5707875All public facilities707875All public facilities excluding CCs768286District and upazila public facilities869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities6Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies7030303 | Private hospital | 98 | 100 | 97 |
| All public facilitiesFoFoFoAll public facilities768286District and upazila public facilities869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies ⁷ All public facilities0303 | Availability of client latrine ⁵ | 70 | 78 | 75 |
| Implane technics statuting CosFo6260District and upazila public facilities869094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies ⁷ 030303 | All public facilities excluding CCs | 76 | 82 | 86 |
| District and updating public facilities505094Union-level public facilities748185Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies ⁷ All public facilities0303 | District and unazila public facilities | 86 | 02 QA | Q/ |
| Community ClinicF46160Community Clinic687771NGO clinic/hospital889375Private hospital949093Availability of average basic amenities ⁶ Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies ⁷ All public facilities0303 | Linion-level public facilities | 74 | 90 81 | 94 85 |
| NGO clinic/hospital0077NGO clinic/hospital889375Private hospital949093Availability of average basic amenities6Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies7030303 | Community Clinic | 68 | 77 | 71 |
| Private hospital005070Private hospital949093Availability of average basic amenities6Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies7All public facilities0303 | NGO clinic/hospital | 88 | 03 | 75 |
| Availability of average basic amenities6Score (out of 6)Score (out of 6)Score (out of 6)All public facilities2.73.43.0All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies7All public facilities0303 | Private hospital | 00 | 00 | 22 10 |
| Availability or average basic amenities*Score (out or b)Score | | 54 Score (cut of 6) | Score (out of 6) | 33 Score (out of 6) |
| All public facilities excluding CCs2.83.23.4NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies7All public facilities030303 | Availability of average basic amenities ⁶ All public facilities | 2.7 | 3.4 | 3.0 |
| NGO clinic/hospital4.75.04.3Private hospital5.45.65.2Transport for emergencies? All public facilities030303 | All public facilities excluding CCs | 2.8 | 3.2 | 3.4 |
| Private hospital5.45.65.2Transport for emergencies7All public facilities030303 | NGO clinic/hospital | 4.7 | 5.0 | 4.3 |
| Transport for emergencies7All public facilities03030303 | Private hospital | 5.4 | 5.6 | 5.2 |
| | Transport for emergencies ⁷ All public facilities | 03 | 03 | 03 |
| All public facilities excluding CCs 10 10 11 | All public facilities excluding CCs | 10 | 10 | 11 |
| District Hospital 92 98 97 | District Hospital | 92 | 98 | 97 |
| Upazila Health Complex 94 88 89 | Upazila Health Complex | 94 | 88 | 89 |

Continued...

| Indicator | BHFS 2014 | BHFS 2017 | BHFS 2022 |
|--|-------------------|-------------------|-------------------|
| NGO clinic/hospital | 20 | 23 | 25 |
| Private hospital | 63 | 65 | 32 |
| Average basic diagnostic testing capacity ⁸ | Score (out of 5) | Score (out of 5) | Score (out of 5) |
| All public facilities | 0.4 | 1.2 | 1.0 |
| All public facilities excluding CCs | 0.6 | 1.0 | 1.8 |
| NGO clinic/hospital | 3.3 | 4.1 | 4.4 |
| Private hospital | 3.6 | 4.1 | 3.7 |
| Availability of functional x-ray machines District Hospital | 73 | 79 | 87 |
| Upazila Health Complex | 22 | 18 | 47 |
| NGO clinic/hospital | 02 | 02 | 00 |
| Private hospital | 62 | 66 | 54 |
| Availability of ultrasound machine | | | |
| District Hospital | 63 | 74 | 79 |
| Upazila Health Complex | 05 | 16 | 49 |
| NGO clinic/hospital | 16 | 27 | 42 |
| Private hospital | 76 | 69 | 75 |
| Availability of essential medicines ⁹ All public facilities | 69 | 60 | 72 |
| All public facilities excluding CCs | 70 | 45 | 54 |
| NGO clinic/hospital | 60 | 47 | 11 |
| Private hospital | 45 | 40 | 10 |
| % of service provider positions functionally vacant in | | | |
| Physicians in district and upazila-level public facilities | 38 | 40 | 26 |
| Nurse/midwife in district and upazila-level public facilities | 19 | 18 | 11 |
| At least one staff trained in IMPAC ¹⁰ in the last 24 months | | | |
| % of public health facilities | 07 | 08 | 04 |
| CHILD HEALTH SERVICES | % | % | % |
| Availability of child curative care services | | | |
| All public facilities | 94 | 99 | 95 |
| All public facilities excluding CCs | 96 | 98 | 96 |
| NGO clinic/hospital | 83 | 94 | 88 |
| Private hospital | 68 | 90 | 57 |
| Availability of child growth monitoring services | 62 | 07 | 00 |
| All public facilities | 68 | 07 70 | 00 |
| All public facilities excluding CCS | 69 | 80 | 82 |
| Rivato bospital | 20 | 45 | 23 |
| Availability of child vaccination services | 20 | -10 | 20 |
| Availability of clinic vaccination services | 80 | 90 | 93 |
| All public facilities excluding CCs | 68 | 77 | 81 |
| NGO clinic/hospital | 71 | 69 | 72 |
| Private hospital | 16 | 07 | 03 |
| Average readiness score ¹¹ of facilities to provide child curative care | Score (out of 10) | Score (out of 10) | Score (out of 10) |
| All public facilities | 7.0 | 7.0 | 6.8 |
| All public facilities excluding CCs | 6.5 | 5.8 | 5.6 |
| NGO clinic/hospital | 7.4 | 7.3 | 6.0 |
| Private hospital | 6.1 | 5.6 | 4.3 |
| FACILLITIES PROVIDING FAMILY PLANNING (FP) SERVICES | % | % | % |
| Any modern FP methods ¹² | | | |
| All public facilities | 79 | 88 | 77 |
| All public facilities excluding CCs | 83 | 88 | 80 |
| NGO clinic/hospital | 84 | 83 | 60 |
| Private hospital | 14 | 25 | 10 |
| Long-acting, reversible contraceptives or permanent methods ¹³ | 20 | 24 | 40 |
| All public facilities | 30 | 24 | 19 |

Continued...

| Indicator | BHFS 2014 | BHFS 2017 | BHFS 2022 |
|---|-------------------|-------------------|-------------------|
| All public facilities excluding CCs | 63 | 74 | 68 |
| NGO clinic/hospital | 66 | 69 | 45 |
| Private hospital | 14 | 19 | 10 |
| Male/female sterilization services ¹⁴ | | | |
| All public facilities | 04 | 04 | 04 |
| All public facilities excluding CCs | 10 | 13 | 12 |
| NGO clinic/hospital | 20 | 11 | 20 |
| Private hospital | 06 | 18 | 10 |
| Average readiness score ¹⁵ to provide family planning (FP) | Score (out of 8) | Score (out of 8) | Score (out of 8) |
| All public facilities | 4.9 | 4.5 | 4.4 |
| All public facilities excluding CCs | 5.9 | 6.0 | 6.1 |
| NGO clinic/hospital | 6.4 | 6.5 | 5.4 |
| Private hospital | 3.3 | 2.2 | 2.2 |
| ANTENATAL CARE SERVICES | % | % | % |
| Availability of antenatal care services | | | |
| All public facilities | 98 | 99 | 100 |
| All public facilities excluding CCs | 96 | 96 | 99 |
| NGO clinic/hospital | 98 | 100 | 100 |
| Private hospital | 79 | 95 | 72 |
| Average readiness score ¹⁶ of health facilities to provide ANC | Score (out of 6) | Score (out of 6) | Score (out of 6) |
| All public facilities | 3.0 | 3.1 | 3.0 |
| All public facilities excluding CCs | 3.2 | 3.4 | 3.8 |
| NGO clinic/hospital | 4.8 | 4.9 | 4.8 |
| Private hospital | 3.9 | 3.7 | 3.4 |
| DELIVERY AND NEWBORN CARE | % | % | % |
| Availability of normal delivery services | ,0 | ,0 | ,,, |
| All public facilities | 16 | 21 | 18 |
| All public facilities excluding CCs | 35 | 57 | 55 |
| NGO clinic/hospital | 31 | 32 | 34 |
| Private hospital | 93 | 95 | 89 |
| Availability of cesarean delivery services | | | |
| All public facilities | 01 | 01 | 02 |
| All public facilities excluding CCs | 04 | 04 | 07 |
| NGO clinic/hospital | 15 | 11 | 21 |
| Private hospital | 95 | 97 | 93 |
| Average readiness score ¹⁷ to provide normal delivery services | Score (out of 13) | Score (out of 13) | Score (out of 13) |
| All public facilities | 4.7 | 4.7 | 5.9 |
| All public facilities excluding CCs | 5.3 | 5.1 | 6.6 |
| NGO clinic/hospital | 8.4 | 9.2 | 10.3 |
| Private hospital | 8.3 | 8.0 | 8.4 |
| NEWBORN SIGNAL FUNCTIONS | % | % | % |
| Primary newborn signal functions ¹⁸ | | | |
| All public facilities | - | - | 43 |
| All public facilities excluding CCs | - | - | 50 |
| NGO clinic/hospital | - | - | 76 |
| Private hospital | - | - | 51 |
| Basic newborn signal functions ¹⁹ | | | |
| All public facilities | - | - | 08 |
| NGO clinic/hospital | - | - | 33 |
| Private hospital | - | - | 22 |
| Comprehensive newborn signal functions ²⁰ | | | |
| All public facilities | - | - | 11 |
| NGO clinic/hospital | - | - | 09 |
| Private hospital | - | - | 07 |
| Advanced newborn signal functions ²¹ | | | |

Continued...
| Indicator | BHFS 2014 | BHFS 2017 | BHFS 2022 |
|-----------------------|-----------|-----------|-----------|
| All public facilities | - | - | 05 |
| NGO clinic/hospital | - | - | 00 |
| Private hospital | - | - | 04 |

¹ Facility routinely provides DPT/pentavalent, polio, and measles vaccinations offered from the facility, excluding any outreach services ² Facility provides, prescribes, or counsels clients on any of the following: contraceptive pills (combined or progestin-only), injectables (progestin-only), one-rod implant, two-rod implant (zadell), IUCDs, male condoms, female sterilization (tubal ligation) or male sterilization (vasectomy), and emergency contraceptive pills.

³ Basic client services include outpatient curative care for sick children, child growth monitoring, facility-based child vaccination services, any modern methods of family planning, antenatal care, and normal delivery.

⁴ Facility is connected to a central power grid and there was no interruption in power supply lasting for more than 2 hours at a time during normal working hours in the 7 days before the survey, or facility has a functioning generator on the day of the survey, or else facility has back-up solar power.

⁵ Facility has a functioning flush or pour-flush toilet, a ventilated improved pit latrine, or a composting toilet.

⁶ Average number of basic amenities include regular electricity, an improved water source, visual and auditory privacy, a client latrine, communication equipment, and a computer with Internet.

⁷ Facility has a functioning ambulance or other vehicle for emergency transport that is stationed at the facility and operates from the facility.
⁸ The capacity to conduct average number of basic diagnostic tests (hemoglobin, blood glucose, urine protein, urine glucose, and urine pregnancy test).

pregnancy test). ⁹ Defined as availability of at least six of eight essential medicines of a DDS kit: amoxicillin tablet/capsule, amoxicillin syrup, cotrimoxazole, paracetamol tablet, paracetamol syrup, tetracycline eye ointment, iron tablet, and vitamin A capsule.

¹⁰ IMPAC stands for integrated management of pregnancy and childbirth.

¹¹ Average readiness score is the average number of 10 items (IMCI guideline, staff trained in IMCI at any time, child weighing scale, thermometer, growth chart, zinc tablets/syrup, ORS, amoxicillin syrup/suspension/dispersible, paracetamol syrup/suspension, and mebendazole/albendazole) available that are necessary for providing outpatient curative care for sick children.

¹² Facility provides any of the following: contraceptive pills (combined or progestin-only), injectables (progestin-only), one-rod implant, tworod implant (zadell), IUCDs, male condoms, female sterilization (tubal ligation), or male sterilization (vasectomy), and emergency contraceptive pills.

¹³ Facility provides any of the following long-term and permanent FP methods: one-rod implant, two-rod implant, IUCDs, female sterilization (tubal ligation), or male sterilization (vasectomy).

¹⁴ Perform male or female sterilization in the facility.

¹⁵ Average readiness score is the average number of 8 items (guidelines on FP, staff trained in FP any time, blood pressure apparatus, combined or progestin-only oral pills, progestin-only injectable, male condom, intrauterine contraceptive device, and implant) available for providing FP services.

¹⁶ Average readiness score is the average number of 6 items (guidelines on ANC, staff trained for ANC any time, blood pressure apparatus, hemoglobin testing capacity, urine protein testing capacity, and iron or folic acid tablets) available for providing antenatal care services.
¹⁷ Average readiness score is the average number of 13 items (guidelines on BEmOC or CEmOC, staff trained in delivery care any time, examination light, delivery pack, suction apparatus, neonatal bag and mask, partograph, gloves, oxytocin, injectable antibiotic, magnesium subbate skin disinfectant, and intravenue fully in the interview pack.

sulphate, skin disinfectant, and intravenous fluid with infusion set) available for providing normal delivery services. ¹⁸ Primary newborn signal functions (all 9 NSFs): Iron and folic acid supplementation in pregnant women, hand wash each time before touching the baby, drying of the baby immediately after birth, delayed umbilical cord clamping, clean cord cutting sterile blade, single application of 7.5% CHX, neonatal resuscitation, perform skin-to-skin care immediately after birth, and early initiation of breast feeding. ¹⁹ Basic newborn signal functions (all 13 NSFs): Basic NSF practices include primary newborn signal functions and Immunize patient with

¹⁹ Basic newborn signal functions (all 13 NSFs): Basic NSF practices include primary newborn signal functions and Immunize patient with tetanus toxoid, administer oral antibiotic, administer intramuscular antibiotic, and suction of airway of newborn. Community clinics don't have the provision to perform all basic newborn signal functions

²⁰ Comprehensive newborn signal function (all 18 NSFs): Comprehensive NSF practices include basic newborn signal functions and administer intravenous antibiotic, administer antibiotic for premature rupture of membrane, oxygen therapy for management of newborn infection and respiratory problem, KMC for pre-term of LBW newborn, phototherapy for term or pre-term with hyperbilirubinemia. Community clinics and union level facilities are not designated to perform all comprehensive newborn signal functions.

clinics and union level facilities are not designated to perform all comprehensive newborn signal functions. ²¹ Advanced newborn signal function (all 20 NSFs): Advanced NSF practices include comprehensive newborn signal functions and incubator support for sick term baby, pre-term baby or if the baby receiving KMC becomes sick, and corticosteroids for pre-term labor. Community clinics and union level facilities are not designated to perform all advanced newborn signal functions.



2022 BANGLADESH HEALTH FACILITY SURVEY

INVENTORY QUESTIONNAIRE

| | | F/ | ACILITY IDENTIFIC | ATION | |
|----------|---|--|-------------------|-------------|-------------------|
| 001 | NAME OF FACI | LITY | | | |
| 001A | ADDRESS: | | | | |
| 002 | DIVISION: | | | | |
| 003 | DISTRICT (ZILA |): | | | |
| 004 | UPAZILA/THAN | IA: | | | |
| 005 | UNION/WARD | : | | | |
| 006 | FACILITY NUM | BER: | | | |
| 007 | 007 TYPE OF FACILITY COUNTRY SPECIFIC: DISTRICT HOSPITAL (DH) 01 UPAZILA HEALTH COMPLEX (UHC) 02 MATERNAL AND CHILD WELFARE CENTER (MCWC) 03 UNION HEALTH AND FAMILY WELFARE CENTER (UnHFWC) 04 UNION SUBCENTER (USC) /RURAL DISPENSARY. 06 COMMUNITY CLINIC. 08 NGO CLINIC/HOSPITAL 09 | | | | |
| 008 | OWNERSHIP (MA GOVERNMENT/PI LOCAL GOVERNM NGO (NAME) PRIVATE FOR PRO | NAGING AUTHORITY) JBLIC (MOHFW) IENT | | | |
| 009 | URBAN/RURAL URBAN RURAL | | | | |
| 010 | YES | | | | |
| | | | INTERVIEWER V | ISITS | 2 |
| | | 1 | 2 | 3 | FINAL VISIT |
| DATE | | | | | DAY |
| | | | | | MONTH |
| INTERVIE | WER'S NAME | | | | YEAR |
| | | | - | - | INTV. CODE NUMBER |
| RESULT* | | | | - <u></u> | RESULT |
| *RESUL | T CODES: | | | | |
| 1 | = FACILITY COM | PLETED | | | |
| 2 | 2 = FACILITY RESPONDENTS NOT AVAILABLE | | | | |
| 3 | = POSTPONED / | PARTIALLY COMPLET | TED | | |
| 4 | = FACILITY REFUS | SED | | | |
| 5 | = FACILITY CLOSI | ED / NOT YET FUNCT | IONAL | | |
| 6 | = OTHER | | | _ | |
| | (SPECIFY) | | | | |

TOTAL NUMBER OF PROVIDER INTERVIEWED

| TOTAL | NUMBER | OF | PROVIDERS | INTERVIEWED | |
|---------|--------------------|--------------|----------------------|------------------------|---------------------------|
| | | | | | |
| | | FA | CILITY GEOGRAPHIC | | |
| | | | | | |
| SET DE | FAULT SETTINGS FO | R GPS UNIT | | | |
| • | SET COORDINATE | SYSTEM TO L | ATITUDE / LONGITUDE | <u>.</u> | |
| • | SET COORDINATE | FORMAT TO I | DECIMAL DEGREE | | |
| • | SET DATUM TO W | GS84 | | | |
| STAND | IN A LOCATION AT | | E OF THE FACILITY WI | TH PLAIN VIEW OF THE S | KY TION!! |
| 1. | | NE ON AND V | AIT UNTIL SATELITE P | AGE CHANGES TO "POSI | HON" |
| 2. | WAILS MINULES | | | | |
| 3. | | | | D " | |
| 4. | | | | n | |
| 5. | | " AND PRESS | 'ENTER" | | |
| 0. 7 | PAGE TO MAIN M | | HT "WAYPOINT LIST" | AND PRESS "ENTER" | |
| 8 | | WAYPOINT | | | |
| 9. | COPY INFORMATI | ON FROM WA | YPOINT LIST PAGE | | |
| 10 |). WRITE ELEVATION | I [ALTITUDE] | | | |
| BE SUF | RE TO COPY THE WA | YPOINT NAM | E FROM THE WAYPOIN | NT LIST PAGE TO VERIFY | THAT YOU ARE ENTERING THE |
| CORRE | CT WAYPOINT INFO | RMATION ON | THE DATA FORM | | |
| | | | | | |
| 008 | WAYPOINT NAME) | NAME (F/ | ACILITY FACILITY | NUMBER | |
| | | | | | |
| 009 | LATITUDE | | N/S | a | |
| | | | DEGREES/D | DECIMb | |

E/Wa

DEGREES/DECIM..b

_°c

010

LONGITUDE

CONSENT

FIND THE MANAGER, THE PERSON IN-CHARGE OF THE FACILITY, OR THE MOST SENIOR HEALTH WORKER RESPONSIBLE FOR CLIENT SERVICES WHO IS PRESENT AT THE FACILITY. READ THE FOLLOWING GREETING:

Good day! My name is ______. We are here on behalf of the National Institute of Population Research and Training (NIPORT), Medical Education & Family Welfare Division (ME&FWD) and Ministry of Health and Family Welfare (MOHFW) conducting a survey of health facilities to assist the Government of knowing more about health services in BANGLADESH.

Now I will read a statement explaining the study.

Your facility was selected to participate in this study. We will be asking you questions about various health services. Information collected about your facility during this study may be used by NIPORT, the MOHFW, organizations supporting services in your facility, and researchers, for planning service improvement or for conducting further studies of health services.

Neither your name nor the names of any other health workers who participate in this study will be included in the dataset or in any report; however, there is a small chance that any of these respondents may be identified later. Still, we are asking for your help in order to collect this information.

You may refuse to answer any question or choose to stop the interview at any time. However, we hope you will answer the questions, which will benefit the services you provide and the nation.

If there are questions for which someone else is the most appropriate person to provide the information, we would appreciate if you introduce us to that person to help us collect that information.

Γ

Hours

Minutes

At this point, do you have any questions about the study? Do I have your agreement to proceed?

| | INTERVIEWER'S SIGNATURE INDICATING CONSENT OBTAINED | | Day | Month | Year | | |
|-----|---|-----------|-----|-------|----------|--------|--|
| 100 | May I begin the interview? | Yes No | | | 1 2 — | → Stop | |
| 101 | INTERVIEW START TIME | | | | | | |

EXPLAIN TO THE RESPONDENT AT THE START OF THIS INTERVIEW THAT THERE ARE QUESTIONS ON MANAGEMENT MEETINGS AND QUALITY ASSURANCE ACTIVITIES THAT REQUIRE LOOKING AT RECORDS OF THOSE MEETINGS AND ACTIVITIES. IT WILL THEREFORE BE HELPFUL IF RECORDS PERTAINING TO MANAGEMENT MEETINGS AND QUALITY ASSURANCE ACTIVITIES ARE GATHERED, IF THEY ARE NOT READILY AVAILABLE AT THE LOCATION WHERE YOU ARE CONDUCTING THE INTERVIEW.

THANK THE RESPONDENT AT THE END OF EACH SECTION OR SUBSECTION BEFORE PROCEDING TO THE NEXT DATA COLLECTION POINT

MODULE 1: GENERAL INFORMATION AND SERVICE AVAILABILITY

SECTION 1: GENERAL SERVICE AVAILABILITY AND INPATIENT SERVICES SERVICE AVAILABILITY

| 102 | Does this facility offer any of the following client services? In other words, is there any location in this facility where clients can receive any of the following services: | YES | NO | DONE |
|-----------------|--|-----|----|------|
| 01 | Child vaccination services. | 1 | 2 | |
| 02 | Growth monitoring services. | 1 | 2 | |
| 03 | Curative care services for children under age 5. | 1 | 2 | |
| 04 | Any family planning services including modern methods, fertility awareness methods (natural family planning), male or female surgical sterilization | 1 | 2 | |
| 05 | Antenatal care (ANC) services | 1 | 2 | |
| 07 | Normal delivery | 1 | 2 | |
| 10 | Diagnosis, treatment prescription or treatment follow-up for TB | 1 | 2 | |
| 14 | Diagnosis or management of non-communicable diseases, specifically diabetes, cardiovascular diseases and chronic respiratory conditions, cervical cancer in adults. | 1 | 2 | |
| 16 | Cesarean delivery (Cesarean section) | 1 | 2 | |
| 17 | Laboratory diagnostic services, including any rapid diagnostic testing. | 1 | 2 | |
| 18 | Blood grouping and typing services | 1 | 2 | |
| 19 | Blood transfusion services | 1 | 2 | |
| 20 | Postnatal care (PNC) services | 1 | 2 | |
| 21 | Postpartum family planning (PPFP) services | 1 | 2 | |
| 22 | Adolescent health services | 1 | 2 | |
| 23 | Nutrition services | 1 | 2 | |
| 24 [‡] | Care and/or referral services for victims of gender-based violence (GBV) | 1 | 2 | |
| 25 [‡] | Post abortion care (PAC) services | 1 | 2 | |
| 26 [‡] | Services for disability | 1 | 2 | |
| 27 [‡] | COVID-19 services | 1 | 2 | |

INPATIENT (INDOOR) SERVICES

| 110 | Does this facility routinely provide in-patient care? | YES 1 112 | |
|-----|---|-----------------|--|
| | | NO2 | |
| 111 | Does this facility have beds for overnight | YES 1 | |
| | observation? | NO2 →200 | |
| 112 | Excluding any delivery and/or maternity beds, how | # OF OVERNIGHT/ | |
| | many (overnight) or (in-patient) beds in total does | INPATIENT BEDS. | |
| | this facility have, both for adults and children? | DON'T KNOW 998 | |
| 113 | How many beds in total does this facility have | | |
| | according to GOB circular/license? | # OF BEDS | |
| | | DON'T KNOW 998 | |
| 114 | Current number of beds with budget allocation? | | |
| | | # OF BEDS | |
| | | DON'T KNOW 998 | |

Section 2: GENERAL FILTER QUESTIONS

PROCESSING OF INSTRUMENTS

| 200 | I have a few questions about how surgical instruments, such as speculums, forceps, and other metal equipment are processed for re-use in this facility. Are instruments that are used in areas processed (i.e., sterilized or high-level disinfected) for re-use? | YES1 NO2 - | >210 |
|-----|---|-----------------------|------|
| 201 | Is the final processing done in this facility, outside this facility, or both? | Only in this Facility | |

STORAGE OF MEDICINES

| 210 | Does this facility store or keep any medicines (including antibiotics, analgesics), vaccines or contraceptive commodities in the facility? I am referring to medicines/commodities that are meant to be given to clients when a provider prescribes them from outpatient departments? PROBE | YES1 Facilities Stocks No Medicine2— | → 300 |
|-----|---|---|-------|
| 211 | CHECK Q102.04 | | |
| | Family Planning services Available | No Family Planning Services | 213 |
| 212 | Are contraceptive commodities generally stored in the family planning service area, or are they stored in a common area with other medicines? | Stored in FP Service area | |
| 213 | CHECK Q102.10 | | |
| | TUBERCULOSIS SERVICES | NO TUBERCULOSIS | 300 |
| 214 | Are medicines for the treatment of TB generally stored in the TB service area or are they stored in a common area with other medicines? | Stored in TB Service area | |

MODULE 2: GENERAL SERVICE READINESS

SECTION 3: 24-HOUR STAFF COVERAGE – INFRASTRUCTURE

EXTERNAL SUPERVISION - USER FEES

24 HOURS STAFF COVERAGE

| 300 | Is there a health care worker present at the facility at all times, or officially on call for the facility at all times (24 hours a day) for emergencies? Specifically, I am referring to medical specialists, medical officers, nurses and paramedics. | YES, 24-Hour Staff 1 No, 24-Hour Staff 2 | |
|-----|---|---|--|
|-----|---|---|--|

COMMUNICATION

| 310 | Does this facility have an official land line or cellular phone that is available to call outside at all times outpatient services are offered? CLARIFY THAT IF FACILITY OFFERS 24-HOUR EMERGENCY SERVICES, THEN THIS REFERS TO 24-HOUR AVAILABILITY. | YES1 NO2 | ▶ 313 |
|-------------------|--|---------------|-------|
| 312 | Is it functioning? ACCEPT REPORTED RESPONSE | YES1 NO2 | |
| 313 | Does this facility have <u>a private cellular phone</u> that is reimbursed by the facility? | YES1 NO2 — | ▶ 319 |
| 315 | Is it functioning? ACCEPT REPORTED RESPONSE | Yes1 No2 | |
| 319 | Does this facility have a computer (Desktop/Laptop)/or a Tablet for outpatient services? | YES1 NO2 — | ▶ 322 |
| 321 | Is it functioning? ACCEPT REPORTED RESPONSE | Yes1 No2 | |
| 321a [‡] | Does this facility use service provision software? | Yes1 No2 | |
| 322 | Is there access to email or internet via computer and/or mobile phone within the facility? ACCEPT REPORTED RESPONSE. | YES1 NO2 — | ▶ 330 |
| 323 | Is the email or internet routinely available for at least 2 hours on days that services are offered? ACCEPT REPORTED RESPONSE. | YES1 NO2 | |

SOURCE OF WATER

| 330 | What is the most commonly used source of water for the facility at this time? OBSERVE THAT WATER IS AVAILABLE FROM SOURCE OR IN THE FACILITY ON THE DAY OF THE VISIT. E.G., CHECK THAT THE PIPE IS FUNCTIONING. | Piped into Facility 01 Piped onto Facility grounds 02 Public Tap/Stand Pipe 03 Tube well/Borehole 04 Protected Dug Well 05 Unprotected Dug Well 06 Protected Spring 07 Unprotected Spring 08 Rain Water 09 Bottled Water 10 CART W/SMALL TANK/DRUM 11 TANKER TRUCK 12 SURFACE WATER 13 (RIVER/DAM/LAKE/POND) 0 Other 96 (SPECIFY) 98 Don't Know 98 No water source 00 |
|-----|--|---|
| 331 | Is water outlet from this source available onsite, within 500 meters of the facility, or beyond 500M of facility? REPORTED RESPONSE IS ACCEPTABLE | ONSITE |

POWER SUPPLY

| 340 | Is this facility connected to the national electricity grid? (including palli biddut) | YES | ▶ 342 |
|-----|--|---|-------|
| 341 | During the past 7 days, was electricity (excluding any back-up generator) available during the times when the facility was open for services, or was it ever interrupted for more than 2 hours at a time? CONSIDER ELECTRICITY TO BE ALWAYS AVAILABLE IF INTERUPTED FOR LESS THAN 2 HOURS AT A TIME. | ALWAYS AVAILABLE | |
| 342 | Does this facility have other sources of electricity such as a generator or solar system? | Yes 1 No other source 2 | ▶ 350 |
| 343 | What other sources of electricity does this facility have? PROBE FOR ANSWERS AND CIRCLE ALL THAT APPLY | FUEL-OPERATED GENERATOR A BATTERY-OPERATED GENERATOR B SOLAR SYSTEM C OTHER(SPECIFY) | |

| 344 | CHECK Q343 GENERATOR OR SOLAR USED (EITHER 'A' OR 'B' OR 'C' CIRCLED) | GENERATOR OR SOLAR NOT USED (NEITHER 'A' OR 'B' OR 'C' CIRCLED) | 350 |
|-----|--|---|-----|
| 345 | Is the generator or solar system functional? ACCEPT REPORTED RESPONSE FROM KNOWLEDGEABLE RESPONDENT. | YES, GENERATOR IS FUNCTIONAL | |

EXTERNAL SUPERVISION

| 350 | Does this facility receive any external supervision from the any upper level facility/higher authority? | ervision from the any upper level NO Iity/higher authority? | | | ▶360 |
|------|---|--|----------|---------------|------|
| 351 | When was the last time a supervisor from any upper level office came here on a supervisory visit? Was it within the past 6 months or more than 6 months ago? | WITHIN THE PAST 6 MONTH MORE THAN 6 MONTHS AG | HS iO | 1 2— | →360 |
| 351A | During the past 6 months, how many supervisory visits has this facility received from an upper level office/higher authority? | # OF SUPERVISORY VISITS DON'T KNOW | | | |
| 352 | The last time during the past 6 months that a s facility or from relevant department visited, of following: | YES | NO | DON'T KNOW | |
| 01 | Use a checklist to assess the quality of availabl | e health services data? | 1 | 2 | 8 |
| 02 | Discuss performance of the facility based o data? | 1 | 2 | 8 | |
| 352A | Does this facility maintain records of written comments made by supervisors from outside the facility when they make their supervisory visits? | n YES, RECORDS MAINTAINED e NO, RECORDS NOT MAINTAINED y | | | |

USER FEES

| 360 | Does this facility have any routine user-fees or charges for client services, including charges for health cards and for client registration? These could be routine fees for some services or for medicines | YES NO | 1 2— | ▶400 |
|-----|--|---|---------|-------|
| 361 | Does the facility charge a fixed fee that covers all services that a client receives, or are there separate fees for different components of the services provided by the facility? PROBE. | FIXED FEE COVERING ALL SERVIO NO, CHARGE FEE FOR SEPARATE ITEMS | ▶ 363 | |
| 362 | Does this facility have a fee for the following items: READ OUT EACH RESPONSE CATEGORY AND CIRCLE APPROPRIATELY | YES | NO | |
| 01 | CLIENT HEALTH CARD | 1 | 2 | |
| 02 | REGISTRATION | 1 | 2 | |
| 03 | CONSULTATION | 1 | 2 | |
| 04 | MEDICINES | 1 | 2 | |
| 05 | VACCINES | 1 | 2 | |
| 06 | CONTRACEPTIVE COMMODITIES | 1 | 2 | |
| 07 | NORMAL DELIVERIES | 1 | 2 | |
| 08 | SYRINGES AND NEEDLES | 1 | 2 | |
| 09 | CESAREAN SECTION | 1 | 2 | |
| 17* | TICKET AND ADMISSION FEES | 1 | 2 | |
| 18* | BED/CABIN CHARGE | 1 | 2 | |
| 19* | CLIENT VOLUNTARY CONTRIBUTION | 1 | 2 | |
| 13 | LABORATORY TESTS | 1 | 2 | |
| 363 | Are the official fees posted or displayed so that the client can easily see them? | YES NO POSTED FEES | 1 2— | → 400 |
| 364 | May I see the posted fees? | OBSERVED, ALL FEES POSTED | 1 | |
| | REVIEW THE POSTED FEES AGAINST THE LIST OF ITEMS IN Q362 TO DETERMINE IF ALL FEES ARE POSTED | OBSERVED, SOME BUT NOT ALL | FEES2 | |

SECTION 4: STAFFING - MANAGEMENT -

QUALITY ASSURANCE - TRANSPORT - HMIS

| 400 | Please tell me: | | | | | | |
|-----|--|--------------------------|-------------------|---------------------------|-------------------|------------------|--|
| a) | How many staff in each of the following qualification / occupational categories are sanctioned (expected) to work in this facility. | | | | | | |
| b1) | For each qualification / occupational category, how many are currently posted based on sanctioned and write down the number. Remember to count health care providers who are (attached out) in other facilities. However, if a health care provider is (deputed out) from the facility then we have to minus that number from (posted). If some healthcare provider come from other facility (deputed in) or (attached in) then we will not be recognized them as posted | | | | | | |
| b2) | For each qualification / occupational cate | egory, how man | y are attached/ | deputed in and wo | orking in the fac | ility. | |
| D3) | For each qualification / occupational cate | egory, how man | y are currently d | eputed out | | | |
| C) | For each qualification / occupational cat many are part-time | egory, among th | nose currently p | oosted/attached a | nd working in t | he facility, how | |
| | Qualification/ Occupational Category | (a) | (b1) | (b2) | (b3) | (C) | |
| | | SANCTIONED (EXPECTED) | POSTED | ATTACHMENT/ DEPUTED-IN | DEPUTED-OUT | PART TIME | |
| 01 | SUPERINTENDANT | | | | | | |
| 02 | DIRECTOR/MANAGER/ COORDINATOR | | | | | | |
| 03 | LIH&EPO | | | | | | |
| 04 | SENIOR CONSULTANT(MEDICINE) | | | | | | |
| 05 | SENIOR CONSULTANT(SURGERY) | | | | | | |
| 06 | SENIOR CONSULTANT(OBS.&GYN.) | | | | | | |
| 07 | SENIORCONSULTANT (PEDIATRICS) | | | | | | |
| 08 | SENIOR CONSULTANT (ORTHOPEDIC SURGURY) | | | | | | |
| 09 | SENIOR CONSULTANT (EYE) | | | | | | |
| 10 | SENIOR CONSULTANT (ANESTHESIA) | | | | | | |
| 11 | CONSULTANT (RADIOLOGY AND IMAGING) | | | | | | |
| 12 | CONSULTANT (PATHOLOGIST) | | | | | | |
| 13 | SENIOR CONSULTANT (ENT) | | | | | | |
| 14 | SENIOR CONSULTANT (SKIN & VD) | | | | | | |
| 15 | SENIOR CONSULTANT (CARDIOLOGY) | | | | | | |
| 16 | JUNIOR CONSULTANT(MEDICINE) | | | | | | |
| 17 | JUNIOR CONSULTANT(SURGERY) | | | | | | |
| 18 | JUNIOR CONSULTANT(OBS.&GYN.) | | | | | | |
| 19 | JUNIOR CONSULTANT (PEDIATRICS) | | | | | | |
| 20 | JUNIOR CONSULTANT (ORTHOPEDIC SURGURY) | | | | | | |
| 21 | JUNIOR CONSULTANT (EYE) | | | | | | |

| | Qualification/ Occupational Category | (a) | (b1) | (b2) | (b3) | (C) |
|----|--|------------|--------|-------------|-------------|-----------|
| | | SANCTIONED | POSTED | ATTACHMENT/ | DEPUTED-OUT | PART TIME |
| | | | | | | |
| 22 | | | | | | |
| 22 | JUNIOR CONSULTANT (ANESTHESIA) | | | | | |
| 25 | JUNIOR CONSULTANT (RADIOLOGY) | | | | | |
| 24 | JUNIOR CONSULTANT (PATHOLOGIST) | | | | | |
| 25 | JUNIOR CONSULTANT(ENT) | | | | | |
| 26 | JUNIOR CONSULTANT (SKIN & VD) | | | | | |
| 27 | JUNIOR CONSULTANT(CARDIOLOGY) | | | | | |
| 28 | RESIDENTIAL MEDICAL OFFICER (RMO) | | | | | |
| 29 | MEDICAL OFFICER (MO)/PHYSICIAN | | | | | |
| 30 | RADIOLOGIST | | | | | |
| 31 | PATHOLOGIST | | | | | |
| 32 | SENIOR CLINICAL PATHOLOGIST | | | | | |
| 33 | ANESTHETIST | | | | | |
| 34 | EMERGENCY MEDICAL OFFICER (EMO) | | | | | |
| 35 | INDOOR MEDICAL OFFICER (IMO) | | | | | |
| 36 | INDOOR MEDICAL OFFICER (IMO) (CARDIOLOGY) | | | | | |
| 37 | INDOOR MEDICAL OFFICER (IMO) (SURGERY) | | | | | |
| 38 | INDOOR MEDICAL OFFICER (IMO) (ORTHOPEDIC SURGERY) | | | | | |
| 39 | INDOOR MEDICAL OFFICER (IMO) (MEDICINE) | | | | | |
| 40 | INDOOR MEDICAL OFFICER (IMO) (PAEDIATRIC) | | | | | |
| 41 | INDOOR MEDICAL OFFICER (IMO) (GYN & OBS) | | | | | |
| 42 | MEDICAL OFFICER (MO-BLOOD TRANSFUSION) | | | | | |
| 43 | DENTAL SURGEON | | | | | |
| 44 | ASSISTANT SURGEON/EQUIVALENT | | | | | |
| 45 | MO (HOMEOPATH/UNANI/AYURVEDIC) | | | | | |
| 46 | ASSIST REGISTRAR (MEDICINE) | | | | | |
| 47 | ASSIST REGISTRAR (SURGERY) | | | | | |
| 48 | ASSIST REGISTRAR (OBS/GYN) | | | | | |
| 49 | ASSIST REGISTRAR (PEDIATRIC) | | | | | |
| 50 | UPAZILA FAMILY PLANNING OFFICER (UFPO) | | | | | |
| 51 | MEDICAL OFFICER (MO-CLINIC) | | | | | |

| | Qualification/ Occupational Category | (a) | (b1) | (b2) | (b3) | (C) | |
|----|---|------------|--------|-------------|-------------|-----------|--|
| | | | POSTED | ATTACHMENT/ | DEPUTED-OUT | PART TIME | |
| | | (EXPECTED) | | DEPOTED-IN | | | |
| 52 | | | | | | | |
| 52 | MEDICAL OFFICER (MCH-FP) | | | | | | |
| 53 | ASSIST UPAZILA FP OFFICER (AUFPO) | | | | | | |
| 54 | ASSIST FAMILY WELFARE OFFICER, MCH- FP(AFWO) | | | | | | |
| 55 | SACMO | | | | | | |
| 56 | FAMILY WELFARE VISITOR (FWV) | | | | | | |
| 57 | MATRON | | | | | | |
| 58 | NURSING SUPERVISOR | | | | | | |
| 59 | SENIOR STAFF NURSE | | | | | | |
| 60 | NURSE MIDWIFE | | | | | | |
| 61 | STAFF NURSE | | | | | | |
| 62 | ASSISTANT NURSE | | | | | | |
| 63 | MIDWIFE | | | | | | |
| 64 | PARAMEDIC | | | | | | |
| 65 | PHARMACIST | | | | | | |
| 66 | MEDICAL TECHNOLOGIST (LAB) | | | | | | |
| 67 | MEDICAL TECHNOLOGIST (BLOOD TRANSFUSION) | | | | | | |
| 68 | MEDICAL TECHNOLOGIST (RADIOLOGY) | | | | | | |
| 69 | MEDICAL TECHNOLOGIST (PHYSIOTHERAPY) | | | | | | |
| 70 | MEDICAL TECHNOLOGIST (DENTAL) | | | | | | |
| 71 | MEDICAL TECHNICIAN EPI/EPI TECHNICIAN | | | | | | |
| 72 | MEDICAL TECHNICIAN | | | | | | |
| 73 | | | | | | | |
| 74 | | | | | | | |
| 75 | ECG TECHNICIAN | | | | | | |
| 76 | ECHO TECHNICIAN | | | | | | |
| 77 | CARDIOGRAPHER | | | | | | |
| 78 | BIOCHEMIST | | | | | | |
| 79 | NUTRITIONIST/DIETICIAN | | | | | | |
| 80 | HEALTH EDUCATOR | | | | | | |
| 81 | SANITARY INSPECTOR | | | | | | |

| | Qualification/ Occupational Category | (a) | (b1) | (b2) | (b3) | (C) |
|-----------------|---|--------------------------|--------|---------------------------|-------------|-----------|
| | | SANCTIONED (EXPECTED) | POSTED | ATTACHMENT/ DEPUTED-IN | DEPUTED-OUT | PART TIME |
| 82 | FEMALE MEDICAL ATTENDANT | | | | | |
| 83 | WARD MASTER | | | | | |
| 84 | ATTENDANT (OT/LAB/DISPENSARY /WARD BOY/EMERGENCY) | | | | | |
| 85 | COMMUNITY HEALTH CARE PROVIDER (CHCP) | | | | | |
| 86 | HEALTH ASSISTANT | | | | | |
| 87 | FAMILY WELFARE ASSISTANT (FWA) | | | | | |
| 88 | COUNSELOR | | | | | |
| 89 | COMMUNITY MOBILIZER/SERVICE PROMOTER | | | | | |
| 90 | STORE KEEPER | | | | | |
| 91 | STATISTICIAN/STATISTICAL OFFICER/STATISTICAL ASSISTANT | | | | | |
| 92 | OFF ASSITANT CUM DATA ENTRY OPERATOR | | | | | |
| 93 | UPAZILA FAMILY PLANNING ASSISTANT (UFPA) | | | | | |
| 94 [‡] | FAMILY PLANNING INSPECTOR (FPI) | | | | | |
| 95 [‡] | HEALTH INSPECTOR | | | | | |
| 96 [‡] | ASSISTANT HEALTH INSPECTOR | | | | | |
| 97 | OTHER-1 | | | | | |
| 98 | OTHER-2 | | | | | |
| 99 | TOTAL | | | | | |

QUALITY ASSURANCE

NOTIFY THE RESPONDENT THAT THIS SUBSECTION REQUIRES LOOKING AT RECORDS OF QUALITY ASSURANCE ACTIVITIES. IT WILL THEREFORE BE HELPFUL IF SUCH RECORDS ARE GATHERED BEFORE PROCEEDING WITH THE INTERVIEW.

| 440 | Does this facility routinely carry out quality assurance activities? An example may be facility-wide review of mortality, or periodic audit of registers. | YES1 NO2 DON'T KNOW |
|-----|---|--|
| 441 | Is there an official record of any quality assurance activities carried out during the past year? | YES1 NO, RECORDS NOT MAINTAINED2 +445 |
| 442 | May I see a record of any quality assurance activity? A REPORT OR MINUTES OF A QA MEETING, A SUPERVISORY CHECKLIST, A MORTALITY REVIEW, AN AUDIT OF RECORDS OR REGISTERS ARE ALL ACCEPTABLE. | OBSERVED1 REPORTED NOT SEEN2 |

TRANSPORT

| 445* | Does this facility have any vehicles including ambulance, rickshaw, boat, three-wheeler? | YES1 NO2 → 460 |
|------|---|---------------------------|
| 446* | For each type of vehicles, please tell me: | Ambulance Engine boat |
| | How many of this type of vehicle the facility currently owns, rents or has full time access to? | Rickshaw/ Speed boat Van |
| | | Boat Auto/ Three Wheelers |
| 450 | Does this facility have a functional ambulance or other vehicle for emergency transportation for clients that is stationed at this facility and that operates from this facility? | YES1 NO2 |

HMIS

FIND THE PERSON RESPONSIBLE FOR HEALTH INFORMATION SYSTEMS. INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE ASSESSMENT BEFORE PROCEEDING WITH QUESTIONS IN THIS SUBSECTION

| 460 | Does this facility have a system in place to | YES, THERE IS A DEDICATED |
|-----|--|---------------------------|
| | regularly collect health/family planning | PERSON1 |
| | services data? | NO DEDICATED PERSON2 500 |

| 464 | Does this facility have a designated person, such as a statistician, who is responsible for health services data in this facility? | YES 1 NO DEDICATED PERSON 2 – | → 500 |
|-----|--|--|-------|
| 465 | Who is responsible for health services data in this facility? PROBE TO DETERMINE WHO THIS PERSON IS | DATA MANAGER/HMIS PERSON/STATISTICIAN1 FACILITY IN-CHARGE2 OTHER SERVICE PROVIDER | |

SECTION 5: PROCESSING OF INSTRUMENTS

ASK TO BE SHOWN THE MAIN LOCATION WHERE SURGICAL INSTRUMENTS ARE PROCESSED/STERILIZED IN THE FACILITY FOR REUSE. FIND THE PERSON MOST KNOWLEDGEABLE ABOUT PROCESSING OF SURGICAL INSTRUMENTS IN THE FACILITY. INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE SURVEY AND PROCEED

| 500 | CHECK 201: ARE ANY EQUIPMENT PROCESSED IN THE FACILITY? YES (CODES "1" OR "2" CIRCLED) | | | | NO (CODE "S | 3" CIRCLED |)) | | |
|-----|---|--|------------------|---------------------------------|---|---------------------------------|-----------------------|-----------------------|---|
| | | | | • | | | | \blacksquare | |
| | | | | | GU TU NEXT | | | | |
| | | | | | SECTION OR SERVIC | CESITE | | | |
| 501 | ASK IF EACH TO SEE IT. AS | OF THE INDICATED K IF IT IS FUNCTION | ITEM ING C | IS BELOW IS DR NOT | USED BY THE FACI | LITY AND . | AVAILA | BLE. IF | AVAILABLE, ASK |
| | FOR EXAMPL | E: "Do you use [ME ⁻ | THOD |] in facility? | "IF YES, ASK: "May | I see it?" 1 | THEN" I | s it func | tioning?" |
| | ITEM | | | (a) USE AND A | VAILABILITY | T | (b) FUN | CTIONING | 3 |
| | | | | OBSERVED | REPORTED NOT SEEN | NOT USED | YES | NO | DON'T KNOW |
| 01 | ELECTRIC AUTOC HEAT) | LAVE (PRESSURE & WET | | 1 → b | 2 — b | 3 02 4 | 1 | 2 | 8 |
| 02 | NON-ELECTRIC A WET HEAT) | UTOCLAVE (PRESSURE 8 | L | 1→b | 2 - ▶b | 3 03 4 | 1 | 2 | 8 |
| 03 | ELECTRIC DRY HE | AT STERILIZER | | 1 → b | 2 → b | 3 04 | 1 | 2 | 8 |
| 04 | ELECTRIC BOILER PRESSURE) | OR STEAMER (NO | | 1 → b | 2 -→ b | 3 05 | 1 | 2 | 8 |
| 05 | NON-ELECTRIC POT WITH COVER FOR BOILING/STEAM | | 1 | 2 | 3 | | | | |
| 06 | HEAT SOURCE FO | DR NON-ELECTRIC DVE OR COOKER) | | 1 → b | 2 -> b | 3 07 4 | 1 | 2 | 8 |
| 07 | AUTOMATIC TIM EQUIPMENT) | IER (MAY BE ON | | 1→b | 2 - ▶ b | 3 08 4 | 1 | 2 | 8 |
| 08 | TST INDICATOR S | TRIPS/OTHER ITEM THA ESS IS COMPLETE | т | 1 | 2 | 3 | | | |
| 09 | ANY CHEMICALS | FOR CHEMICAL HLD | | 1 | 2 | 3 | | | |
| 502 | CHECK Q501. FOR RESPONDENT AN | EACH OF THE FOLLOWIN D INDICATE THE PROCESS | ig met Ing de | HODS OF STERI | LIZATION/HIGH LEVEL DIS NG PROCESSING TIME, RE | INFECTION T | HAT IS US D PRESSU | ED IN THE RE, ETC. | FACILITY, ASK YOUR |
| | | (1) AUTOCLAVE (Steam with pressure) | S | (2) DRY HEAT TERILIZATION | (3) BOILING (HLD) | (4) STEAM HIG DISINFECTIO | GH LEVEL ON (HLD) | CHE | (5) MICAL HIGH LEVEL SINFECTION (HLD) |
| А | Method | USED 1 NOT USED 2 2 | USED NOT L | $\frac{1}{2} \rightarrow 3$ | USED 1 NOT USED 2 - 4 | USED NOT USED | 1 2 → 5 | USED | ¹ |
| В | Temperature (centigrade) | Temperature | Temperature | |] | | | | |
| | | DON'T KNOW 998 | DON'T | KNOW 998 | | | | | |
| C | Pressure | Pressure | | | | | | | |
| | | AUTOMATIC 666 DON'T KNOW 998 | | | | | | | |
| D | Units of | UNITS OF PRESSURE: | | | | | | | |
| | pressure | KG/SQ CM1 | | | | | | | |

| | | ATM PRESSURE2 KILOPASCAL (LB/IN ²) 3 MILLIMETER HG4 DON'T KNOW8 | | | | | |
|-----|--|--|---|-----------------------------------|----------------|-------------------------|--|
| E | What is the duration in minutes when instrument is not wrapped in cloth for | | | | | | |
| | [METHOD]? | NOT USED | DON'T KNOW | DON'T KNOW 998 | DON'T KNOW 998 | DON'T KNOW 998 | |
| F | What is the duration in minutes when instrument is wrapped in cloth for autoclave? | AUTOMATIC | | | | | |
| G | Chemical disinfectant used | | | | | ALCOHOL | |
| 503 | Does this fa processing or | Does this facility have any guidelines on final processing or sterilization of surgical instruments? | | | | 1 2 →Next section | |
| 504 | May I see sterilization o HAND-WRITTE AREA WHERE E IS ACCEPTABLI | the guidelines or of equipment? N GUIDELINES POST EQUIPMENT IS PROCES | n processing or ED ON WALLS IN ISED OR STERILIZED | OBSERVED 1 REPORTED NOT SEEN 2 | | | |

| | E DERSON RESPONSIBLE FOR WASTE MANAGEM | | | | | | | |
|-----|---|---|--|--|--|--|--|--|
| | EXPLAIN THE PURPOSE OF THE ASSESSMEN | IT BEFORE PROCEEDING WITH THE QUESTIONS | | | | | | |
| 600 | Now I would like to ask you a few questions | BURN IN INCINERATOR | | | | | | |
| | about waste management practices for | 2-CHAMBER INDUSTRIAL | | | | | | |
| | sharps waste such as needles or blades. | (800-1000+°C)02 | | | | | | |
| | How does this facility <i>finally</i> dispose of <i>sharps waste</i> (e.g., filled sharps hoves)? | 1-CHAMBER DRUM/BRICK03 | | | | | | |
| | Since waste (e.g., med sharps boxes): | OPEN BURNING | | | | | | |
| | PROBE TO ARRIVE AT CORRECT RESPONSE | FLAT GROUND-NO | | | | | | |
| | NOTE | PROTECTION04 | | | | | | |
| | IF ANY OF THE RESPONSE 02-09 TAKE PLACE OUTSIDE | PIT OR PROTECTED FLAT | | | | | | |
| | THE FACILITY, THEN THE CORRECT | GROUND05 | | | | | | |
| | RESPONSE TO CIRCLE WILL BE IN THE CATEGORY OF "REMOVE OFE SITE" | DUMP WITHOUT BURNING | | | | | | |
| | | FLAT GROUND-NO- | | | | | | |
| | | PROTECTION06 | | | | | | |
| | | COVERED PIT OR PIT | | | | | | |
| | | LATRINE07 | | | | | | |
| | | OPEN PIT-NO PROTECTION08 | | | | | | |
| | | PROTECTED GROUND OR PIT09 | | | | | | |
| | | REMOVE OFF SITE | | | | | | |
| | | STORED IN COVERED | | | | | | |
| | | CONTAINER | | | | | | |
| | | STORED IN OTHER PROTECTED | | | | | | |
| | | ENVIRONMENT11 | | | | | | |
| | | STORED UNPROTECTED12 | | | | | | |
| | | OTHER96 | | | | | | |
| | | (specify) | | | | | | |
| | | NEVER HAVE SHARPS WASTE | | | | | | |
| 601 | Now I would like to ask you a few questions | SAME AS FOR SHARP ITEMS01 | | | | | | |
| | about waste management practices for | BURN IN INCINERATOR | | | | | | |
| | medical waste other than sharps, such as | 2-CHAMBER INDUSTRIAL | | | | | | |
| | used bandages | (800-1000+°C)02 | | | | | | |
| | How does this facility <i>finally</i> dispose of <i>medical waste</i> other than sharps boxes? | 1-CHAMBER DRUM/BRICK03 | | | | | | |
| | | OPEN BURNING | | | | | | |
| | PROBE TO ARRIVE AT CORRECT RESPONSE | FLAT GROUND-NO | | | | | | |
| | NOTE! | PROTECTION04 | | | | | | |
| | IF ANY OF THE RESPONSES 02-09 TAKE | PIT OR PROTECTED FLAT | | | | | | |
| | PLACE | GROUND | | | | | | |
| | OUTSIDE THE FACILITY, THEN THE | | | | | | | |
| | CORRECT RESPONSE TO CIRCLE WILL BE IN | FLAT GROUND-NO- | | | | | | |
| | | | | | | | | |
| | CATEGORT OF REIVIOVE OFF SITE | | | | | | | |
| | | | | | | | | |

SECTION 6: HEALTH CARE WASTE MANAGEMENT AND CLIENT LATRINE

| | | OPEN PIT-NO PROTECTION 08 |
|--|--|--|
| | | PROTECTED GROUND OR PIT 09 |
| | | REMOVE OFF SITE |
| | | STORED IN COVERED |
| | | CONTAINER 10 |
| | | STORED IN OTHER PROTECTED |
| | | ENVIRONMENT11 |
| | | STORED UNPROTECTED 12 |
| | | OTHER96 |
| | | (SPECIFY) |
| | | NEVER HAVE SHARPS WASTE |
| 602 | СНЕСК Q.600 | |
| | FACILITY-BASED WASTED | NEITHER FACILITY-BASED |
| | DISPOSAL OR WASTE | WASTE DISPOSAL NOR |
| | REMOVED OFFSITE | REMOVAL OFF SITE |
| | (ANY CODE OTHER THAN"95"CIRCLED) | (CODE "95"CIRCLED) |
| 603 | ASK TO SEE THE PLACE USED BY THIS FACILITY | NO WASTE VISIBLE1 |
| | FOR DISPOSAL OF SHARPS WASTE AND | WASTE VISIBLE, BUT PROTECTED AREA2 |
| | INDICATE THE CONDITION OBSERVED. IF SHARPS WASTE IS DISPOSED OFF-SITE, OBSERVE | WASTE VISIBLE, NOT PROTECTED3 |
| | THE SITE WHERE IT IS STORED PRIOR TO | WASTE SITE NOT INSPECTED8 |
| | COLLECTION FOR OFF-SITE DISPOSAL. | |
| | IF SITE NOT INSPTECTED, CIRCLE '8'. | |
| | | |
| | | |
| 604 | CHECK Q.601 | |
| 604 | CHECK Q.601 FACILITY-BASED WASTE | NEITHER FACILITY-BASED |
| 604 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE |
| 604 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO "96"CIRCLED) | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OB "95"CIRCLED) |
| 604 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) |
| 604 605 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE |
| 604 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFE SUTE OPSERVE THE SUTE WHERE IT IS | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE |
| 604 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE |
| 604 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE |
| 604 605 606 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) 606 NO WASTE VISIBLE |
| 604 605 606 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) 606 NO WASTE VISIBLE |
| 604 605 606 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) 606 NO WASTE VISIBLE |
| 604 605 606 607 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE 'S'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) ASK TO BE SHOWN THE INCINERATOR | NEITHER FACILITY-BASED 606 WASTE DISPOSAL NOR 606 REMOVAL OFFSITE 606 (CODE "01" OR "95"CIRCLED) 606 NO WASTE VISIBLE 1 WASTE VISIBLE, BUTPROTECTED 1 AREA 2 WASTE VISIBLE, NOT PROTECTED 3 WASTE SITE NOT INSPECTED 8 INCINERATOR NOT USED 610 INCINERATOR OBSERVED 1 INCINERATOR BEPORTED NOT SEEN 2 |
| 604 605 606 607 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) ASK TO BE SHOWN THE INCINERATOR | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE WASTE VISIBLE, BUTPROTECTED AREA 2 WASTE VISIBLE, NOT PROTECTED AREA 2 WASTE SITE NOT INSPECTED INCINERATOR NOT USED (NEITHER "2" NOR "3" CIRCLED) INCINERATOR OBSERVED INCINERATOR REPORTED NOT SEEN |
| 604 605 606 607 608 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) ASK TO BE SHOWN THE INCINERATOR Is the incinerator functional today? ACCEPT BEDORTED DESDONSE FROM | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE WASTE VISIBLE, BUTPROTECTED AREA 2 WASTE VISIBLE, NOT PROTECTED AREA 2 WASTE SITE NOT INSPECTED INCINERATOR NOT USED (NEITHER "2" NOR "3" CIRCLED) INCINERATOR OBSERVED INCINERATOR REPORTED NOT SEEN 2 YES NO |
| 604 605 606 607 608 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) ASK TO BE SHOWN THE INCINERATOR Is the incinerator functional today? ACCEPT REPORTED RESPONSE FROM KNOWLEDGEABLE RESPONDENT | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE |
| 604 605 606 607 608 609 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) ASK TO BE SHOWN THE INCINERATOR IS the incinerator functional today? ACCEPT REPORTED RESPONSE FROM KNOWLEDGEABLE RESPONDENT Is fuel available today for the incinerator? | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE MASTE VISIBLE, BUTPROTECTED AREA 2 WASTE VISIBLE, NOT PROTECTED AREA 2 WASTE SITE NOT INSPECTED INCINERATOR NOT USED (NEITHER "2" NOR "3" CIRCLED) INCINERATOR REPORTED NOT SEEN 2 YES 1 NO 2 YES |
| 604 605 606 607 608 609 | CHECK Q.601 FACILITY-BASED WASTE DISPOSAL OR WASTE REMOVED OFFSITE (ANY CODE "02" TO"96"CIRCLED) ASK TO SEE THE PLACE USED BY THIS FACILITY FOR DISPOSAL OF MEDICAL WASTE AND INDICATE THE CONDITION OBSERVED. IF MEDICAL WASTE IS DISPOSED OFF-SITE, OBSERVE THE SITE WHERE IT IS STORED PRIOR TO COLLECTION FOR OFF-SITE DISPOSAL. IF SITE NOT INSPTECTED, CIRCLE '8'. CHECK Q.600 and/or, Q.601 INCINERATOR USED (EITHER "2" OR "3" CIRCLED) ASK TO BE SHOWN THE INCINERATOR IS the incinerator functional today? ACCEPT REPORTED RESPONSE FROM KNOWLEDGEABLE RESPONDENT Is fuel available today for the incinerator? ACCEPT REPORTED RESPONSE | NEITHER FACILITY-BASED WASTE DISPOSAL NOR REMOVAL OFFSITE (CODE "01" OR "95"CIRCLED) NO WASTE VISIBLE MASTE VISIBLE, BUTPROTECTED AREA 2 WASTE VISIBLE, NOT PROTECTED AREA 2 WASTE SITE NOT INSPECTED 1 INCINERATOR NOT USED (NEITHER "2" NOR "3" CIRCLED) 610 610 610 YES 1 NO 2 YES 1 NO 2 |

| 610 | Do you have any guidelines on health care waste management available in this service area? This may be part of the infection prevention guideline or protocol. | YES 1 NO GUIDELINE AVAILABLE 2— | →620 |
|-----|---|------------------------------------|------|
| 611 | May I see the guidelines on health care waste management? | OBSERVED1 REPORTED NOT SEEN2 | |

CLIENT LATRINE

| - | | | 1 |
|------|--|---------------------------------------|-----|
| 620 | Is there a toilet (latrine) in functioning | FLUSH OR POUR FLUSH TOILET | |
| | condition that is available for general | FLUSH TO PIPED SEWER SYSTEM11 | |
| | outpatient client use? | FLUSH TO SEPTIC TANK12 | |
| | IF YES, ASK TO SEE THE CLIENT TOILET AND | FLUSH TO PIT LATRINE13 | |
| | INDICATE THE TYPE. THIS MUST BE TOILET | FLUSH TO SOMEWHERE ELSE14 | |
| | FACILITIES FOR THE MAIN OUTPATIENT | FLUSH, DON'T KNOW WHERE15 | |
| | SERVICE AREA. | PIT LATRINE | |
| | | VENTILATED IMPROVED PIT LATRINE21 | |
| | | PIT LATRINE WITH SLAB22 | |
| | | PIT LATRINE WITHOUT SLAB / OPEN PIT23 | |
| | | COMPOSTING TOILET | |
| | | BUCKET TOILET41 | |
| | | HANGING TOILET / HANGING LATRINE51 | |
| | | NO FUNCTIONING TOILET61 | 700 |
| 620A | ASK TO SEE THE CLIENT TOILET AND | PRIVACY | |
| | INDICATE THE PRIVACY AND FUNCTION OF | UNLOCKED DOOR WHEN NOT IN USEA | |
| | THE TOILET THIS MUST BE TOILET FACILITIES | CAN BE LOCKED FROM INSIDE WHEN USEB | |
| | FOR THE MAIN OUTPATIENT SERVICE AREA. | | |
| | | FUNCTION | |
| | | | |
| | | | |
| | | NO CRACK OR LEAK IN THE | |
| | | TOILET STRUCTUREF | |
| | | HOLE OR PIT IS NOT BLOCKED. | |
| 620B | Is there a separate sanitary | SEPARATE SANITARY TOILET | |
| | toilet/latrine facility for the use of | FACILITY FOR THE USE OF FEMALE | |
| | female clients? | CLIENTS1 | |
| | | NO SEPARATE TOILETS, ONLY | |
| | | COMBINED TOILETS2 | |

SECTION 7: BASIC SUPPLIES - CLIENT EXAMINATION ROOM CLIENT WAITING AREA

AT THIS POINT TELL YOUR RESPONDENT THAT YOU WOULD LIKE TO SEE SOME BASIC SUPPLIES AND EQUIPMENT USED IN THE PROVISION OF CLIENT SERVICES. YOU WOULD LIKE TO SEE IF THESE SUPPLIES AND EQUIPMENT ARE AVAILABLE IN THE GENERAL OUTPATIENT AREA. IF YOU ARE NOT IN THE GENERAL OUTPATIENT AREA, ASK TO BE TAKEN TO THE GENERAL OUTPATIENTAREA.

| | BASIC SUPPLIES AND EQUIPMENT | | | | | | | | | |
|-----|--|---------------|-----------------------------|----------------------|-----------------|----|---------------|--|--|--|
| 700 | I would like to know if the following | a) AVAILABLE | E | | (b) FUNCTIONING | | | | | |
| | items are available today in the main service area and are functioning ASK TO SEE ITEMS | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABL E | YES | NO | DON'T KNOW | | | |
| 01 | ADULT WEIGHING SCALE | 1_ ▶ b | ²→ ^b | 3 02◀ | 1 | 2 | 8 | | | |
| 02 | CHILD WEIGHING SCALE [250 GRAM GRADATION] | 1_ _ b | ² → ^b | 3 ₀3 ◀ | 1 | 2 | 8 | | | |
| 03 | INFANT WEIGHING SCALE [100 GRAM GRADATION] | 1_ → b | 2 _ ▶ ^b | 3 ₀4 ◀ | 1 | 2 | 8 | | | |
| 04 | STADIOMETER (OR HEIGHT ROD) FOR MEASURING HEIGHT | 1 | 2 | 3 | | | | | | |
| 05 | MEASURING TAPE (FOR HEAD CIRCUMFERENCE) | 1_ _ b | 2 _ ▶b | 3 06 ◀ | 1 | 2 | 8 | | | |
| 06 | THERMOMETER | 1 b | ² → ^b | 3 07 ◀ | 1 | 2 | 8 | | | |
| 07 | STETHOSCOPE | 1_ → b | ² → ^b | 3 ₀8 ◀ | 1 | 2 | 8 | | | |
| 08 | DIGITAL BP APPARATUS | 1_ _ b | ² → ^b | 3 09 ◀ | 1 | 2 | 8 | | | |
| 09 | MANUAL BP APPARATUS | 1_ → b | ² → ^b | 3 10 ◀ | 1 | 2 | 8 | | | |
| 10 | LIGHT SOURCE (FLASHLIGHT ACCEPTABLE) | 1_ → b | ² → ^b | 3 11 ◀ | 1 | 2 | 8 | | | |
| 11 | SELF-INFLATING BAG AND MASK [ADULT] | 1_ _ b | ² → ^b | 3 12 ◀ | 1 | 2 | 8 | | | |
| 12 | SELF-INFLATING BAG AND MASK [PEDIATRIC] | 1_ → b | ² → ^b | 3 13 ◀ | 1 | 2 | 8 | | | |
| 13 | NEBULIZER | 1_ → b | ² → ^b | 3 14 ◀ | 1 | 2 | 8 | | | |
| 14 | SPACERS FOR INHALERS | 1 | 2 | 3 | | | | | | |
| 15 | PEAK FLOW METERS | 1_ → b | ² → ^b | 3 16 ◀ | 1 | 2 | 8 | | | |
| 16 | PULSE OXIMETER | 1_ b | ² → ^b | 3 17 ◀ | 1 | 2 | 8 | | | |

| 17 | OXYGEN CONCENTRATORS | 1_ ▶ b | ²→ ^b | 3 | 1 | 2 | 8 |
|----|--|---------------|-------------------------|-----------|---|---|---|
| | | | | 18 | | | |
| 18 | FILLED OXYGEN CYLINDER WITH FLOW METER | 1 _ ▶b | ² → ^b | 3 19 ◀ | 1 | 2 | 8 |
| 19 | FILLED OXYGEN CYLINDER WITHOUT FLOW METER | 1_ → b | ² → ^b | 3 20 ◀ | 1 | 2 | 8 |
| 20 | OXYGEN DISTRIBUTION SYSTEM | 1_ _ b | ² → ^b | 3 21 ◀ | 1 | 2 | 8 |
| 21 | INTRAVENOUS INFUSION KITS - ADULT | 1 | 2 | 3 | | | |
| 22 | INTRAVENOUS INFUSION KITS - PEDIATRIC | 1 | 2 | 3 | | | |
| 23 | GLUCOMETER | 1 _ ▶b | ² → ^b | 3 24 ◀ | 1 | 2 | 8 |
| 24 | TAPE FOR MID-UPPER ARM CIRCUMFERENCE (MUAC) | 1 | 2 | 3 | | | |
| 25 | TONGUE DEPRESSOR (WOODEN/METALLIC) | 1 | 2 | 3 | | | |

CLIENT EXAMINATION ROOM

AT THIS POINT ASK TO BE SHOWN THE ROOM OR AREA IN THE GENERAL OUTPATIENT AREA WHERE MOST CLIENT SERVICES ARE OFFERED. OBSERVE THE CONDITION UNDER WHICH MOST CLIENT EXAMINATION TAKE PLACE. INDICATE IF THE FOLLOWING ITEMS ARE AVAILABLE IN THE ROOM OR AREA. ASK TO BE SHOWN ITEMS THAT YOU DO NOT SEE.

| 710 | STANDARD PRECAUTIONS AND CONDITIONS FOR CLIE EXAMINATION | NT | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | | |
|-----|--|---|-----------|-----------------------|---------------|--|--|
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POUR PITCHE | R) | 1 | 2 | 3 | | |
| 02 | HAND-WASHING SOAP (MAY BE LIQUID SOAP) | | 1 | 2 | 3 | | |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 | | |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLASTIC BIN L | INER | 1 ₀6 ◀ | 2 | 3 | | |
| 05 | OTHER WASTE RECEPTACLE | | 1 | 2 | 3 | | |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | 1 | 2 | 3 | | | |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 | | |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITANE, ALCO | HOL] | 1 | 2 | 3 | | |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH NEEDLES AUTO-DISABLE SYRINGES WITH NEEDLES | OR | 1 2 | | 3 | | |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 | | |
| 11 | GOWNS | | 1 | 2 | 3 | | |
| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 | | |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 | | |
| 711 | DESCRIBE THE SETTING OF THE CLIENT EXAMINATION ROOM OR SERVICE AREA | SCRIBE THE SETTING OF THE CLIENT AMINATION ROOM OR SERVICE AREA AND V VISUA NO PR | | ATE ROOM | | | |

CLIENT WAITING AREA

| 720 | Is there a waiting area for clients where they | YES1 | |
|-----|---|-----------------------------------|---|
| | are protected from the sun and rain? | NO PROTECTED CLIENT WAITING AREA2 | |
| | ASK TO SEE THE CLIENT WAITING AREA. MUST BE THE | | |
| | ASK TO SEE THE CLIENT WAITING AREA. MOST BE THE | | L |

| | WAITING AREA IN THE MAIN OUTPATIENT SERVICE AREA. | |
|---|---|--|
| 1 | | |

SECTION 8: DIAGNOSTICS

| 800 | | CHECK Q.102.17 | | | | | | | | | |
|---------------------------|--|--|---------------|---|-------------------------------|------------------------------------|--|---|----------------------------------|-----------------|------------------------------------|
| | | DIAGNOSTIC | SERVICE | NO DIAGNOSTIC SERVIVCES | | | | | | | |
| | | AVAILABLE IN FACII | LITY | GO TO NEXT SECTION OR | | | | | | | |
| | | | | | | | SERVI | CESITE | | | |
| ASK TO EXPLAIN BE A | BE SHO N THE P VAILA | DWN THE MAIN LABORATORY OR LOC, PURPOSE OF THE SURVEY. FOR EACH C BLE. IF INFORMATION IS NOT IN THAT | ATION IN TH | E FACILITY W DF INTEREST, ASK IF IT IS AN | HERE MO ASK AND NYWHERE | ST TESTIN GO TO TI ELSE IN T | IG IS DONE TO START HE MAIN LOCATION I THE FACILITY AND GC | DATA COLLECTION. N THE FACILITY WHE THERE TO COMPLE | INTRODU RE THE IN | CE YOU IFORM | RSELF AND ATION WILL NNAIRE. |
| | | | | HEM | ίατο | | , | | | | |
| 0.01 | De | as this facility do the follow | ing tosti | | | VES | | | | 1 | |
| 801 | in the facility? | | | | | | | .1 .2 — | → ₈₃₀ | | |
| 802 | Plea | ase tell me if: | (a | ı) | | | (b) | | | (c) | |
| a) | Any hen use | of the following noglobin test equipment is d in this facility, | US | ED | EQU | IPMENT, | ALL ITEMS FOR TE | ST AVAILABLE | IS WO | THE IT RKING | EM IN ORDER? |
| b) | All i ava | tems needed for the test are ilable, and | YES | NO | OBSE | RVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| c) | Equ | ipment is in working order | | | | | | | | | |
| 01 | Her | natology analyzer | 1 - ▶b | 2 | 1 - | С | 2 → c | 3 | 1 | 2 | 8 |
| | (for full cou | total lymphocyte count, blood count, platelet nt, etc.) | | 02 | | | | 02 ◀² | | | |
| 02 | Her | noCue | 1 — ▶b | 2 03 ◀ | 1 → c | | 2 -> c | 3 ₀₃ ◀ | 1 | 2 | 8 |
| 03 | Mic | rocuvette | 1 - ▶b | 2 | 1 | L | 2 | 3 | | | |
| | (wit | h valid expiration date) | | ₀₄ 🕈 | | | | 04 ◀ | | | |
| 04 | Colo hen | orimeter or noglobinometer | 1 → b | 2 05 ◀ | 1 - | ►с | 2 -> c | 3 05 ◀ | 1 | 2 | 8 |
| 05 | Dra | bkin's solution | 1 - ▶b | 2 | 1 | L | 2 | 3 | | | |
| | (for hen | colorimeter and noglobinometer) | | 06 🕶 | | | | 06 | | | |
| 06 | Pipe volu | ette (for measuring blood ume) | 1 → b | 2 07 | 1 | L | 2 | 3 ₀₇ ◀ | | | |
| 07 | Tall test | quist paper for hemoglobin (with valid expiration date) | 1 - ▶b | 2 830 | 1 | L | 2 | 3 | | | |
| | | | | CLINICA | | EMIST | RY | | | | |
| 830 | Doe | es this facility do any blood | glucose | testing ir | n the | YES | | | | .1 | |
| | faci | lity? | - | • | | NO | | | | .2 | → ₈₃₆ |
| 831 | Plea | ase tell me if: | (á | a) | | | (b) | | | (c) | |
| a) | Any gluc in tl | of the following blood cose test equipment is used nis facility | US | USED EQU | | | ALL ITEMS FOR TE | ST AVAILABLE | IS THE ITEM IN WORKING ORDER? | | |
| b) | Equ | ipment is available, and | YES | NO | OBSERVED | | REPORTED, | NOT AVAILABLE | YES | NO | DON'T KNOW |
| c) | Equ | ipment is in working order | 4.51 | | | <u> </u> | | | | - | |
| 01 | Equ | ipment is in working order | 1 -₽ b | ² 836 | 1- | ₽ C | 2 C | 3 836 ∢ | 1 | 2 | 8 |
| 02 | Glu | cometer test strips | | | 1- | ►c | 2 -> c | 3 836 | 1 | 2 | 8 |
| 836 | Does this facility do any urine chemistry testing using YES1 | | | | | | | | | | • |

| | dipsticks and/or urine preg | d/or urine pregnancy test on sit | | | | ? NO | | | | | | 2 | | |
|-----|--|----------------------------------|--------------|-----------------------|----------------|----------------|------------------------|----------------------|-------------------------|-------------|----------------------------------|--------------|---------------|------------------------------------|
| 837 | Please tell me if any of th | ne followi | ing | A) | USED | | (B) OBSERVED AVAILABLE | | | | | | | |
| | dipstick test is done (or u location. If done or used, see one. IF DONE/USED ASK TO SEE IT VALID/UNEXPIRED | ased) in ti I will like | his to | YES | Ν | 10 | AT LEA VA | ST ONE ILID | AVAILAE NONE V/ | BLE ALID | REPOI NOT S | RTED SEEN | N AV/ N | ORMALLY AILABLE BUT OT TODAY |
| 01 | Dip sticks for urine protein | | | 1- > b | | 2 02 | | 1 | 2 | | 3 | | 4 | |
| 02 | Dip sticks for urine glucose | | | 1 - ▶b | | 2 03 | | 1 | 2 | | 3 | | 4 | |
| 03 | Urine pregnancy test | | | 1 - ▶b | 8 | 2 40◀ | | 1 | 2 | | 3 | 5 | | 4 |
| | | PAR | ASIT | OLOGY/E | BAC | TERIO | OLOG | Ϋ́ | | | | | | |
| 840 | Please tell me if: | (a) |) | | | | (b) | | | | | (c |) | |
| a) | Any of the following EQUIPMENT is used in the facility | EQUIPMEN | NT/ TES D | στ eq | UIPM | ENT/AL AVAI | L ITEMS | 5 FOR TE | ST | | IS THE ITEM IN WORKING ORDER? | | | |
| b) | Is available, and | YES | NO | OBSERV | /ED | REPO | RTED, | NORI AVAILA | MALLY BLE NOT | Y | ΈS | NC |) | DON'T KNOW |
| c) | Equipment is functioning | | | | | NOT | SEEN | TO | DAY | | | | | - |
| 01 | LIGHT MICROSCOPE | 1 → b | 2- 02 | ▲ 1 → | С | 2 → c | | 3 02 | | | 1 | 2 | | 8 |
| 02 | ELECTRON MICROSCOPE | 1 → b | 2- 03 | ▲ 1 → | С | 2 → c | | 3 03 | | | 1 | 2 | | 8 |
| 03 | REFRIGERATOR IN LAB AREA | 1 - ▶b | 2- 04 | ▲ 1 → | ►c | 2 - | ►c | | 3 04 | | 1 | 2 | | 8 |
| 04 | INCUBATOR | 1 - ▶b | 2- 05 | ▲ 1 → | •c 2- | | ►c | | 3 05 ◀ | | 1 | 2 | | 8 |
| 05 | TEST TUBES | 1 → b | 2- 06 | ↓ ¹ | | 2 | | | ³ | | | | | |
| 07 | CULTURE MEDIUM | 1 → b | 2- 08 | ↓ ¹ | | 2 | | | 3 | | | | | |
| 08 | GLASS SLIDES AND COVERS | 1 - ▶b | 2- 848 | ↓ ¹ | | 2 | 2 | | 3 | | | | | |
| 848 | Does this facility do any GRAM STAINING? | | | | YE | S D | | | | | | | L 2 | ▶850 |
| 849 | Please tell me if the following | (a) |) | | | | (b) | | | | | | | |
| | are used and are available USED | | EQ | UIPM | ENT/AL AVAI | L ITEMS | FOR TE | ST | | | | | | |
| | IF USED ASK TO SEE IT | YES | YES NO | | /ED | REPOI | RTED, SEEN | NORI AVAILA TO | MALLY BLE NOT DAY | | | | | |
| 01 | Crystal violet or Gentian violet | 1 → b | 2- 02 | ↓ ¹ | | 2 | 2 | | 3 | | | | | |
| 02 | Lugol's iodine / Lugol's solution | 1 → b | 2 03 | ▲ 1 | | 2 | 2 | | 3 | | | | | |
| 03 | Acetone or Acetone alcohol | 1 b | 2- | 1 | | 2 | 2 | | 3 | | | | | |

| 04 | Neutral red, carbol fuchsin, or other counter stain | 1 b | 04 2 850 | 1 | | 2 | 3 | | | |
|--|--|---------------|----------------|---------|-----|-----------------------|------------------------------------|----------------------------------|---------------------|------------------|
| 850 | Do you ever send any spe staining, India Ink staining o | YE N(| /ES1 NO | | | | | | | |
| 851 | INDICATE IF THERE IS AN OBSERVED RECORD OF RESULTS FOR TESTS CONDUCTED OUTSIDE | | | | | (A) SENI OUTSIE | D SPECIMEN DE FOR TEST | (B) RECORD OF TEST OBSERVED | | |
| | | | | | | YES | NO | YE | S | NO |
| 01 | Gram stain | | | | | 1 → b | 2 02 ∢ _ | 1 | | 2 |
| 02 | India ink stain | | | | | 1 → b | 2 04 ∢ _ | 1 | | 2 |
| 04 | Specimen for culture | | | | | 1 → b | 2 | 1 | | 2 |
| 852 | Does this facility do STOOL | MICROS | COPY? | | YE | S | | | 1 2- | → 861 |
| 853 | Please tell me if the following | (; | a) | | | (b) | | | | |
| | are used and are available today. | US | SED | EQU | IPM | ENT/ALL ITEM | IS FOR TEST | | | |
| | | YES | NO | OBSERVE | D | REPORTED, NOT SEEN | NORMALLY AVAILABLE NOT TODAY | | | |
| 01 | Formal saline (for concentration method) | 1 - ▶b | 2 02 | 1 | | 2 | 3 | | | |
| 02 | Normal saline (for direct microscopy) | 1 → b | 2 03 |] 1 | | 2 | 3 | | | |
| 03 | Lugol's iodine / Lugol's solution | 1 - ▶b | 2 861 |] 1 | | 2 | 3 | | | |
| | | | ти | IBERCU | | SIS | | | | |
| 861 Does this facility do any TB tests on site? YES NO | | | | | | | | | 1 2 [—] | ► ₈₆₅ |
| 862 | Please tell me if: | (; | a) | | | (b) | | (c) | | |
| a) | Any of the following TB tests or equipment is used in the facility | TEST COI | NDUCTED | EQU | IPM | ENT/ALL ITEM | IS FOR TEST | IS THE ITEM IN WORKING ORDER? | | |
| b) | All items needed for the test are available, | YES | NO | OBSERVE | D | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| c) | Equipment is functioning | | | | | | | | | |
| 01 | Ziehl-Neelson test for AFB | 1 | 2 02 ◀ | | | | | | | |
| 02 | Carbol-Fuchsin | ¹→b | 2 03 ◀ | 1 | | 2 | 3 | | | |
| 03 | Sulphuric Acid (20 - 25% concentration) or Acid Alcohol or Hydrochloric Acid (HCl) | 1_▶b | 2 04 ◀ | 1 | | 2 | 3 | | | |
| 04 | Methylene Blue | 1_▶b | 2 05 ◀ | 1 | | 2 | 3 | | | |
| 05 | Fluorescence Microscope (FM) - LED | 1_ b | 2 06 ◀ | 1 → 0 | : | 2 - ▶ c | 3 06 ◀ | 1 | 2 | 8 |
| 06 | Culture / growth medium for Mycobacterium Tuberculosis (e.g., MGIT 960) | 1_▶b | 2 07 ◀ | 1 | | 2 | 3 | | | |

| 07 | Biosafety hood / cabinet | 1_▶b | 2 08 ◀ | 1 | | 2 | 3 | | | | |
|--------------------|--|---------------------------|-----------------------|----------------|--|---|-----------|--------|----------------------|--------------|--|
| 08 | Auramine stain for Fluorescence Microscope | 1b | 2 09 ◀ | 1 | | 2 | 3 | | | | |
| 09 [‡] | Binocular / Monocular Light Microscope | 1_ b | 2 863◀ | 1 → | C | 2 — ▶ c | 3 863_ | 1 | 2 | 8 | |
| 863 | Do you use TB rapid diagnostic diagnose TB at this laboratory | test (such / service s | n as GeneX site? | (pert) to | YE | YES1 | | | | | |
| 864 | May I see a sample TB rapid diagnostic test (RDT) kit (Such as GeneXpert Cartridge)? | | | | OBSERVED, AT LEAST 1 VALID 1 OBSERVED, NONE VALID 2 REPORTED AVAILABLE, NOT SEEN | | | | | | |
| 865 | Do you maintain any spu service site for collecting sp | utum con | ntainers ecimen? | at this | YE | S | | | 1 | → 867 | |
| 866 | May I see a sample sputum container? | | | | | OBSERVED | | | | | |
| 867 | Does this laboratory send sputum outside the facility for TB testing? | | | | YES | | | | | ₩870 | |
| 868 | Do you maintain records of result of sputum tests conducted elsewhere? | | | | | YES1 NO | | | | | |
| 869 | May I see the record or register? | | | | | OBSERVED 1 REPORTED, NOT SEEN 2 | | | | | |
| 870 | Is there a system for quality control (either internal or external) for the TB sputum smear microscopy assessed in this laboratory? | | | | | YES1 NO2 | | | | | |
| 871 | Please tell me which type of Quality Control / Quality Assurance practice is followed by this facility PROBE TO DETERMINE WHICH TYPE OF QUALTY CONTROL IS USED | | | | | INTERNAL QC / QA ONLY 1 EXTERNAL QC / QA ONLY 2 INTERNAL & EXTERNAL QC / QA 3 SEND SLIDE FOR RE-READING 4 OTHER 6 (SPECIFY) 6 | | | | | |
| 872 | Are records maintained of quality control (internal or e | of the re external) | esults fro procedu | om the res? | YES1 NO | | | | | | |
| 873 | Are records maintained for the internal QC / QA procedures, the external QC / QA procedures, or for both internal and external QC / QA procedures? | | | | | RECORDS FOR IQC / IQA ONLY | | | | | |
| DIAGNOSTIC IMAGING | | | | | | | | | | | |
| 880 | 30 Does this facility perform diagnostic X-rays, ultrasound, or computerized tomography? IF YES, ASK TO GO TO WHERE THE EQUIPMENT IS LOCATED AND SPEAK WITH THE MOST KNOWLEDGEAPLE REPORT | | | | | YES | | | | | |
| 881 | Please tell me if: | (4 | a) | | | (b) | | | (c) | | |
| a) | Any of the following imaging EQUIPMENT is used in the facility | EQUIPMI | ENT USED | | EQU | IPMENT AVAI | LABLE? | IS THE | ITEM IN WO ORDER? | ORKING | |
| b) | if it is available today, and | YES | NO | OBSERVE | D | REPORTED, | NORMALLY | YES | NO | DON'T | |

| c) | if it is functioning today | | | | NOT SEEN | AVAILABLE NOT TODAY | | | KNOW |
|---|---|--------------|-----------|---------------|-----------------------------|------------------------|----------------|---------|------|
| 01 | DIGITAL X-RAY MACHINE NOT REQUIRING FILM | 1 ▶ b | 2 02 ◀ | 1_ → c | ² → C | 3 02 | 1 | 2 | 8 |
| 02 | X-RAY MACHINE | 1 ▶ b | 2 03 ◀ | 1_ → c | ² → ^c | 3 03 _ | 1 | 2 | 8 |
| 03 | UNEXPIRED FILM FOR X-RAY | 1 ▶ b | 2 04 ◀ | 1_ → c | ² → C | 3 04 | 1 | 2 | 8 |
| 04 | ULTRASOUND SYSTEM / MACHINE | ⊥ ▶ b | 2 05 ◀ | 1_ → c | ² → ¢ | 3 05 _ | 1 | 2 | 8 |
| 05 | CT SCAN | 1 ▶ b | 2 | 1_ → c | ² → | 3 | 1 _ | 2 _ | 8 _ |
| | | NEXT SECTION | | SKIP | ▲J SKIP TO NEXT SECTION | | NEXT SECTIO | SECTION | NEXT |
| THANK YOUR RESPONDENT FOR THE TIME AND HELP PROVIDED AND PROCEED TO THE NEXT DATA COLLECTION SITE | | | | | | | | | |

SECTION 9: MEDICINES AND COMMODITIES

FACILITY STORES MEDICINES

CHECK Q210

FACILITY STORES NO MEDICINES

GO TO NEXT SECTION

SECTION 9.1: GENERAL MEDICINES AND SUPPLY ITEMS

ASK TO BE SHOWN THE MAIN LOCATION IN THE FACILITY WHERE MEDICINES AND OTHER SUPPLIES ARE STORED. FIND THE PERSON MOST KNOWLEDGEABLE ABOUT STORAGE AND MANAGEMENT OF MEDICINES AND SUPPLIES IN THE FACILITY. INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE SURVEY AND ASK THE FOLLOWING QUESTIONS

I would like to know if the following medicines are available today in this facility. If any of the medicines I mention is stored in another location in the facility, please tell me where in the facility it is stored so I can go there to verify.

ANTIBIOTICS

| 901 | Are any of the following antibiotics | (A)OBSERVE | D/ AVAILABLE | (B) NOT OBSERVED | | | |
|-----|--|-----------------------|-------------------------|------------------------|---------------------------|--------------------|--|
| | available in this facility/location TODAY? | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED, AVAILABLE | NOT AVAILABLE TODAY/DK | NEVER AVAILABLE | |
| | CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | | | NOT SEEN | | | |
| 01 | AMOXICILLIN TABLET/CAPSULE (Bacterial infections in adults) | 1 | 2 | 3 | 4 | 5 | |
| 02 | AMOXICILLIN SYRUP/ SUSPENSION OR DISPERSIBLE PEDIATRIC-DOSED TABLETS (Oral antibiotics for children) | 1 | 2 | 3 | 4 | 5 | |
| 03 | AMOXICILIN+CLAVULINATE (AUGMENTIN) TABS (broad spectrum antibiotics) | 1 | 2 | 3 | 4 | 5 | |
| 04 | AMPICILLIN INJECTION (Broad spectrum antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 05 | AZITHROMYCIN TABS/CAPS (antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 06 | AZITHROMYCIN SYR/SUSPENSION (antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 07 | BENZATHINE BENZYLPENICILLIN (POWDER) FOR INJECTION | 1 | 2 | 3 | 4 | 5 | |
| 08 | CEFIXIME TABS/CAPS (antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 09 | CEFTRIAXONE INJECTION (Injectable antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 10 | CIPROFLOXACIN (2nd-line oral antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 11 | CO-TRIMOXAZOLE (TABS) (Oral antibiotics- adult formation) | 1 | 2 | 3 | 4 | 5 | |
| 12 | CO-TRIMOXAZOLE SUSPENSION OR DISPERSIBLE PEDIATRIC-DOSED TABLET (Oral antibiotics for children) | 1 | 2 | 3 | 4 | 5 | |
| 13 | DOXYCYCLINE /CAP [Broad spectrum antibiotic] | 1 | 2 | 3 | 4 | 5 | |
| 14 | ERYTHROMYCIN [Broad spectrum antibiotic, oral tabs] | 1 | 2 | 3 | 4 | 5 | |
| 15 | ERYTHROMYCIN [oral suspension] | 1 | 2 | 3 | 4 | 5 | |
| 16 | GENTAMYCIN INJECTION (Broad spectrum injectable antibiotic) | 1 | 2 | 3 | 4 | 5 | |
| 17 | METRONIDAZOLE TABLETS [antibiotic/amebecide/antiprotozoal] | 1 | 2 | 3 | 4 | 5 | |
| 18 | METRONIDAZOLE INJECTION | 1 | 2 | 3 | 4 | 5 | |
| 19 | PENICILLIN INJECTION (Broad spectrum injectable antibiotic) | 1 | 2 | 3 | 4 | 5 | |

| 20 | TETRACYCLINE [Broad spectrum antibiotic, oral caps] | 1 | 2 | 3 | 4 | 5 | | | | |
|-----------------|---|--------------|--------------|----------|------------------|-------|--|--|--|--|
| 21 | TETRACYCLINE EYE OINTMENT | 1 | 2 | 3 | 4 | 5 | | | | |
| 23 | CHLORAMPHENICOL EYE DROP 0.5% | 1 | 2 | 3 | 4 | 5 | | | | |
| 24 | CHLORAMPHENICOL EYE OINTMENT1% | 1 | 2 | 3 | 4 | 5 | | | | |
| 25* | CIPROFLOXACIN EYE DROP 0.3% | 1 | 2 | 3 | 4 | 5 | | | | |
| 26* | AMPICILLIN CAPSULE (Broad spectrum antibiotic) | 1 | 2 | 3 | 4 | 5 | | | | |
| 27* | CEFALEXIN CAPSULE | 1 | 2 | 3 | 4 | 5 | | | | |
| 28* | CEFALEXIN DRY SYRUP | 1 | 2 | 3 | 4 | 5 | | | | |
| 29* | CEFRADINE CAPSULE | 1 | 2 | 3 | 4 | 5 | | | | |
| 30* | CEFRADINE DRY SYRUP | 1 | 2 | 3 | 4 | 5 | | | | |
| 31* | CIPROFLOXACIN DRY SYRUP | 1 | 2 | 3 | 4 | 5 | | | | |
| 32* | CLOXACILLIN CAPSULE | 1 | 2 | 3 | 4 | 5 | | | | |
| 33* | CLOXACILLIN DRY SYRUP | 1 | 2 | 3 | 4 | 5 | | | | |
| 34* | FLUCLOXACILLIN CAPSULES | 1 | 2 | 3 | 4 | 5 | | | | |
| 35* | FLUCLOXACILLIN DRY SYRUP | 1 | 2 | 3 | 4 | 5 | | | | |
| 36* | PENICILLIN-V TABLETS | 1 | 2 | 3 | 4 | 5 | | | | |
| 37* | PENICILLIN-V DRY SYRUP | 1 | 2 | 3 | 4 | 5 | | | | |
| 38* | LEVOFLOXACIN TABLETS | 1 | 2 | 3 | 4 | 5 | | | | |
| 39 [‡] | Nalidixic Acid | 1 | 2 | 3 | 4 | 5 | | | | |
| 40 [‡] | Gentamycin Eye Drop | 1 | 2 | 3 | 4 | 5 | | | | |
| 41 [‡] | Gentamicin + Hydrocortisone | 1 | 2 | 3 | 4 | 5 | | | | |
| | MEDICINES FOR WORM INFESTATION | | | | | | | | | |
| 902 | Are any of the following medicines for | (A) OBSERVE | ED AVAILABLE | | (B) NOT OBSERVED | | | | | |
| | the treatment of worm infestations | AT LEAST ONE | AVAILABLE | REPORTED | NOT AVAILABLE | NEVER | | | | |

| 902 | Are any of the following medicines for | (A) OBSERVE | | (B) NOT OBSERVED | | | |
|-----|--|-----------------------|-------------------------|-----------------------------------|---------------------------|--------------------|--|
| | the treatment of worm infestations available in the facility/location today? CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NEVER AVAILABLE | |
| 01 | ALBENDAZOLE TABLET (400mg) | 1 | 2 | 3 | 4 | 5 | |
| 02 | MEBENDAZOLE CHEWABLE TABLET (100mg OR 500mg) | 1 | 2 | 3 | 4 | 5 | |
| 03* | MEBENDAZOLE SUSPENSION (100mg/5ml) | 1 | 2 | 3 | 4 | 5 | |
| 04* | LEVAMISOLE TABLET (40mg) | 1 | 2 | 3 | 4 | 5 | |
| 05* | LEVAMISOLE SYRUP (40mg/5ml) 100ML/250ML/450ML | 1 | 2 | 3 | 4 | 5 | |

| MEDICINES FOR NON-COMMUNICABLE DISEASES | | | | | | | | | |
|---|---|-----------------------|-------------------------|-----------------------------------|---------------------------|--------------------|--|--|--|
| 903 | Are any of the following medicines | (A) OBSERVE | ED AVAILABLE | (B) NOT OBSERVED | | | | | |
| | for the treatment of non- communicable diseases available in the facility/location today? CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NEVER AVAILABLE | | | |
| 01 | AMITRIPTYLINE (Depression) | 1 | 2 | 3 | 4 | 5 | | | |
| 02 | AMLODIPINE TABLETS (CCB for high blood | 1 | 2 | 3 | 4 | 5 | | | |

| | pressure) | | | | | |
|-----|--|---|---|---|---|---|
| 03 | ATENOLOL (Beta-blocker, Angina/hypertension) | 1 | 2 | 3 | 4 | 5 |
| 04 | BECLOMETHASONE INHALER | 1 | 2 | 3 | 4 | 5 |
| 05 | BETAMETHASONE INJECTION | 1 | 2 | 3 | 4 | 5 |
| 06 | CAPTOPRIL (Vaso-dilatation, cardiac hypertension) | 1 | 2 | 3 | 4 | 5 |
| 07 | DEXAMETHASONE TABLET/INJECTION (ORADEXON) | 1 | 2 | 3 | 4 | 5 |
| 08 | DIAZEPAM INJECTION (Anxiety/muscle relaxant/anticonvulsant) | 1 | 2 | 3 | 4 | 5 |
| 09 | ENALAPRIL CAPSULE/TABLET (A.C.E INHIBITOR) | 1 | 2 | 3 | 4 | 5 |
| 10 | LOSARTAN POTASSIUM 25 mg, 50 MG, 100 mg | 1 | 2 | 3 | 4 | 5 |
| 11 | EPINEPHRINE INJECTION | 1 | 2 | 3 | 4 | 5 |
| 12 | FRUSEMIDE (DIURETIC) tablet/injection 40 mg | 1 | 2 | 3 | 4 | 5 |
| 13 | THIAZIDE DIURETIC | 1 | 2 | 3 | 4 | 5 |
| 14 | GLIBENCLAMIDE (Oral treatment for type-2 diabetes) | 1 | 2 | 3 | 4 | 5 |
| 15 | GLUCOSE INJECTABLE SOLUTION | 1 | 2 | 3 | 4 | 5 |
| 16 | HEPARIN INJECTION | 1 | 2 | 3 | 4 | 5 |
| 17 | HYDROCORTISONE INJECTION 100 mg | 1 | 2 | 3 | 4 | 5 |
| 18 | INSULIN INJECTIONS [ANTI DIABETIC] | 1 | 2 | 3 | 4 | 5 |
| 19 | ISOSORBIDE DINITRATE TABLET 10 mg | 1 | 2 | 3 | 4 | 5 |
| 20 | METFORMIN TABLETS 500 mg, 850 mg | 1 | 2 | 3 | 4 | 5 |
| 21 | NIFEDIPINE TABLETS/CAPSULES (CCB for high blood pressure) | 1 | 2 | 3 | 4 | 5 |
| 22 | OMEPRAZOLE CAP 20 mg, 40 mg (Gastro- esophageal reflux) INJECTION 40 mg | 1 | 2 | 3 | 4 | 5 |
| 23 | PREDNISOLONE Tablet 5 mg, 10mg, 20 mg / Suspension | 1 | 2 | 3 | 4 | 5 |
| 24 | SALBUTAMOL INHALER (Bronchospasms/Chronic asthma) | 1 | 2 | 3 | 4 | 5 |
| 25 | SIMVASTATIN (High cholesterol) | 1 | 2 | 3 | 4 | 5 |
| 26 | ASPIRIN TABLETS 75 mg | 1 | 2 | 3 | 4 | 5 |
| 27* | SALBUTAMOL TABLET 2mg or 4 mg (Bronchospasms/Chronic asthma) | 1 | 2 | 3 | 4 | 5 |
| 28* | SALBUTAMOL SYRUP (2mg or 4 mg) (Bronchospasms/Chronic asthma) | 1 | 2 | 3 | 4 | 5 |
| 29* | AMINOPHYLLINE TABLET (100mg) | 1 | 2 | 3 | 4 | 5 |
| 30* | ANTACID CHEWABLE TABLET/SUSPENSION (ALUM. HYDROXIDE + MAG. HYDROXIDE) | 1 | 2 | 3 | 4 | 5 |
| 31* | DIAZEPAM 5MG TABLETS | 1 | 2 | 3 | 4 | 5 |
| 32* | GLICLAZIDE Tablet, 80 Mg | 1 | 2 | 3 | 4 | 5 |
| 33* | GLYCERINE TRINITRATE Tablet 2.6 mg | 1 | 2 | 3 | 4 | 5 |
| 34* | GLIMEPIRIDE TABLET (SECRIN) 1,2, 3,4 mg | 1 | 2 | 3 | 4 | 5 |
| 35* | RANITIDINE Tablet 150 Mg | 1 | 2 | 3 | 4 | 5 |
| 36* | PANTOPRAZOLE Tablet 20 mg, 40 mg / Injection 40 mg | 1 | 2 | 3 | 4 | 5 |
|-----|--|-----------------------|-------------------------|-----------------------|---------------------------|-----------|
| 37* | ESOMEPRAZOLE Tablet 20 mg, 40 mg / Injection 40 mg | 1 | 2 | 3 | 4 | 5 |
| 38* | ATORVASTATIN Tablet 10 mg, 20 mg | 1 | 2 | 3 | 4 | 5 |
| 39* | ROSUVASTATIN Tablet 10 mg, 20 mg | 1 | 2 | 3 | 4 | 5 |
| 40* | CLOPIDOGRELTABLET 75 mg | 1 | 2 | 3 | 4 | 5 |
| 41* | THEOPHYLLINE Tablet 200 mg, /Syrup | 1 | 2 | 3 | 4 | 5 |
| 42* | IPRATROPIUM BROMIDE (IPREX) , Respiratory Solution | 1 | 2 | 3 | 4 | 5 |
| 43* | SALBUTAMOL RESPIRATORY SOLUTION | 1 | 2 | 3 | 4 | 5 |
| 44* | MANNITOL 20 % Infusion (Osmosol 20% IV) | 1 | 2 | 3 | 4 | 5 |
| 45* | VITAMIN B COMPLEX TABLETS | 1 | 2 | 3 | 4 | 5 |
| 46* | CHLORPHENIRAMINE TAB 4 mg | 1 | 2 | 3 | 4 | 5 |
| 47* | MICRONUTRIENT POWDER | 1 | 2 | 3 | 4 | 5 |
| | AN | TI-FUNGA | | NES | | |
| 004 | Are any of the following anti fungal | | | // | | |
| 504 | medicines available in the facility/ | AVAI | LABLE | (1 | S) NOT OBSERVE | |
| | location today? | AT LEAST | AVAILABLE | REPORTED | NOT AVAILABLE | NEVER |
| | CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | ONE VALID | NONE VALID | AVAILABLE NOT SEEN | TODAY/DK | AVAILABLE |
| 01 | FLUCONAZOLE CAPS 50mg, 150mg, 200 mg | 1 | 2 | 3 | 4 | 5 |
| 02 | MICONAZOLE VAGINAL PESSARY | 1 | 2 | 3 | 4 | 5 |
| 03 | MICONAZOLE CREAM | 1 | 2 | 3 | 4 | 5 |
| 04 | NYSTATIN ORAL SUSPENSION | 1 | 2 | 3 | 4 | 5 |
| 05 | NYSTATIN VAGINAL PESSARY/ CREAM | 1 | 2 | 3 | 4 | 5 |
| 06* | GRISEOFULVIN TABLET (500mg) | 1 | 2 | 3 | 4 | 5 |
| 07* | GENTIAN VIOLET 2% TOPICAL SOLUTION | 1 | 2 | 3 | 4 | 5 |
| 08* | NEOMYCIN & BACITRACIN SKIN OINTMENT 10mg | 1 | 2 | 3 | 4 | 5 |
| 09* | KETOCONAZOLE Tablet 200 mg / ointment | 1 | 2 | 3 | 4 | 5 |
| 11* | BENZOIC ACID 6% AND SALICYLIC ACID 3% (WHITFIELDS OINTMENT) | 1 | 2 | 3 | 4 | 5 |
| | MATE | RNAL ANI | D CHILD HI | EALTH | | |
| 906 | Are any of the following medicines | (A) OB | SERVED | (E | 3) NOT OBSERVI | ED |
| | for maternal health available in the | AVAI | LABLE | | , | |
| | | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE | NOT AVAILABLE TODAY/DK | NEVER |
| | (NOT EXPIRED) | | | NOT SEEN | | AVAILABLE |
| 01 | CALCIUM GLUCONATE INJECTION | 1 | 2 | 3 | 4 | 5 |
| 02 | FOLIC ACID TABLETS 5 mg | 1 | 2 | 3 | 4 | 5 |
| 03 | IRON TABLETS 400 mg | 1 | 2 | 3 | 4 | 5 |
| 04 | IRON 200 mg + FOLIC ACID 60 mg COMBINATION TABLET | 1 | 2 | 3 | 4 | 5 |
| 05 | MAGNESIUM SULPHATE INJECTION 2.5 gm /5ml/INFUSION 4% | 1 | 2 | 3 | 4 | 5 |

| 06 | MISOPROSTOL TABLET | 1 | 2 | 3 | 4 | 5 |
|-----------------|--|---|---|---|---|---|
| 07 | OXYTOCIN OR OTHER INJECTABLE UTEROTONIC | 1 | 2 | 3 | 4 | 5 |
| 08 | TETANUS TOXOID VACCINE | 1 | 2 | 3 | 4 | 5 |
| 09 | ORAL REHYDRATION SALTS (ORS) SACHETS | 1 | 2 | 3 | 4 | 5 |
| 10 | VITAMIIN A CAPSULE 50,000 IU/ 2 Lac IU | 1 | 2 | 3 | 4 | 5 |
| 11* | ZINC SULPHATE TABLETS 10mg, 20 mg / SYRUP | 1 | 2 | 3 | 4 | 5 |
| 12* | CALCIUM LACTATE TABLET 300 mg | 1 | 2 | 3 | 4 | 5 |
| 13* | FERROUS SULPHATE + FOLIC ACID + ZINC CAPSULE | 1 | 2 | 3 | 4 | 5 |
| 14* | FERROUS FUMERATE + FOLIC ACID TABLET | 1 | 2 | 3 | 4 | 5 |
| 15* | ZINC SULPHATE SYRUP | 1 | 2 | 3 | 4 | 5 |
| 16* | METHYL ERGOMETRINE TABLET/INJECTION 125 micro gram, 200 micro gram | 1 | 2 | 3 | 4 | 5 |
| 17* | ERGOMETRINE TABLET | 1 | 2 | 3 | 4 | 5 |
| 18* | GENTIAN VIOLET 1% | 1 | 2 | 3 | 4 | 5 |
| 19* | NYSTATIN DROPS | 1 | 2 | 3 | 4 | 5 |
| 20* | INJECTION NALOXONE HYDROCHLORIDE | 1 | 2 | 3 | 4 | 5 |
| 21* | 7.1 % CHLORHEXIDINE | 1 | 2 | 3 | 4 | 5 |
| 22* | CHLORHEXIDINE CREAM (60 ml bottle) | 1 | 2 | 3 | 4 | 5 |
| 23* | COTRIMOXAZOLE VT 200 mg | 1 | 2 | 3 | 4 | 5 |
| 24* | CALCIUM CARBONATE TABLET 500 mg | 1 | 2 | 3 | 4 | 5 |
| 25* | CHLORPHENIRAMINE MALEATE SYRUP (2mg/5ml) 60 ml | 1 | 2 | 3 | 4 | 5 |
| 26* | LIDOCAINE INJECTION 1% (50ml) | 1 | 2 | 3 | 4 | 5 |
| 28 [‡] | Tab. Cotrimoxazole 120 mg dispersible | 1 | 2 | 3 | 4 | 5 |
| 29 [‡] | Tab. Cotrimoxazole DS 960mg | 1 | 2 | 3 | 4 | 5 |
| 30 [‡] | Cotrimoxazole suspension 60 ml | 1 | 2 | 3 | 4 | 5 |
| 31 [‡] | Inj. Corticosteroid 100mg (Amp) | 1 | 2 | 3 | 4 | 5 |
| | | | | | | |

| | INTRAVENOUS FLUIDS | | | | | | |
|-----|--|-----------------------|-------------------------|-----------------------------------|---------------------------|--------------------|--|
| 907 | Are any of the following intravenous fluids available in the | (A) OB AVAI | SERVED LABLE | (E | B) NOT OBSERVE | Ð | |
| | facility/location today? CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NEVER AVAILABLE | |
| 01 | NORMAL SALINE 500 ML, 1000ML | 1 | 2 | 3 | 4 | 5 | |
| 02 | RINGERS LACTATE SOLUTION 1000 ML | 1 | 2 | 3 | 4 | 5 | |
| 03 | 0.9 % DEXTROSE IN NORMAL SALINE (DNS) 500 ML, 1000ML | 1 | 2 | 3 | 4 | 5 | |
| 04* | 5 % DEXTROSE IN AQUA (D/A) 500 ML, 1000ML | 1 | 2 | 3 | 4 | 5 | |
| 05* | CHOLERA SALINE 500 ML, 1000 ML | 1 | 2 | 3 | 4 | 5 | |
| 06* | HARTMAN'S SOLUTION 1000 ML | 1 | 2 | 3 | 4 | 5 | |

| 07 [‡] | W | ater for Injection (sterile/pyrogen free) | 1 | 2 | 3 | 4 | 5 | | |
|-----------------|--|---|--|--------------------------------|-----------------------------------|---------------------------|--------------------|--|--|
| | FEVER REDUCING AND PAIN MEDICINES | | | | | | | | |
| 908 | Ar m | e any of the following OTHER edicines available in the | (A) OB AVAI | SERVED LABLE | (E | 3) NOT OBSERVE | Ð | | |
| | fac CH | cility/location today? IECK TO SEE IF AT LEAST ONE IS VALID (NOT | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NEVER AVAILABLE | | |
| 01 | DI | CLOFENAC TABLETS (50 mg) OR | 1 | 2 | 3 | 4 | 5 | | |
| 02 | PA | RACETAMOL TABLET 500 mg | 1 | 2 | 3 | 4 | 5 | | |
| 03* | PA | RACETAMOL SYRUP OR SUSPENSION | 1 | 2 | 3 | 4 | 5 | | |
| 04* | PA DC | RACETAMOL DISPERSIBLE PEDIATRIC- DZED TABLET | 1 | 2 | 3 | 4 | 5 | | |
| 05* | IN | DOMETHACIN CAPSULE 25 mg | 1 | 2 | 3 | 4 | 5 | | |
| 07* | DI | CLOFENAC SODIUM INJECTION 75 mg | 1 | 2 | 3 | 4 | 5 | | |
| 09* | IB | UPROFEN TABLET 200 mg, 400mg/syrup | 1 | 2 | 3 | 4 | 5 | | |
| 10* | NA | APROXEN SODIUM 250mg, 500mg | 1 | 2 | 3 | 4 | 5 | | |
| 11* | AC | CECLOFENAC TAB, 100 mg | 1 | 2 | 3 | 4 | 5 | | |
| 12* | TR mį | AMADOL HYDROCHLORIDE CAP, 50 g, 100mg | 1 | 2 | 3 | 4 | 5 | | |
| 13* | KE IN | TOROLAC HYDROCHLORIDE, 10 mg/ J, 30 MG, 60 MG | 1 | 2 | 3 | 4 | 5 | | |
| 14* | * DICLOFENAC SODIUM SUPPOSITORY 12.5 mg. 25 mg. 50mg | | 1 | 2 | 3 | 4 | 5 | | |
| 15* | DF IN | ROTAVERINE HYDROCHLORIDE JECTION 40 mg | 1 | 2 | 3 | | 5 | | |
| 16* | НҮ (В | /OSCINE – N – BUTYL BROMIDE TABS UTAPEN) / INJECTION, 20 mg | 1 | 2 | 3 | 4 | 5 | | |
| 17* | TII 50 | EMONIUM METHYL SULPHATE TAB, mg /Injection5mg/2 ml | 1 | 2 | 3 | 4 | 5 | | |
| | | STORAGE CONDITIO | N: ANTIBI | OTICS & G | | | | | |
| 909 | | OBSERVE THE PLACE WHERE THE N AND INDICATE THE PRESENCE (OR STORAGE CONDITIONS. | /EDICINES AS ABSENCE) OF | SESSED SO FAF F EACH OF THE | R ARE STORED E FOLLOWING | YES | NO | | |
| | 01 | ARE THE MEDICINES OFF THE FLO | OR? | | | 1 | 2 | | |
| | 02 | ARE THE MEDICINES PROTECTED F | ROM THE WA | ATER? | | 1 | 2 | | |
| | 03 | ARE THE MEDICINES PROTECTED F | ROM THE SU | N? | | 1 | 2 | | |
| | 04 | IS THE ROOM CLEAN OF EVIDEN (ROACHES, ETC)? | CE OF RODEN | ITS (BATS, RA | TS) OR PESTS | 1 | 2 | | |
| | 05 | IS THE STORAGE ROOM WELL VEN | ITILATED? | | | 1 | 2 | | |
| 910 | 910 ARE THE MEDICINES ORGANIZED ACCORDING TO DATE OFEXPIRATION ("first expire, first out")? VES, ALL MEDICINES | | | | | | | | |
| 911 | | What system does this facility us the amount of medicines received issued, and the amount present to ASK TO SEE THE SYSTEM A | system does this facility use to monitor nount of medicines received, the amount , and the amount present today? TO SEE THE SYSTEM AND RECORD | | | | r | | |

| | OBSERVATION | LEDGER/STOCK CARD NOT UPDATED DAILY, BUT THERE IS DAILY RECORD OF DISTRIBUTED MEDICINES | | | | |
|-----|--|--|----------|----------------------|-----|-----------|
| 912 | Do you have the following supply items availab the facility/location today? | le in | OBSERVED | REPORTED NOT SEEN | NOT | AVAILABLE |
| 01 | SINGLE-USE STANDARD DISPOSABLE SYRINGES NEEDLES OR AUTO-DESTRUCT SYRINGES WITH NEEL | WITH DLES | 1 | 2 | | 3 |
| 02 | INFUSION SET FOR IV SOLUTION | | 1 | 2 | | 3 |
| 03 | CANULA FOR ADMINISTERING IV FLUIDS | | 1 | 2 | | 3 |
| 04 | LATEX GLOVES | | 1 | 2 | | 3 |
| 05 | ALCOHOL-BASED HAND RUB | | 1 | 2 | | 3 |
| 06 | HAND WASHING SOAP | | 1 | 2 | | 3 |
| 07 | DISINFECTING SOLUTION | | 1 | 2 | | 3 |

SECTION 9.2: CONTRACEPTIVE COMMODITIES

| 920 | CHECK Q212 CONTRACEPTIVES STORED WITH OTHER MEDICINES IN COMMON LOCATION (RESPONSE 2 CIRCLED) | ↓ ↓ | CONTR. IN FP S OR 3 CIRCLED) PROCEE (TB MEE | ACEPTIVES STOF SERVICE MEDICII D TO NEXT SECT DS?) | RED NES (RESPONSE 1 | |
|-----|---|--------------------------|--|---|-------------------------------|--------------------|
| 921 | Are any of the following CONTRACEPTIVE commodities | (A) OB AVAI | SERVED | (B) NOT OBSERVED | | |
| | available in the facility/location today? CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED, AVAILABLE NOT SEEN | NOT AVAILABLE/TO DAY/DK | NEVER AVAILABLE |
| 01 | COMBINED ORAL CONTRACEPTIVE PILLS | 1 | 2 | 3 | 4 | 5 |
| 02 | PROGESTIN-ONLY CONTRACEPTIVE PILLS | 1 | 2 | 3 | 4 | 5 |
| 04 | PROGESTIN-ONLY INJECTABLE CONTRACEPTIVES (DEPO) | 1 | 2 | 3 | 4 | 5 |
| 05 | MALE CONDOMS | 1 | 2 | 3 | 4 | 5 |
| 07 | INTRAUTERINE CONTRACEPTIVE DEVICE | 1 | 2 | 3 | 4 | 5 |
| 08* | ONE ROD IMPLANT | 1 | 2 | 3 | 4 | 5 |
| 09* | TWO ROD IMPLANT (ZADELL) | 1 | 2 | 3 | 4 | 5 |
| 10 | EMERGENCY CONTRACEPTIVE PILLS (E.G., PROSTINOL 2) | 1 | 2 | 3 | 4 | 5 |

STORAGE CONDITION - CONTRACEPTIVE COMMODITIES

| 922 | OBSERVE THE LOCATION WHERE CONTRACEPTIVE COI INDICATE THE PRESENCE (OR ABSENCE) OF EACH O CONDITIONS | MMODITIES ARE STORED AND F THE FOLLOWING STORAGE | YES | NO |
|------------|--|---|--|----|
| 01 | ARE THE COMMODITIES OFF THE FLOOR? | | 1 | 2 |
| 02 | ARE THE COMMODITIES PROTECTED FROM WATER? | | 1 | 2 |
| 03 | ARE THE COMMODITIES PROTECTED FROM THE SUN? | | 1 | 2 |
| 04 | IS THE ROOM CLEAN OF EVIDENCE OF RODENTS (BAT: ETC)? | S, RATS) OR PESTS (ROACHES, | 1 | 2 |
| 05 | IS THE STORAGE ROOM WELL VENTILATED? | | 1 | 2 |
| 923 924 | ARE THE CONTRACEPTIVE COMMODITIES ORGANIZED ACCORDING TO DATE OF EXPIRATION ("first expire, first out") What type of system does this facility use to monitor the amount of contraceptive commodities received, the amount issued, and the amount present today? ASK TO SEE THE SYSTEM AND RECORD OBSERVATION | YES, ALL COMMODITIES. NOT ALL COMMODITIES. NO. COMPUTER SYSTEM UPDATE LEDGER/STOCK CARD UPDAT DAILY. COMPUTER SYSTEM NOT UP THERE IS DAILY RECORD OF DISTRIBUTED CO LEDGER/STOCK CARD NOT UP BUT THERE IS DAILY RECORD COMMODITIES. OTHER SYSTEM (SPECIFY). | 1 2 8 D DAILY 1 ED 2 DATED DAILY, BUT MMODITIES 3 PDATED DAILY, OF DISTRIBUTED 4 | |
| 925 | PRESENTLY INTERVIEWING IN PHARMACY PROCEED TO NEXT SECTION OR SERVICE SITE | PRESENTLY INTERVIEWING I PLANNING SERVICE AREA THANK THE RESPONDENT IN TH AND CONTINUE TO NEXT SECTION OR SERVICE SITE | E FP SERVICE AREA | |

SECTION 9.3: ANTI-TB DRUGS

| 930 | CHECK Q214 | | | | |] | |
|-----------------|--|-----------------------|--|------------------------|-------------------------------|--------------------|--|
| | ANTI-TB MEDICINES STORED WITH OTHER MEDICINESIN COMMON LOCATION (RESPONSE 2 CIRCLED) | □ | ANTI-TB MEDICINES STORED IN TB SERVICE AREA OR NOT STOCKED AT ALL IN FACILITY (RESPONSE 1 OR 3 CIRCLED) PROCEED TO NEXT SECTION | | | | |
| 931 | Are any of the following TB | (A) OBSERVI | ED AVAILABLE | | (B) NOT OBSERVED | OT OBSERVED | |
| | medicines available in the facility/location today? | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED, AVAILABLE | NOT AVAILABLE/TO DAX/DK | NEVER AVAILABLE | |
| | CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | | | NOT SEEN | DATION | | |
| 01 | ETHAMBUTOL TABS (E) | 1 | 2 | 3 | 4 | 5 | |
| 02 | ISONIAZID TABS (INH, H) | 1 | 2 | 3 | 4 | 5 | |
| 03 | PYRAZINAMIDE (Z) | 1 | 2 | 3 | 4 | 5 | |
| 04 | RIFAMPICIN (R) | 1 | 2 | 3 | 4 | 5 | |
| 05 | RIFAMPICIN+ISONIAZID- RH 75/50 mg- (2FDC) Child | 1 | 2 | 3 | 4 | 5 | |
| 06 | RIFAMPICIN + ISONIAZID -RH 150/75 mg- 2FDC Adult | 1 | 2 | 3 | 4 | 5 | |
| 07 | RIFAMPICIN + ISONIAZID + PYRAZINAMIDE (RHZ)- 75/50/150 mg- 3FDC Child | 1 | 2 | 3 | 4 | 5 | |
| 08 | RIFAMPICIN + ISONIAZID + ETHAMBUTOL (RHE)- 150/75/275- 3FDC Adult | 1 | 2 | 3 | 4 | 5 | |
| 09 | RIFAMPICIN + ISONIAZID+ PYRAZINAMIDE + ETHAMBUTOL | 1 | 2 | 3 | 4 | 5 | |
| | (RHZE)- 150/75/400/275- 4FDC Adult | | | | | | |
| 10 | STREPTOMYCIN (S 1000) INJECTABLE | 1 | 2 | 3 | 4 | 5 | |
| 11 [‡] | LEVOFLOXACIN 250MG TABLET | 1 | 2 | 3 | 4 | 5 | |
| 935 | PRESENTLY INTERVIEWING IN PHARMACY | | PRESENTLY IN TB SERVICE AF | TERVIEWING IN REA | | | |
| | PROCEED TO NEXT SECTION OR SERVICE SITE | | THANK THE RESPONDENT IN THE TB SERVICE AREA AND CONTINUE TO NEXT SECTION OR SERVICE SITE | | | | |

MODULE 3: SERVICE-SPECIFIC READINESS

CHILD HEALTH SERVICES

SECTION 10: CHILD VACCINATION

| 1000 | CHECK Q102.01 | | | | | | | |
|-------|--|---|--------------------------------------|--|----------------------|--|--|--|
| | CHILD VACCINATION SERVICES | NO CHILD VA | CCINATION SERVIC | ES 🗌 | | | | |
| | | NEXT SECTI | NEXT SECTION OR SERVICE SITE | | | | | |
| ΛΟΥ Τ | | | | | | | | |
| FIND | FIND THE PERSON MOST KNOWLEDGEABLE ABOUT CHILD VACCINATION SERVICES IN THE FACILITY. INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE SURVEY AND ASK THE FOLLOWING QUESTIONS. | | | | | | | |
| 1001 | Now I would like to ask you specifically about vac the following services, please tell me whether the days per month the service is provided at the facil | cination service e service is offe ity. | es for children u red by your fac | under 5 years. I cility, and if so, | For each of how many | | | |
| | CHILD VACCINATION SERVICE (a) | | | | | | | |
| | (USE A 4-WEEK MONTH TO CALCULATE # OF DAYS | # OF DAYS | PER MONTH SERVI | CE IS PROVIDED AT | FACILITY | | | |
| 01 | Routine DPT+HepB+Hib (i.e., pentavalent) | # OF DAYS 00=NO SERVI | CE | | | | | |
| 02 | Routine polio vaccination (i.e., OPV) | # OF DAYS 00=NO SERVI | CE | | | | | |
| 03 | Routine measles vaccination (i.e., Measles - Rubella) | # OF DAYS | | | | | | |
| 04 | BCG vaccination | | | | | | | |
| | | # OF DAYS | | | | | | |
| 05* | Inactivated polio vaccination (i.e., IPV) | # OF DAYS 00=NO SERVI | CE | | | | | |
| 06* | Pneumococcal vaccination (i.e., PCV) | # OF DAYS 00=NO SERVI | CE | | | | | |
| 1002 | Do you have the national guidelines for child vaccinations (i.e. EPI SOHAYIKA) AVAILABLE in this service area today? | YES NO | | 1 2— | →1004 | | | |
| 1003 | May I see the guidelines? | OBSERVED | T SFFN | | → 1006 | | | |
| 1004 | Do you have any other guidelines for child vaccinations available in this service area today? | YES | | | →1006 | | | |
| 1005 | May I see the other guidelines? | OBSERVED REPORTED, NO | T SEEN | 1 2 | | | | |
| 1006 | ASK YOUR RESPONDENT TO SHOW YOU ITEMS REQUIRED FOR VACCINATION SERVICES | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | | | | |
| 01 | Blank/unused individual child vaccination cards or booklets (EPI card) | 1 | 2 | 3 | | | | |
| 02 | Tally sheets | 1 | 2 | 3 | | | | |
| 03 | Summary forms / monthly report | 1 | 2 | 3 | | | | |

| 1007 | Does this facility routinely store any vaccines, or | ROUTINELY STORE VACCINES1 | |
|------|---|---------------------------|-------|
| | are all its vaccines either picked up from | STORES NO VACCINES 2 — | ▶1050 |
| | another facility or delivered when services are | | |
| | being provided? | | |

| 1008 | ASK TO BE TAKEN TO THE AREA WHERE VACCINES | REFRIGEF | ATOR OBSER | VED | 1 | | |
|------|--|-----------|----------------------------|-----------|---------------|-----------|--|
| 1000 | ARE STORED. ASK TO SEE THE VACCINE | REFRIGEF | REFRIGERATOR NOT OBSERVED2 | | | | |
| | REFRIGERATOR. | | | | | | |
| 1009 | Do you maintain a cold-chain temperature | YES | | ••••• | 1 | | |
| | monitoring chart? | NO | | | 2 — | ▶ 1012 | |
| 1010 | May I see the cold-chain temperature monitoring | OBSERVE | D | | 1 | | |
| | chart? | REPORTE | D, NOT SEEN. | | | ▶ 1012 | |
| 1011 | CHECK WHETHER THE TEMPERATURE RECORD WAS | YES, CON | IPLETED | | 1 | | |
| 1011 | COMPLETED TWICE DAILY FOR EACH OF THE PAST | NO, NOT | COMPLETED. | | 2 | | |
| | 30 DAYS, INCLUDING WEEKENDS AND PUBLIC | | | | | | |
| | HOLIDAYS. | | | | | | |
| 1012 | Please tell me if each of the following vaccines is | (A) O | BSERVED | (6 | B) NOT OBSERV | ED | |
| 1012 | available in the facility today. If available, I would | AVA | AILABLE | | | | |
| | like to see it. | ATLEAST | AVAILABLE | REPORTED | NOT | NEVER | |
| | ΙΕ ΔΥΔΙΙ ΔΒΙ Ε. CHECK ΤΟ SEE ΙΕ ΔΤΙ ΕΔST ONE IS VALID | ONE VALID | NONE VALID | NOT SEEN | TODAY/DK | AVAILABLE | |
| | (NOT EXPIRED, VACCINE VIAL MONITOR UNCHANGED | | | | | | |
| | (NOT FROZEN)) | | | | | | |
| 01 | DPT+HepB+Hib [PENTAVALENT] | 1 | 2 | 3 | 4 | 5 | |
| | | | | | | | |
| 02 | ORAL POLIO VACCINE (I.e., OPV) | 1 | 2 | 3 | 4 | 5 | |
| 03 | MEASLES VACCINE AND DILUENT (i.e., MR) | 1 | 2 | 3 | 4 | 5 | |
| | | _ | _ | - | - | | |
| 04 | BCG VACCINE AND DILUENT | 1 | 2 | 3 | 4 | 5 | |
| 05 | INACTIVATED POLIO VACCINE (i.e., IPV) | 1 | 2 | з | 1 | E | |
| | | - | 2 | 5 | 4 | 5 | |
| 06 | PNEUMOCOCCAL VACCINE (i.e., PCV) | 1 | 2 | 3 | 4 | 5 | |
| 1012 | WHAT IS THE TEMPERATURE IN THE VACCINE | BETWEEN | N +2 AND +8 D | EGREES | 1 | t | |
| 1013 | REERIGERATOR? | ABOVE +8 | B DEGREES | | 2 | | |
| | | BELOW + | 2 DEGREES | | 3 | | |
| | CENTIGRATE/CELSIUS DEGREES | THERMO | METER NOT F | UNCTIONAL | 4 | | |
| | | NO THER | MOMETER | | 5 | | |

| 1050 | ASSESS THE ROOM OR AREA FOR THE ITEMS LISTED BELOW. FOR ITEMS THAT YOU DO NOT SEE, ASK | GENERAL INFOR | RMATION [Q710] E CARE [Q1251] . | 11 | |
|--------|---|----------------|------------------------------------|-----------------|--------------|
| | YOUR RESPONDENT TO SHOW THEM TO YOU. | FAMILY PLANNI | NG [Q1351] RE [01451] | 14 15 | - |
| | IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, | DELIVERY [Q16 | 51] | | |
| | INDICATE WHERE THE DATA ARE RECORDED | TUBERCULOSIS | [Q1951] | 19 | NEXT |
| | | NCD [Q2351] | | | SECTION / |
| | | | | | SERVICE SITE |
| 1051 | STANDARD PRECAUTIONS AND CONDITIONS | FOR CLIENT | OBSERVED | D NOT | AVAILABLE |
| | EXAMINATION | | | SEEN | |
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POL | JR PITCHER) | 1 | 2 | 3 |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOA | P) | 1 | 2 | 3 |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLAS | STIC BIN LINER | 1 | 2 | 3 |
| | | | ₀₆ 🚽 | | |
| 05 | | | 1 | 2 | 2 |
| 05 | | | 1 | Ζ | 3 |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITA | NE, ALCOHOL] | 1 | 2 | 3 |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH | H NEEDLES OR | 1 | 2 | 3 |
| | | | | | |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 |
| 11 | GOWNS | | 1 | 2 | 3 |
| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 |
| 1052 | DESCRIBE THE SETTING OF THE CHILD VACCINATION | PRIVATE ROOM | | 1 AND VISUAI | |
| | | PRIVACY | | | |
| | | VISUAL PRIVAC | Y ONLY | 3 | |
| | | NO PRIVACY | | 4 | |
| THANK | YOUR RESPONDENT AND MOVE TO YOUR NEXT D | | ON POINT IF DIF | FERENT FR | OM CURRENT |
| LOCATI | ON. | | | | - |

| 1100* | CHECK Q102.02 | | | | | | | |
|---------|--|--------------------------------|--------|----------------|---------------|----------|---------|-----------|
| | GROWTH MONITORING SERVICES | NO GROWTH MONITORING SERVICES | | | | es [| | |
| | AVAILABLE | NEXT SECTION OR SERVICE SITE | | | | | ← | |
| ٨sk | | GROV | | | | | | ТНЕ |
| FACILIT | Y. FIND THE PERSON MOST KNOWLEDGEABL | E ABC | DUT GF | ROWTH MON | NITORING SER | VICES IN | THE F | ACILITY. |
| IN | TRODUCE YOURSELF, EXPLAIN THE PURPOSE | OF TH | HE SUR | VEY AND AS | K THE FOLLOV | VING QL | JESTIC | NS. |
| 1101 | Please tell me the number of days per month that(a)growth monitoring services are offered in this facility.# OF DAYS PER MONTH SERVICE IS PROVIDED AT FACILITY | | | | | | ACILITY | |
| | USE A 4-WEEK MONTH TO CALCULATE # OF DAYS | | | | | | | |
| 01 | Child growth monitoring | | | | | | | |
| | | | # O | F DAYS | | | | |
| 1102 | Do you have any guidelines for growth | | YES | | | | 1 | |
| | monitoring available in this service area too | ay? NO GUIDELINE AVAILABLE | | | /AILABLE | | 2 | ► 1103A |
| 1103 | (family health card) | OBSERVED REPORTED, NOT SEEN | | | | | . 2 | |
| 1103A | Do you have any guidelines for SAM and M | AM | YES | | | | 1 | → 1104 |
| 11020 | May I see the other guidelines? | | OBSE | RVED | | <u></u> | . 1 | P 1104 |
| 11020 | (family health card) | | REPO | RTED, NOT SE | EN | | . 2 | |
| 1104 | I would like to know if the following items are | | | (a) AVAILA | BLE | (| b) Fur | nctioning |
| 1104 | available in this service area and are | OBSE | RVED | REPORTED, | NOT | YES | NO | DON'T |
| | functioning. I would like to see them. | | | NOT SEEN | AVAILABLE | | | KNOW |
| 01 | CHILD WEIGHING SCALE (250 GRAM GRADATION) | 1- | ►b | 2 - ▶ b | 3 — | 1 | 2 | 8 |
| | | | | | 02 ◀┘ | | | |
| 02 | INFANT WEIGHING SCALE (100 GRAM GRADATION) | 1— | ►b | 2 -→ b | 3 | 1 | 2 | 8 |
| 03 | HEIGHT OR LENGTH BOARD | 1- | ►b | 2 - ▶ þ | 3 - | 1 | 2 | 8 |
| | | | | | ₀₄ | | | |
| 04 | TAPE FOR MEASURING HEAD CIRCUMFERENCE | | 1 | 2 | 3 | | | |
| 05 | GROWTH CHARTS | | 1 | 2 | 3 | | | |
| 07 | TAPE FOR MID-UPPER ARM CIRCUMFERENCE (MUAC) | | 1 | 2 | 3 | | | |
| | YOUR RESPONDENT AND MOVE TO YOUR N | EXT D | ΑΤΑ Ο | OLLECTION I | POINT IF DIFF | ERENT F | ROM | CURRENT |

| SECTION 12 : | CHILD CURATIVE | CARE SERVICES |
|---------------------|-----------------------|----------------------|
|---------------------|-----------------------|----------------------|

| 1200 | CHECK Q102.03 | | | |
|-----------------|---|---|---------------------------------|---------------------|
| | CURATIVE CARE SERVICES | NO CURATIVE CARE SERVICES | | |
| | AVAILABLE | NEXT SERVICE SITE / 01253* | \mathbf{L} | |
| | + | | | |
| ASK TO PERSO | BE SHOWN THE LOCATIONS IN THE FACILITY WHE ON MOST KNOWLEDGEABLE ABOUT CURATIVE CA EXPLAIN THE PURPOSE OF THE SURVEY A | RE CURATIVE CARE SERVICES ARE RE SERVICES IN THE FACILITY. INT IND ASK THE FOLLOWING QUESTI | PROVIDED. RODUCE YOU ONS. | FIND THE JRSELF, |
| 1201 | Please tell me the number of days per month tha consultations or curative care for children under 5 are offered in this facility. | t (a) # OF DAYS PER MONTH SERVICE I | IS PROVIDED AT | FACILITY |
| | USE A 4-WEEK MONTH TO CALCULATE # OF DAYS | | | |
| 01 | Consultation or curative care services for sick children | # OF DAYS | | |
| 1202 | Please tell me if providers of child health serv following services | ices in this facility provide the | YES | NO |
| 01 | DIAGNOSE AND/OR TREAT CHILD MALNUTRITION | | 1 | 2 |
| 02 | PROVIDE VITAMIN A SUPPLEMENTATION TO CHILDREN | | 1 | 2 |
| 03 | PROVIDE IRON SUPPLEMENTATION TO CHILDREN | | 1 | 2 |
| 04 | PROVIDE ZINC SUPPLEMENTATION TO CHILDREN | | 1 | 2 |
| 05* | PROVIDE DEWORMING TO CHILDREN | | 1 | 2 |
| 06* | PROVIDE SEPSIS MANAGEMENT FOR CHILDREN | 1 | 2 | |
| 07 | PROVIDE ORAL ANTIBIOTIC TO CHILDREN | | | 2 |
| 08 | PROVIDE INJECATABLE ANTIBIOTIC TO CHILDREN | | 1 | 2 |
| 12 | PROVIDE SCANU SUPPORT TO SEVERELY ILL NEWBORN | | 1 | 2 |
| 13 [‡] | PROVIDE DIARROHEA MANAGEMENT TO CHILDREN | | 1 | 2 |
| 14 [‡] | PROVIDE PNEUMONIA MANAGEMENT TO CHILDREN | | 1 | 2 |
| 1203 | Do providers of services for sick children in this facility follow the IMCI guidelines in the provision of services to children under 5 years? | YES NO | 1 2 | |
| 1203A | Do this facility have separate IMCI corner? | YES NO | 1 2 | |
| 1204 | Do you have the <i>IMCI guidelines</i> (chart booklet) for the diagnosis and management of childhood illnesses available in this service area today? | YES NO | 1 2— | →1206 |
| 1205 | May I see the IMCI guidelines? | OBSERVED REPORTED, NOT SEEN | 1 2 | |
| 1206 | Do you have <i>any (other) guidelines</i> for the diagnosis and management of childhood illnesses available in this service site today? | YES NO | <u>1</u> 2— | →1208 |
| 1207 | May I see the other guidelines? | OBSERVED REPORTED, NOT SEEN | 1 2 | |
| 1208 | Does this facility have a system whereby certain observations and parameters are routinely carried out on sick children before the consultation for the presenting illness? | YES | | →1210 |
| 1207 1208 | illnesses available in this service site today? May I see the other guidelines? Does this facility have a system whereby certain observations and parameters are routinely carried out on sick children before the consultation for the presenting illness? IF YES, ASK TO SEE THE PLACE WHERE THESE ACTIVITIES | OBSERVED REPORTED, NOT SEEN YES NO | | - |

| | TAKE PLACE BEFORE THE CONSULTATION | | | | | | | |
|-----------------|--|-------|-----------|------------------|----------------------|------------------------|----------|---------------|
| 1209 | OBSERVE IF THE BELOW ACTIVITIES ARE BEING DONE ROUTINELY. IF YOU DO NOT SEE AN ACTI | VITY, | AC OBS | TIVITY SERVED | ACTIVITY REPORTED | ACTIVITY N CONDUCTE | OT ED | DON'T KNOW |
| | ASK: | | | | NOT SEEN | | | |
| | Is [ACTIVITY YOU DO NOT SEE] routinely | | | | | | | |
| | conducted for all sick children? | | | | | | | |
| 01 | Weighing the child | | | 1 | 2 | 3 | | 8 |
| 02 | Plotting child's weight on graph | | | 1 | 2 | 3 | | 8 |
| 03 | Taking child's temperature | | | 1 | 2 | 3 | | 8 |
| 04 | Assessing child's vaccination status | | | 1 | 2 | 3 | | 8 |
| 05 | Providing group health education | | | 1 | 2 | 3 | | 8 |
| 06 | Administer fever-reducing medicines and/or spo for fever | onge | | 1 | 2 | 3 | | 8 |
| 07 | Triaging of sick children, i.e., prioritizing sick chi based on the severity of their condition | ldren | | 1 | 2 | 3 | | 8 |
| 08 | Taking child's height/length measurement | | | 1 | 2 | 3 | | 8 |
| 09 | Taking MUAC measurement | | | 1 | 2 | 3 | | 8 |
| 1210 | I would like to know if the following items are | | (a) A\ | /AILABLE | | (b) | FUNC | TIONING |
| _ | available in this service area. I would like to | OBSE | RVED | REPORTED | D, NOT | YES | NO | DON'T |
| | see them. For equipment and instruments, I | | | NOT SEEM | N AVAILABLE | | | KNOW |
| | (Outpatient department) | | | | | | | |
| 01 | CHILD WEIGHING SCALE | 1_ | ►b | 2 b | 3 — | 1 | 2 | 8 |
| 01 | (250 GRAM GRADATION) | | | | 02 🚽 | _ | | |
| 02 | INFANT WEIGHING SCALE | 1- | ►b | 2 ⊳ b | 3 — | 1 | 2 | 8 |
| | (100 GRAM GRADATION) | | - | F | 03 ┥ | | | |
| 03 | THERMOMETER | 1_ | ►b | 2 - ▶ b | 3 04 4 | 1 | 2 | 8 |
| 04 | STETHOSCOPE | 1_ | ►b | 2 → b | 3 05 4 | 1 | 2 | 8 |
| 05 | Timer or watch with seconds hand | 1– | ►b | 2 → b | 3 06 4 | 1 | 2 | 8 |
| 06 | Staff has watch with seconds hand or other device (e.g., cell phone) that can measure seconds | | 1 | 2 | 3 | | | |
| 07 | Calibrated 1/2 or 1-liter measuring jar for ORS | | 1 | 2 | 3 | | | |
| 08 | Cup and spoon | | 1 | 2 | 3 | | | |
| 09 | ORS PACKETS OR SACHETS | | 1 | 2 | 3 | | | |
| 10 | At least 3 buckets (for cleaning used cups) | | 1 | 2 | 3 | | | |
| 11 | Examination bed or couch | | 1 | 2 | 3 | | | |
| 12 [‡] | Pulse oximeter | | 1 | 2 | 3 13 4 | 1 | 2 | 8 |
| 13 [‡] | Torches | | 1 | 2 | 3 14 | 1 | 2 | 8 |
| 14 [‡] | Measuring tape | | 1 | 2 | 3 | | | |
| 15 [‡] | Growth monitoring chart | | 1 | 2 | 3 | | | |
| 16 [‡] | Height measuring board | | 1 | 2 | 3 | | | |
| 17 [‡] | Tape for Mid-upper Arm Circumference (MUAC) | | 1 | 2 | 3 | | | |

| 18 [‡] | Tongue depress | or (wooden/metallic) | 1 | 2 | 3 | | | | |
|--|--|---|--|--|---|--|---|--|--|
| `19 [‡] | Syringe (5ml) | | 1 | 2 | 3 | | | | |
| 20 [‡] | Insulin syringe | | 1 | 2 | 3 | | | | |
| 21 [‡] | Alcohol based h | and rub | 1 | 2 | 3 | | | | |
| 22 [‡] | Cotton ball | | 1 | 2 | 3 | | | | |
| 23 [‡] | Disposable latex | k/other gloves | 1 | 2 | 3 | | | | |
| 1211 | Please tell me materials. IF YES, ASK TO | if you have any of the following | | | | | | | |
| 01 | IMCI chart book | let | 1 | 2 | 3 | | | | |
| 02 | IMCI mother's c | ards (IMCI card) | 1 | 2 | 3 | | | | |
| 03 | Other visual aid | s for teaching caretakers | 1 | 2 | 3 | | | | |
| 04* | IMCI register | | 1 | 2 | 3 | | | | |
| 05* | Sick newborn ca | are register | 1 | 2 | 3 | | | | |
| 1212 | Are individua card or book this service s | al health records (i.e., child we let) for sick children maintaine ite? | fare YES. d at NO . | | | | 1 2 — • | ► 1214 [‡] | |
| 1213 | May I see an records? | unused copy of the individual | YES. NO . | | | | 1 2 | | |
| | | Inpatient department YES (Inpatient department) | | | NO (Inpatient | departme | ent) | | |
| | | | Ļ | | | 2.2200 | | ▲ | |
| 1214A [‡] | l would like to | know if the following items are | (a) A | VAILABLE | | (b) | FUNCT | | |
| 1214A [‡] | I would like to available in th see them. For would like to k (Inpatient dep | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. partment) | (a) A OBSERVED | VAILABLE REPORTED, NOT SEEN | NOT AVAILABLE | (b) YES | FUNCT | IONING DON'T KNOW | |
| 1214A [‡] | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. bartment) IG SCALE (250GRAM GRADATION) | (a) A OBSERVED 1→b | VAILABLE REPORTED, NOT SEEN 2-b | NOT AVAILABLE | (b) YES | FUNCT NO 2 | IONING DON'T KNOW 8 | |
| 1214A [‡] 01 02 | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHI | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. bartment) IG SCALE (250GRAM GRADATION) | (a) A OBSERVED 1→b 1→b | VAILABLE REPORTED, NOT SEEN 2- b 2- b | NOT AVAILABLE | (b) YES 1 | FUNCT NO 2 | IONING DON'T KNOW 8 8 | |
| 1214A [‡] 01 02 03 | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHI THERMOMETER | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. bartment) IG SCALE (250GRAM GRADATION) NG SCALE (100 GRAM GRADATION) | (a) A OBSERVED 1→b 1→b 1→b | VAILABLE REPORTED, NOT SEEN 2- b 2- b 2- b | NOT AVAILABLE 3 02 3 03 3 03 4 | (b) YES 1 1 | FUNCT NO 2 2 2 | IONING DON'T KNOW 8 8 8 8 | |
| 1214A [‡] 01 02 03 04 | l would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHI THERMOMETER STETHOSCOPE | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. hartment) IG SCALE (250GRAM GRADATION) ING SCALE (100 GRAM GRADATION) | (a) A OBSERVED 1→b 1→b 1→b 1→b | VAILABLE REPORTED, NOT SEEN 2 b 2 b 2 b 2 b 2 b 2 b | NOT AVAILABLE $3 \\ 02 \\ 3 \\ 03 \\ 4 \\ 3 \\ 04 \\ 4 \\ 3 \\ 05 \\ 4 \end{bmatrix}$ | (b) YES 1 1 1 1 | FUNCT NO 2 2 2 2 2 2 | IONING DON'T KNOW 8 8 8 8 8 8 | |
| 1214A [‡] 01 02 03 04 05 | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHI THERMOMETER STETHOSCOPE Timer or watch | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. bartment) IG SCALE (250GRAM GRADATION) ING SCALE (100 GRAM GRADATION) | (a) A OBSERVED $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ | VAILABLE REPORTED, NOT SEEN 2→ b 2→ b 2→ b 2→ b 2→ b 2→ b | NOT AVAILABLE 3 02 3 03 4 3 04 4 3 05 4 3 05 4 | (b) YES 1 1 1 1 1 1 | FUNCT NO 2 2 2 2 2 2 2 2 2 2 | IONING DON'T KNOW 8 8 8 8 8 8 8 8 8 | |
| 1214A [‡] 01 02 03 04 05 06 | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHIN THERMOMETER STETHOSCOPE Timer or watch Nebulizer | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. bartment) G SCALE (250GRAM GRADATION) NG SCALE (100 GRAM GRADATION) | (a) A OBSERVED $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ $1 \rightarrow b$ | VAILABLE REPORTED, NOT SEEN 2→ b 2→ b 2→ b 2→ b 2→ b 2→ b | NOT AVAILABLE $3 \\ 02 \\ 3 \\ 03 \\ 4 \\ 3 \\ 04 \\ 4 \\ 3 \\ 05 \\ 4 \\ 3 \\ 05 \\ 4 \\ 3 \\ 06 \\ 4 \\ 3 \\ 07 \\ 4 \end{bmatrix}$ | (b) YES 1 1 1 1 1 1 1 | FUNCT NO 2 2 2 2 2 2 2 2 2 2 2 2 2 | IONING DON'T KNOW 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 | |
| 1214A [‡] 01 02 03 04 05 06 07 | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHI THERMOMETER STETHOSCOPE Timer or watch Nebulizer Spacers for inha | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. hartment) IG SCALE (250GRAM GRADATION) ING SCALE (100 GRAM GRADATION) R with seconds hand | (a) A OBSERVED $1 \rightarrow b$ $1 \rightarrow b$ | VAILABLE REPORTED, NOT SEEN 2→ b 2→ b 2→ b 2→ b 2→ b 2→ b 2→ b 2→ b | NOT AVAILABLE $3 \\ 02 \\ 3 \\ 03 \\ 4 \\ 3 \\ 04 \\ 4 \\ 3 \\ 05 \\ 4 \\ 3 \\ 06 \\ 4 \\ 3 \\ 06 \\ 4 \\ 3 \\ 06 \\ 4 \\ 3 \\ 07 \\ 4 \\ 3 \\ 08 \\ 4 \end{bmatrix}$ | (b) YES 1 1 1 1 1 1 1 1 1 | FUNCT NO 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | IONING DON'T KNOW 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 | |
| 1214A [‡] 01 02 03 04 05 06 07 08 | I would like to available in th see them. For would like to k (Inpatient dep CHILD WEIGHIN INFANT WEIGHI THERMOMETER STETHOSCOPE Timer or watch Nebulizer Spacers for inha Peak flow meter | know if the following items are is service area. I would like to equipment and instruments, I know if they are functioning. bartment) IG SCALE (250GRAM GRADATION) ING SCALE (100 GRAM GRADATION) R with seconds hand | (a) A OBSERVED $1 \rightarrow b$ $1 \rightarrow b$ | VAILABLE REPORTED, NOT SEEN 2 \rightarrow b 2 \rightarrow b | NOT AVAILABLE 3 02 3 03 4 3 04 4 3 05 4 3 06 4 3 06 4 3 07 4 3 08 4 3 09 4 | (b) YES 1 1 1 1 1 1 1 1 1 1 1 1 | FUNCT NO 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | IONING DON'T KNOW 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 | |

| 10 | FILLED OXYGEN CYLINDER WITH FLOW METER | 1 b | 2_ b | 3 11 4 | 1 | 2 | 8 |
|--|--|--|---|--|---|---|---|
| 11 | FILLED OXYGEN CYLINDER WITHOUT FLOW METER | 1 b | 2 -> b | 3 12 | 1 | 2 | 8 |
| 12 | OXYGEN DISTRIBUTION SYSTEM | 1 b | 2 > b | 3 13 ↓ | 1 | 2 | 8 |
| 13 | RADIANT WARMER | 1 _ ▶b | 2 _ b | 3 14 4 | 1 | 2 | 8 |
| 14 | Staff has watch with seconds hand or other device (e.g., cell phone) that can measure seconds | 1 | 2 | 3 | | | |
| 15 | Calibrated 1/2 or 1-liter measuring jar for ORS | 1 | 2 | 3 | | | |
| 16 | Cup and spoon | 1 | 2 | 3 | | | |
| 17 | ORS PACKETS OR SACHETS | 1 | 2 | 3 | | | |
| 18 | At least 3 buckets (for cleaning used cups) | 1 | 2 | 3 | | | |
| 20 | Examination bed or couch | 1 | 2 | 3 | | | |
| 21 | DIGITAL BP APPARATUS | 1 b | 2- > b | 3 22 | 1 | 2 | 8 |
| 22 | MANUAL BP APPARATUS | 1 b | 2- > b | 3 23 4 | 1 | 2 | 8 |
| 23 | Pulse oximeter | 1 b | 2 b | 3 24 | 1 | 2 | 8 |
| 24 | Diagnostic set or otoscope | 1 _ ▶b | 2 b | 3 25 | 1 | 2 | 8 |
| 25 | Vision testing chart | 1 | 2 | 3 | | | |
| | | | | | | | |
| 26 | Torches | 1 - ▶b | 2 - ▶ b | 3 28 | 1 | 2 | 8 |
| 26 27 | Torches Measuring tape | 1- → b 1 | 2→ b 2 | 3 28 ◀ 3 | 1 | 2 | 8 |
| 26 27 28 | Torches Measuring tape Growth monitoring chart | 1-→b 1 1 | 2→ b 2 2 | 3 | 1 | 2 | 8 |
| 26 27 28 29 | Torches Measuring tape Growth monitoring chart Height measuring board | 1→b 1 1 | 2→ b 2 2 2 | 3 28 4 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand | 1→b 1 1 1 1 | 2→ b 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) | 1→b 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 32 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe | 1→b 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 4 3 3 3 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 32 33 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids | 1→b 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 32 33 34 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution | 1→b 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 32 33 33 34 35 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub | 1→b 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 31 32 33 34 35 36 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 32 33 33 34 35 36 37 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | | 2 | 8 |
| 26 27 28 29 30 31 32 33 34 35 36 37 38 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube Disposable latex/other gloves | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | 1 | 2 | 8 |
| 26 27 28 29 30 31 32 33 33 34 35 36 37 38 39 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube Disposable latex/other gloves Micropore tape | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | | 2 | 8 |
| 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube Disposable latex/other gloves Micropore tape Gauze | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | | 2 | 8 |
| 26 27 28 29 30 31 32 33 33 34 35 36 37 38 39 40 41 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube Disposable latex/other gloves Micropore tape Gauze Nasal catheter | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | | 2 | 8 |
| 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube Disposable latex/other gloves Micropore tape Gauze Nasal catheter Nasal cannula | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | | 2 | 8 |
| 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 1214B [‡] | Torches Measuring tape Growth monitoring chart Height measuring board Drip Stand Syringe (5ml) Insulin syringe Cannula for administering IV fluids Infusion set for IV solution Alcohol based hand rub Cotton ball Nasogastric tube Disposable latex/other gloves Micropore tape Gauze Nasal catheter Nasal cannula Please tell me if you have any of the following materials. | 1→b 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2→ b 2 2 2 2 2 2 2 2 2 2 2 2 2 | 3 28 ◀ 3 3 3 3 3 3 3 3 3 3 3 3 3 | | 2 | 8 |

| 02 | NSU/SCANU register | 1 | 2 | 3 |
|----|-------------------------------|---|---|---|
| 03 | Paediatric inpatient register | 1 | 2 | 3 |

| 1250 | ASSESS THE ROOM OR AREA FOR THE ITEMS LISTED BELOW. FOR ITEMS THAT YOU DO NOT SEE, ASK YOUR RESPONDENT TO SHOW THEM TO YOU. IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECORDED | GENERAL INFOI CHILD VACCINA FAMILY PLANNI ANTENATAL CA DELIVERY [Q16: TUBERCULOSIS NCD [Q2351] NOT PREVIOUS | RMATION [Q710 ITION [Q1051] ING [Q1351] RE [Q1451] 51] [Q1951] LY SEEN | 'NEXT SECTION / SERVICE SITE | |
|------|--|--|--|------------------------------------|------------------|
| 1251 | STANDARD PRECAUTIONS AND CONDITIONS EXAMINATION | OBSERVED | REPORTED NOT SEEN | NOT AVAILABLE | |
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POL | IR PITCHER) | 1 | 2 | 3 |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOA | P) | 1 | 2 | 3 |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLAS | 1 | 2 | 3 | |
| 05 | OTHER WASTE RECEPTACLE | | 1 | 2 | 3 |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITA | NE, ALCOHOL] | 1 | 2 | 3 |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH AUTO-DISABLE SYRINGES WITH NEEDLES | H NEEDLES OR | 1 | 2 | 3 |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 |
| 11 | GOWNS | | 1 | 2 | 3 |
| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 |
| 1252 | DESCRIBE THE SETTING OF THE SICK CHILD SERVICE PRIVATE ROOM OTHER ROOM V DELIVERY ROOM OR AREA. OTHER ROOM V PRIVACY | | 1 WITH AUDITORY Y ONLY | AND VISUAL | - 2 3 4 |

ADOLESCENT HEALTH SERVICES

| 1253* | CHECK Q102.22 | | | | | F | - | |
|--------------------|---|-----------------------|----------|---|-----------------------------|---------------------|----------|-------------------|
| | ADOLESCENT HEALTH SERVICE PROVIDED | | | ADOLE NOT F | SCENT HEALTH SE PROVIDED | RVICE | | →1261 |
| 1254* | Is there a dedicated room/area for | • adolescent | | YES | | | 1 | |
| | health service? | | | NO 2— | | | | ▶1256 |
| 1255 | DESCRIBE THE SETTING OF THE ADOLE | SCENT HEALTH | l | PRIVATE RO | DOM | | 1 | |
| | SERVICE ROOM OR AREA. | | | OTHER ROOM WITH AUDITORY AND VISUAL | | | | |
| | | | | PRIVACY | | | 2 | |
| | | | | VISUAL PRI | VACY ONLY | | 3 | |
| 1250 | How many days each week are adoles | cent health | | | <u>Y</u> | ······ | 4 | |
| 1256 | service available at this facility? | | | | Ι ΤΗΛΝ ΟΝCΕ/Μ | | | |
| 1257 | Do you have the adolescent health gui | delines/manua | ı | VES | | | . 0 1 | |
| 1237 | available in this service area today? | | | NO | | | - 2 | → ₁₂₅₉ |
| 1258 | May I see the adolescent guidelines/m | ianual? | | | | | - 1 | →1260a |
| 1250 | ACCEPTABLE IF PART OF OTHER GUIDE | LINES | | REPORTED | NOT SEEN | | 2 | |
| 1259 | Do you have any other adolescent hea | lth | | YES | | | - | |
| | guidelines/job aid available in this serv | vice area today | ? | NO | | | 2 | → 1260a |
| 1260 | May I see the other guidelines/job aid | ? | | OBSERVED | | | 1 | |
| | | | | REPORTED, NOT SEEN | | | | |
| 1260a‡ | Does this corner have sitting arrangem | nent for | | YES 1 | | | | |
| | adolescent? | | | NO | | | 2 | |
| 1260b [‡] | Does this corner have safe drinking wa | ater for | | YES1 | | | | |
| | adolescent? | | | NO | | | 2 | |
| 1260c [‡] | Does this corner have 7 adolescent he | alth related | | YES | | | 1 | |
| | drugs? | | | NO | | | 2 | ▶1016 |
| 1260d [‡] | Are any of the following adolescent | (A) OB | SEF | RVED | (E | 3) NOT OBSERVE | ED | |
| | health related drugs available in the location today? | AVAI | LAE | BLE | | | r | |
| | | AT LEAST ONE VALID | A\ N(| VAILABLE ONE VALID | REPORTED AVAILABLE | NOT AVAILABLE/TO | NE | VER |
| | CHECK TO SEE IF AT LEAST ONE IS | 0112 171210 | | | NOT SEEN | DAY/DK | A١ | AILABLE |
| | VALID (NOT EXPIRED) | | | | | | | |
| 01 | Fumarate 200mg+Folic acid (Ferrous | 1 | | 2 | 3 | 4 | | 5 |
| 02 | Cap. Doxycycline (100 mg) | 1 | | 2 | 3 | 4 | | 5 |
| 03 | Tab. Metronidazole 400 mg | 1 | | 2 | 3 | 4 | | 5 |
| 04* | Tab. Ciprofloxacin (500 mg) | 1 | | 2 | 3 | 4 | | 5 |
| 05* | Tab. Naproxen (250 mg) | 1 | | 2 | 3 | 4 | | 5 |
| 06* | Tab. Pantoprazole (20 mg) | 1 | | 2 | 3 | 4 | | 5 |
| 07 | Tab. Albendazole | 1 | | 2 | 3 | 4 | | 5 |
| 1016 | Now I would like to ask you specific | ally about vac | cin | ation servic | es for adolesce | ent. For each of | the | following |
| | services, please tell me whether th | e service is of | ter | ed by your f | acility, and if s | o, how many da | iys p | er month |
| | ADOLESCENT VACCINATION SERVI | CE | | | | (a) | | |
| | USE A 4-WEEK MONTH TO CALCULAT | E # OF DAYS | | # OF DAYS PER MONTH SERVICE IS PROVIDED AT FACILITY | | | | |

| 01 | TT vaccine | # OF DAY 00=NO SI | ′S ERVICE | | | |
|-------|---|---|---|-----------------------------------|------------------------------|--------------------|
| 1017* | Do you have the national guidelines for adolescent vaccinations available in this service area today? | YES NO | | | 1 2— | → 1019 |
| 1018* | May I see the guidelines? | OBSERVE REPORTE | D D, NOT SEEN. | | 1 — 2 | → 1021* |
| 1019* | Do you have any other guidelines for adolescent vaccinations available in this service area today? | YES NO | | | 1 2— | ➡1021* |
| 1020* | May I see the other guidelines? | OBSERVE REPORTE | D D, NOT SEEN. | | 1 2 | |
| 1021* | Please tell me if each of the following vaccines is available in the facility today. If available, I | (A) O AVA | BSERVED | (B) | NOT OBSER | VED |
| | would like to see it. IF AVAILABLE, CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED, VACCINE VIAL MONITOR UNCHANGED (NOT FROZEN)) | ATLEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NEVER AVAILABLE |
| 01 | TT vaccine | 1 | 2 | 3 | 4 | 5 |
| | NUTRITION | SERVI | CES | | | |
| 1261* | CHECK Q102.23 NUTRITION SERVICES PROVIDED | NUTR | ITION SERVICES | NOT PROVIDE | D | →1300 |
| 1262* | Is there a dedicated room/area for nutrition services? | YES NO | | | 1 2- | 1264 |
| 1263* | DESCRIBE THE SETTING OF THE NUTRITION SERVICE ROOM OR AREA. | PRIVAT OTHER PRIVAC VISUAL NO PRI | re Room Room with Cy Privacy oni | AUDITORY A | 1 ND VISUAL 2 3 | |
| 1264* | How many days each week are nutrition service available at this facility? | # OF D. LESS O | AYS FTEN THAN O | NCE/WEEK | 0 | |
| 1265* | Do you have the SAM/MAM guidelines available in this service area today? | ⁵ YES NO | | | 1 2 ⁻ | ► 1267 |
| 1266* | May I see the SAM/MAM guidelines? ACCEPTABLE IF PART OF OTHER GUIDELINES | OBSER REPOR | VED TED, NOT SEE | N | | 1300 |
| 1267* | Do you have any other nutrition guidelines available in this service area today? | YES NO | | | | → 1300 |
| 1268* | May I see the other nutrition guidelines/job aid? | OBSER | VED | N | | |
| | THANK YOUR RESPONDENT AND MOVE TO YOUR NE LOCATION. | EXT DATA | COLLECTION | POINT IF DIF | FERENT FROM | / CURRENT |

SECTION 13: FAMILY PLANNING

| 1300 | CHECK Q102.04 | | | | |
|------|--------------------------|---|-----------------------------|---|---|
| | FAMILY PLANNING SERVICES | | NO FAMILY PLANNING SERVICES | | |
| | | | | • | |
| | , | 7 | | | _ |

| | | NEXT SECTIO | ON OR SERVICE SITE | <u>-</u> | | | | |
|--|---|---|--------------------|-----------------------------|--------|--|--|--|
| ASK TO BE SHOWN THE LOCATION IN THE FACILITY WHERE FAMILY PLANING SERVICES ARE PROVIDED. FIND THE PE MOST KNOWLEDGEABLE ABOUT FAMILY PLANNING SERVICES IN THE FACILITY. INTRODUCE YOURSELF, EXPLAIN THE P OF THE SURVEY AND ASK THE FOLLOWING QUESTIONS. | | | | | | | | |
| 1301 | How many days in a month are family planning services offered at this facility? USE A 4-WEEK MONTH TO CALCULATE # OF DAYS | # OF DAYS | # OF DAYS | | | | | |
| 1302 | Does this facility provide (i.e., stock the commodit counsel or refer clients for any of the following mod family planning? | bes this facility provide (i.e., stock the commodity) or <i>prescribe, unsel or refer clients for</i> any of the following modern methods of mily planning? | | | NO | | | |
| 01 | COMBINED ORAL CONTRACEPTIVE PILLS | | 1 | 2 | 3 | | | |
| 02 | PROGESTIN-ONLY CONTRACEPTIVE PILLS | | 1 | 2 | 3 | | | |
| 04 | PROGESTIN-ONLY INJECTABLE CONTRACEPTIVES (D | EPO) | 1 | 2 | 3 | | | |
| 05 | MALE CONDOMS | | 1 | 2 | 3 | | | |
| 07 | INTRAUTERINE CONTRACEPTIVE DEVICE (IUCD) / Cu | ı-T- 380A | 1 | 2 | 3 | | | |
| 08 | ONE ROD IMPLANT | | 1 | 2 | 3 | | | |
| 09 | TWO RODS IMPLANT | | 1 | 2 | 3 | | | |
| 10 | EMERGENCY CONTRACEPTIVE PILLS (ECP) (e.g., PRC | OSTINOL 2) | 1 | 2 | 3 | | | |
| 12 | No Scalpel VASECTOMY (NSV) (MALE STERILIZATION | 1 | 2 | 3 | | | | |
| 13 | TUBAL LIGATION /Tubectomy (EEMALE STERILIZATI | 1 | 2 | 2 | | | | |
| 15 | COUNSEL CLIENTS ON LACTATIONAL AMENORRHEA | A METHOD | - | 2 | 3 | | | |
| 1303 | Do you have the <i>family planning guidelines/</i> | YES | | | ▶ 1305 | | | |
| 1304 | May I see the family planning guidelines/manual? | OBSERVED | SEEN | 1 | ▶ 1305 | | | |
| 1304a‡ | Please observe the last publishing year of the family planning guideline/manual | Year | | | ▶1307 | | | |
| 1305 | Do you have any other guidelines/instructions/ job aid/checklist on family planning available at this service area today? | YES NO | | 1 2— | ▶1307 | | | |
| 1306 | May I see the other guidelines/instructions/ job aid/ checklist? (e.g., IUCD card, IMPLANT card, Sthayi Paddhati Grahita Form consent form, admission form, payment record, admission register etc.,) | OBSERVED | | | | | | |
| 1307 | Are individual records or cards (e.g., IUCD card, IMPLANT card, Sthayi Paddhati Grahita Form etc.,) maintained at this service site for family planning clients? | YES | | | ▶1309 | | | |
| 1308 | May I see a blank copy of the individual records or card? | OBSERVED REPORTED, NOT | SEEN | 1 2 — | ▶1309 | | | |
| 1308a‡ | List IUCD Card Implant Card Injectable Client Card Sthayi Paddhati Grahita Form (Male/female) | YES 1 1 1 1 1 1 1 | | NO 2 2 2 2 2 | | | | |
| 1309 | Does this facility have a system whereby certain observations and parameters are routinely carried out on family planning clients before the | YES NO | | 1 | ▶1311 | | | |

| | consultation takes place? IF YES, ASK TO SEE THE PLACE WHERE THESE ACTIVITIES | | | | |
|------|--|---|----------------------------------|----------------------|---------------|
| 1310 | OBSERVE IF THE BELOW ACTIVITIES ARE BEING DONE ROUTINELY. IF YOU DO NOT SEE AN ACTIVITY, ASK: Is [ACTIVITY YOU DO NOT SEE] routinely done for all family planning clients? | ACTIVITY OBSERVED | ACTIVITY REPORTED NOT SEEN | ACTIVITY NOT DONE | DON'T KNOW |
| 01 | Weighing of clients | 1 | 2 | 3 | 8 |
| 02 | Taking blood pressure | 1 | 2 | 3 | 8 |
| 03 | Conducting group health education sessions | 1 | 2 | 3 | 8 |
| 1311 | Do family planning providers in this facility routinely diagnose and treat RTIs/STIs, or are RTIs/STIs clients referred to another provider or location for STI diagnosis and treatment? PROBE TO ARRIVE AT THE RIGHT ANSWER | ROUTINELY DIAG DIAGNOSE BUT FOR TREATMEN REFER ELSEWHER TREATMENT REFER OUTSIDE F TREATMENT NO DIAGNOSIS/T | | | |

EQUIPMENT AND SUPPLIES

| 121/ | I would like to know if the following items are | (a |) AV | 'AILABLE? | | (b) | FUNCTI | ONING |
|------|---|----------|-------|----------------------|-----------------------------|-----------|--------|---------------|
| 1314 | available in this service area today and are functioning | OBSERV | 'ED | REPORTED NOT SEEN | 9, NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01 | DIGITAL BP APPARATUS | 1 | b | 2 - ▶ b | 3 02 | 1 | 2 | 8 |
| 02 | MANUAL BP APPARATUS | 1 | b | 2 -→ b | 3 03 4 | 1 | 2 | 8 |
| 03 | STETHOSCOPE | 1 | b | 2 → b | 3 04 4 | 1 | 2 | 8 |
| 04 | EXAMINATION LIGHT/ SPOT LIGHT | 1 | b | 2 -→ b | 3 05 4 | 1 | 2 | 8 |
| 05 | EXAMINATION BED OR COUCH | 1→ | b | 2 → b | 3 | | | |
| 06 | SAMPLE OF FP METHODS/ COUNSELLING KIT | 1 | | 2 | 3 | 1 | | |
| 07 | OTHER FP-SPECIFIC VISUAL AIDS [e.g., FLIP CHARTS, LEAFLETS] | 1 | | 2 | 3 | | | |
| 08 | PELVIC MODEL FOR IUCD COUNSELLING | 1 | | 2 | 3 | 7 | | |
| 09 | MODEL FOR SHOWING CONDOM USE | 1 | | 2 | 3 | | | |
| 1315 | CHECK Q1302.07 & Q1302.08. &09 IUCD OR IMPLANT PROVIDED IN FACILITY | | NEIT | HER IUCD N | IOR IMPLANT PRO\ ACILITY | /IDED | | 1 321 |
| | ASK TO BE TAKEN TO THE ROOM OR LOCATION W | HERE IUC | Ds AN | D/OR IMPL | ANTS ARE INSERTE | D OR REMO | VED | |
| 1316 | Please show me the following items for the provi of IUCD or Implant methods: | ision | OB | SERVED | REPORTED, NOT SEEN | NOT AVAIL | ABLE | |
| 01 | STERILE GLOVES | | | 1 | 2 | 3 | | |
| 02 | ANTISEPTIC SOLUTION (E.G., POVIDON IODINE) | | | 1 | 2 | 3 | | |
| 03 | SPONGE HOLDING FORCEPS | | | 1 | 2 | 3 | | |
| 04 | STERILE GAUZE PAD OR COTTON WOOL | | | 1 | 2 | 3 | | |
| 05 | GALLIPOT (FOR ANTISEPTIC SOLUTION) | | | 1 | 2 | 3 | | |

| 1317 | CHECK Q1302.07 IUCD PROVIDED IN FACILITY | IUCD N | OT PROVIDED IN F | | → 1319 |
|------|--|---|---|---------------|----------------|
| 1318 | Please show me the following items for the provision of IUCD: | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | |
| 01 | CUSCO VAGINAL SPECULUM - SMALL | 1 | 2 | 3 | |
| 02 | CUSCO VAGINAL SPECULUM - MEDIUM | 1 | 2 | 3 | |
| 03 | CUSCO VAGINAL SPECULUM - LARGE | 1 | 2 | 3 | |
| 04 | TENACULAM/ VOLSELLUM FORCEPS | 1 | 2 | 3 | |
| 05 | UTERINE SOUND | 1 | 2 | 3 | |
| 06 | STRAIGHT ARTERY FORCEPS (LONG) | 1 | 2 | 3 | |
| 07 | STRAIGHT CUTTING SCISSORS | 1 | 2 | 3 | |
| 08 | HIGH STOOL FOR SITTING | 1 | 2 | 3 | |
| 09 | 0.5% CHLORINE SOLUTION IN RED BUCKET WITH LID | 1 | 2 | 3 | |
| 10 | BLUE BUCKET (FOR WASTE DISPOSAL) | 1 | 2 | 3 | |
| 11 | IUCD IN STERILE PACKAGING | 1 | 2 | 3 | |
| 12 | KELLY FORCEPS for PPIUCD | 1 | 2 | 3 | |
| 1319 | CHECK Q 1302.08, 1302.09 IMPLANT PROVIDED IN FACILITY | IMPLANT N | OT PROVIDED IN FA | | →1321 |
| 1320 | Please show me the following items for the provision of Implant: | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | |
| 01 | LOCAL ANESTHETIC (e.g., 1% LIDOCAINE) | 1 | 2 | 3 | |
| 02 | STERILE SYRINGE AND NEEDLE | 1 | 2 | 3 | |
| 03 | CANNULA AND TROCHAR FOR INSERTING IMPLANT | 1 | 2 | 3 | |
| 04 | SEALED IMPLANT PACK | 1 | 2 | 3 | |
| 05 | SCALPEL WITH BLADE (SURGICAL BLADE WITH HANDLE) | 1 | 2 | 3 | |
| 07 | ARM REST / SIDE TABLE | 1 | 2 | 3 | |
| 08 | MARKER PEN | 1 | 2 | 3 | |
| 09 | SURGICAL DRAPE | 1 | 2 | 3 | |
| 10 | NORMAL BANDAGE OR BUTTERFLY BANDAGE | 1 | 2 | 3 | |
| 11 | BAND AID | 1 | 2 | 3 | |
| 12 | ELASTOMETRIC MATTRESS DRESSING | 1 | 2 | 3 | |
| 1321 | Where are equipment such as speculam or forceps that are used in the provision of family planning services processed for re-use? | FP SERVICE SITE . CENTRAL LOCATI BOTH LOCATIONS NO EQUIPMENT | ON IN FACILITY S PROCESSED IN FACILIT | | →1350 →1350 |
| 1322 | What is the final processing method used for family planning equipment at this service site? PROBE FOR ALL METHODS USED | AUTOCLAVE DRY HEAT STEF BOIL OR STEAN WASH WITH SC | RILIZATION 1 DAP AND WATER | A | |
| | | SOAK IN OTHER | R CHEMICAL SOLUT | IONE | |

| 1350 | ASSESS THE ROOM OR AREA FOR THE ITEMS LISTED BELOW. FOR ITEMS THAT YOU DO NOT SEE, ASK YOUR RESPONDENT TO SHOW THEM TO YOU. IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECORDED | GENERAL INFORMATION [Q710] 11 CHILD VACCINATION [Q1051] 12 CHILD CURATIVE CARE [Q1251] 13 ANTENATAL CARE [Q1451] 15 DELIVERY [Q1651] 17 TUBERCULOSIS [Q1951] 19 NCD [Q2351] 22 NOT PREVIOUSLY SEEN 31 | 'NEXT SECTION / ERVICE SIT/ 1353 |
|------|--|---|---|
| 1351 | STANDARD PRECAUTIONS AND CONDITIONS | FOR CLIENT OBSERVED REPORTED NC |)/ NOT |
| | | | |

| | EXAMINATION | | | NOT SEEN | AVAILABLE | |
|------|---|---|--------------------------------------|---------------|-----------|--|
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POL | JR PITCHER) | 1 | 2 | 3 | |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOA | 1 | 2 | 3 | | |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 | |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLAS | STIC BIN LINER | 1 | 2 | 3 | |
| 05 | OTHER WASTE RECEPTACLE | | 1 | 2 | 3 | |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 | |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 | |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITA | 1 | 2 | 3 | | |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH AUTO-DISABLE SYRINGES WITH NEEDLES | H NEEDLES OR | 1 | 2 | 3 | |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 | |
| 11 | GOWNS | | 1 | 2 | 3 | |
| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 | |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 | |
| 1352 | DESCRIBE THE SETTING OF THE FP SERVICE ROOM OR AREA. | PRIVATE ROOM OTHER ROOM V PRIVACY VISUAL PRIVACY | i WITH AUDITORY Y ONLY | | | |
| 1353 | CHECK Q212 FP COMMODITIES STORED IN OTHER LOCATION OR NOT STOCKED (RESPONSE 1 NOT CIRCLED) | 'FP COMMO 'SERVICE AREA | DITIES STORED I A (RESPONSE 1 CIR | N FP CLED) | 921 | |

| | No Scalpel VASECTOMY (NSV) AND TUBAL LIGATION (TUBECTOMY) | | | | | | | | | |
|-------------------|--|--|--------|--|--|--|--|--|--|--|
| 1353A | CHECK Q1302.12 NSV PROVIDED IN FACILITY | NSV NOT PROVIDED IN FACILITY | ►1353F | | | | | | | |
| ASK TO B NSV S | ASK TO BE SHOWN THE LOCATION IN THE FACILITY WHERE NSV SERVICES ARE PROVIDED. FIND THE PERSON MOST KNOWLEDGEABLE ABOUT NSV SERVICES IN THE FACILITY. INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE SURVEY AND ASK THE FOLLOWING QUESTIONS. | | | | | | | | | |
| 1353B | B I would like to know if the following items are available in this service area today. For some of the items, I would like to know if they are functioning. If any of the items are maintained elsewhere in the facility, please tell me where in the facility they are kept. | | | | | | | | | |
| | ASSESS THE ROOM OR AREA FOR THE ITEMS BELOW. FOR ITEMS THAT YOU DO NOT SEE, YOUR RESPONDENT TO SHOW THEM TO YOU | S LISTED ROOM / AREA ASSESSED IN Q1353H 11 + 1 , ASK ROOM/AREA NOT PREVIOUSLY DU. ASSESSED | 1353D | | | | | | | |
| | IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECO NOT PREVIOUSLY ASSESSED, CIRCLE "31" AND ASS SERVICE SITE. | N DRDED. IF ISESS THIS | | | | | | | | |

| 13530 | | (a) AV | /AILABLE? | | (b) | FUNCTI | ONING |
|-------|--|---------------|-----------------------|------------------|-----|--------|---------------|
| 15550 | | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01* | OPERATING THEATRE TABLE | 1 b | 2 _ ▶ b | 3 02 | 1 | 2 | 8 |
| 02* | OPERATING THEATRE LIGHT | 1 _ ▶b | 2 b | 3 03 ◀ | 1 | 2 | 8 |
| 03* | INSTRUMENT TROLLEY | 1 b | 2 b | 3 04 4 | 1 | 2 | 8 |
| 04* | STERILIZER DRUM | 1 b | 2_ b | 3 05 4 | 1 | 2 | 8 |
| 05* | BIG CURVE SCISSORS FOR CUTTING GAUZE OR BANDAGE | 1 b | 2 b | 3 06 | 1 | 2 | 8 |
| 06* | SCISSORS FOR CUTTING CLOTH | 1 b | 2 b | 3 07 | 1 | 2 | 8 |
| 07* | DIGITAL BP APPARATUS | 1 b | 2_ b | 3 09 4 | 1 | 2 | 8 |
| 09* | MANUAL BP APPARATUS | 1- - b | 2 b | 3 10 ◀ | 1 | 2 | 8 |
| 10* | STETHOSCOPE | 1 b | 2_ b | 3 11 4 | 1 | 2 | 8 |
| 11* | WEIGHING SCALE (ADULT) | 1 b | 2_ b | 3 12 | 1 | 2 | 8 |
| 12* | GALLIPOT OR CUP FOR ANTISEPTIC SOLUTION | 1 | 2 | 3 | | | |
| 13* | KIDNEY TRAY | 1 | 2 | 3 | | | |
| 15* | LIFTER AND LIFTER JAR | 1 | 2 | 3 | | | |
| 16* | DRESSING JAR WITH LID/COVER | 1 | 2 | 3 | | | |
| 18* | TALQUEST BOOK | 1 | 2 | 3 | | | |
| 19* | IRON COT | 1 | 2 | 3 | | | |
| 20* | THERMOMETER | 1 b | 2 b | 3 21 | 1 | 2 | 8 |
| 21* | GOWN FOR CLIENTS | 1 | 2 | 3 | | | |
| 22* | GOWN FOR PROVIDER | 1 | 2 | 3 | | | |
| 23* | TROLLEY SHEET | 1 | 2 | 3 | | | |
| 24* | DRAW SHEET | 1 | 2 | 3 | | | |
| 25* | САР | 1 | 2 | 3 | | | |
| 26* | MASK | 1 | 2 | 3 | | | |
| 34* | COTTON | 1 | 2 | 3 | | | |
| 35* | SURGICAL GAUZE | 1 | 2 | 3 | | | |
| 36* | POVIDON IODINE SOLUTION | 1 | 2 | 3 | | | |
| 37* | SURGICAL GLOVES (SIZE 6.5) | 1 | 2 | 3 | | | |
| 38* | SURGICAL GLOVES (SIZE 7) | 1 | 2 | 3 | | | |
| 39* | DISPOSABLE STERILE SYRINGE (5 ML) | 1 | 2 | 3 | | | |
| 40* | DISPOSABLE STERILE SYRINGE (10 ML) | 1 | 2 | 3 | | | |
| 41* | URISTICK GP (FOR GLUCOSE/ALBUMIN TEST) | 1 | 2 | 3 | | | |
| 42* | ELASTOMETRIC DRESSING MATTRESS | 1 | 2 | 3 | | | |

| 13530 | | (a) AV | /AILABLE? | | (b) | FUNCTI | ONING |
|-------|---------------------------|----------|-----------------------|------------------|-----|--------|---------------|
| 13530 | | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| | | | | | ļ | | |
| 43* | DISPOSABLE STERILE LANCET | 1 | 2 | 3 | | | |
| | SILK THREAD | 1 | 2 | 3 | | | |
| 44* | | | | | | | |

| 12520 | Do you have any of the following items available at | (A) A | VAILABLE? | | (B) | FUNC | TIONING |
|-----------|--|--|---|---|----------------------------|----------------------------|------------------------------|
| 13330 | this service site today? If available I will like to see them. For some of the items I will like to know if they are functioning. | OBSERVED | REPORTED, | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 1* | NON-SURGICAL VASECTOMY KIT | 1 | 2 | 3 | | | |
| 2* | RING FORCEPS*** | 1 - ▶b | 2 > b | 3 03 ◀ | 1 | 2 | 8 |
| 3* | VAS DISECTING FORCEPS*** | 1- > b | 2 - ▶ b | 3 04 4 | 1 | 2 | 8 |
| 4* | SMALL SURGICAL SCISSORS*** | 1 _ ▶b | 2 _ b | 3 05 ◀ | 1 | 2 | 8 |
| 5* | CONDOM | 1 | 2 | 3 | | | |
| 1353E | Please tell me if any of the following medicines are | (A) OBSERV | ED AVAILABLE | (| B) NOT OBSE | RVED | |
| | available at this services site today. I would like to see them. CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILAB TODAY/DK | LE | NO, OR NEVER AVAILABLE |
| 01* | INJECTION XYLOCAINE 1% | 1 | 2 | 3 | 4 | | 5 |
| 02* | ANTIBIOTIC (CIPROFLOXACIN OR AZITHROMYCIN) | 1 | 2 | 3 | 4 | | 5 |
| 03* | PARACETAMOL TABLETS | 1 | 2 | 3 | 4 | | 5 |
| 04* | VITAMIN B-COMPLEX TABLETS | 1 | 2 | 3 | 4 | | 5 |
| 1353F | CHECK Q1302.13 TUBAL LIGATION (TUBECTOMY) PROVIDED IN FACILITY TUBAL LIGATION (TUBECTOMY) NOT PROVIDED IN FACILITY | | | | | | |
| PR FA(| ASK TO BE SHOWN THE LOCATION IN THE FAC OVIDED.FIND THE PERSON MOST KNOWLEDGE CILITY. INTRODUCE YOURSELF, EXPLAIN THE PU | ILITY WHERE ABLE ABOU RPOSE OF T | E TUBECTOMY T TUBECTOMY HE SURVEY AN | / TUBAL LIGA / / TUBAL LIGA ID ASK THE FO | TION SER | VICES / VICES i QUES | ARE IN THE STIONS. |
| 1353G | I would like to know if the following items are like to know if they are functioning. If any of where in the facility they are kept. | available in the items ar | this service ar e maintained | ea today. For s elsewhere in t | some of th the facility | ie item y, plea | ns, I would se tell me |
| | ASSESS THE ROOM OR AREA FOR THEITEMS L BELOW. FOR ITEMS THAT YOU DO NOT SEE YOUR RESPONDENT TO SHOW THEM TO YOU | ISTED ROO , ASK ROO . ASS | M / AREA ASSESSE DM / AREA NOT F SESSED | D IN Q1353C PREVIOUSLY | | 11 31 | _ 1353I |
| | IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECOR IF NOT PREVIOUSLY ASSESSED, CIRCLE "31" AND AS THIS SERVICE SITE. | DED. SESS | | | | | |
| 1353H | | (a) A | VAILABLE? | | (b) | FUNC | TIONING |
| | | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01* | OPERATING THEATRE TABLE | 1- > b | 2 - ▶ b | 3 02 | 1 | 2 | 8 |
| 02* | OPERATING THEATRE LIGHT | 1 _ ▶b | 2 > b | 3 — | 1 | 2 | 8 |
| | | | | | | | |

| | | | | 03 | | | |
|-------|---|------------------|-----------------------|------------------|-----|---------|---------------|
| 03* | | 1_ > h | 2 b h | 3 | 1 | 2 | 8 |
| 05 | | 1-0 | 2 U | 04 | 1 | 2 | 0 |
| 04* | STERILIZER DRUM | 1 b | 2 _ ▶ b | 3 05 4 | 1 | 2 | 8 |
| 05* | BIG CURVE SCISSORS FOR CUTTING GAUZE OR BANDAGE | 1 b | 2 b | 3 06 4 | 1 | 2 | 8 |
| 06* | SCISSORS FOR CUTTING CLOTH | 1 - ▶b | 2 b | 3 07 ◀ | 1 | 2 | 8 |
| 07* | DIGITAL BP APPARATUS | 1- - b | 2 _ ▶ b | 3 09 ↓ | 1 | 2 | 8 |
| 09* | MANUAL BP APPARATUS | 1- b | 2 → b | 3 10 ◀ | 1 | 2 | 8 |
| 10* | STETHOSCOPE | 1 b | 2 b | 3 11 4 | 1 | 2 | 8 |
| 11* | WEIGHING SCALE (ADULT) | 1 - ▶b | 2 - ▶ b | 3 12 | 1 | 2 | 8 |
| 12* | GALLIPOT OR CUP FOR ANTISEPTIC SOLUTION | 1 | 2 | 3 | | | |
| 13* | KIDNEY TRAY | 1 | 2 | 3 | | | |
| 15* | LIFTER AND LIFTER JAR | 1 | 2 | 3 | | | |
| 16* | DRESSING JAR WITH LID/COVER | 1 | 2 | 3 | | | |
| 18* | TALQUEST BOOK | 1 | 2 | 3 | | | |
| 19* | IRON COT | 1 | 2 | 3 | | | |
| 20* | THERMOMETER | 1 - ▶b | 2 _▶ b | 3 21 | 1 | 2 | 8 |
| 21* | GOWN FOR CLIENTS | 1 | 2 | 3 | | | |
| 22* | GOWN FOR PROVIDER | 1 | 2 | 3 | | | |
| 23* | TROLLEY SHEET | 1 | 2 | 3 | | | |
| 24* | DRAW SHEET | 1 | 2 | 3 | | | |
| 25* | САР | 1 | 2 | 3 | | | |
| 26* | MASK | 1 | 2 | 3 | | | |
| 34* | COTTON | 1 | 2 | 3 | | | |
| 35* | SURGICAL GAUZE | 1 | 2 | 3 | | | |
| 36* | POVIDON IODINE SOLUTION | 1 | 2 | 3 | | | |
| 37* | SURGICAL GLOVES (SIZE 6.5) | 1 | 2 | 3 | | | |
| 38* | SURGICAL GLOVES (SIZE 7) | 1 | 2 | 3 | | | |
| 39* | DISPOSABLE STERILE SYRINGE (5 ML) | 1 | 2 | 3 | | | |
| 40* | DISPOSABLE STERILE SYRINGE (10 ML) | 1 | 2 | 3 | | | |
| 41* | URISTICK GP (FOR GLUCOSE/ALBUMIN TEST) | 1 | 2 | 3 | | | |
| 42* | ELASTOMETRIC DRESSING | 1 | 2 | 3 | | | |
| 43* | DISPOSABLE STERILE LANCET | 1 | 2 | 3 | | | |
| 44* | SILK THREAD | 1 | 2 | 3 | | | |
| 12521 | Do you have any of the following items | (A) AV | AILABLE? | | (B) | FUNCTIO | ONING |
| 13231 | available at this service site today? If available I will like to see them. For some of the items I will like to know if they are functioning. | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |

| 01* | ТИВЕСТОМҮ КІТ | 1 b | 2 b | 3 02 | 1 2 | 8 | | |
|-------|--|-----------------------|-------------------------|-----------------------------------|------------------------------|------------------------------|--|--|
| 02* | BP HANDLE*** | 1- - b | 2 b | 3 03 4 | 1 2 | 8 | | |
| 03* | NEEDLE HOLDER*** | 1- - b | 2 b | 3 04 ◀ | 1 2 | 8 | | |
| 04* | BABCOCK TISSUE FORCEP*** | 1- - b | 2_ b | 3 05 4 | 1 2 | 8 | | |
| 05* | LONG STRAIGHT ARTERY FORCEP (MEDIUM)*** | 1 b | 2_ b | 3 06 | 1 2 | 8 | | |
| 06* | CURVED MOSQUITO ARTERY FORCEP*** | 1 b | 2_ b | 3 07 | 1 2 | 8 | | |
| 07* | ALICE TISSUE FORCEP*** | 1- > b | 2 b | 3 08 4 | 1 2 | 8 | | |
| 08* | PLAIN DISSECTING FORCEP*** | 1 b | 2 _ ▶ b | 3 09 4 | 1 2 | 8 | | |
| 09* | TOOTH DISSECTING FORCEP*** | 1 - ▶b | 2 - ▶ b | 3 10 | 1 2 | 8 | | |
| 10* | MAYO SCISSORS*** | 1 - ▶b | 2 → b | 3 11 ↓ | 1 2 | 8 | | |
| 11* | SPONGE HOLDING FORCEP*** | 1 b | 2 b | 3 12 | 1 2 | 8 | | |
| 18‡ | VICRYL | 1 | 2 | 3 | | | | |
| 1353J | Please tell me if any of the following | (A) OBSERVE | D AVAILABLE | (В |) NOT OBSERVE |) | | |
| | medicines are available at this services site today. I would like to see them. | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NO, OR NEVER AVAILABLE | | |
| 01* | INJECTION ATROPINE SULPHATE (0.6MG/ML) | 1 | 2 | 3 | 4 | 5 | | |
| 02* | INJECTION PROMETHAZINE (12.5MG/ML) | 1 | 2 | 3 | 4 | 5 | | |
| 03* | INJECTION PETHIDINE (25MG/ML) | 1 | 2 | 3 | 4 | 5 | | |
| 04* | INJECTION PENTAZOSIN (30MG/ML) | 1 | 2 | 3 | 4 | 5 | | |
| 05* | INJECTION LIGNOCAINE (1%) | 1 | 2 | 3 | 4 | 5 | | |
| 06* | DIAZEPAM TABLETS (5MG) | 1 | 2 | 3 | 4 | 5 | | |
| 07* | ANTIBIOTIC (CIPROFLOXACIN OR AZITHROMYCIN) | 1 | 2 | 3 | 4 | 5 | | |
| 08* | PARACETAMOL TABLETS | 1 | 2 | 3 | 4 | 5 | | |
| 09* | IRON + FOLIC ACID TABLETS | 1 | 2 | 3 | 4 | 5 | | |
| THAN | THANK YOUR RESPONDENT AND MOVE TO YOUR NEXT DATA COLLECTION POINT IF DIFFERENT FROM CURRENT LOCATION. | | | | | | | |

SECTION 14: ANTENATAL CARE

| 1400 | 0 CHECK Q102.05 | | | | | | |
|------------------|--|---|--------------------------------|-------------------------------------|--------------------------------|----------------------------------|---------------------------------------|
| | ANC SERVICES AVAILABLE | | ANC SEF | | AILABLE | | |
| | | NEXT SECTION OR SERVICE SITE | | | | | |
| ASK T KNOWLE | O BE SHOWN THE LOCATION IN THE FACILITY WHERE DGEABLE ABOUT ANTENATAL CARE SERVICES IN THE AND ASK THE | ANTENAT FACILITY. FOLLOWII | AL CARE SERVINTRODUCE Y | /ICES ARE PRC OURSELF, EXF S. | VIDED. FIND 1 PLAIN THE PUR | THE PERSON POSE OF TH | N MOST HE SURVEY |
| 1401 | How many days in a month are antenata | al care | | | | | |
| | offered at this facility? # OF DAYS PER MONTH | | | | | | |
| | USE A 4-WEEK MONTH TO CALCULATE # OF DAYS | | | | | | |
| 1402 | Do this facility provide any of the following so women as part of routine ANC? | ervices to | o pregnant | YES | NO | | |
| 01 | IRON SUPPLEMENTATION | | | 1 | 2 | | |
| 02 | FOLIC ACID SUPPLEMENTATION | | | 1 | 2 | | |
| 02a* | COMBINED IRON AND FOLIC ACID SUPPLEMENTATIO | ON | | 1 | 2 | | |
| 05* | PROVIDE MISOPROSTOL TABLET/CAPSULE FOR HOM | IE-BASED | DELIVERIES | 1 | 2 | | |
| 06 [‡] | ANC Counselling | | | 1 | 2 | | |
| 06a [‡] | Post-partum Family Planning (PPFP) counselling | | | 1 | 2 | | |
| 07 [‡] | BP measurement | 1 2 | | | | | |
| 08 [‡] | Antenatal Corticosteroid administration | 1 2 | | | | | |
| 09 [‡] | Weight monitoring | 1 2 | | | | | |
| 10 [‡] | Height monitoring | 1 2 | | | | | |
| 1406 | Do ANC providers in this facility provide any of the following tests from this site to pregnant women as | (A) C AV | DBSERVED AILABLE | (B) NOT OBSERVED | | | |
| | IF YES, ASK TO SEE THE TEST KIT OR EQUIPMENT. IF TEST NOT DONE IN ANC, PROBE TO DETERMINE. IF THE TEST IS DONE ELSEWHERE IN THE FACILITY | AT LEAST ONE VALID | AVAILABL E NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NO, OR NEVER AVAILABL E | AVAILABLE ELSEWHERE IN FACILITY |
| | CHECK TO SEE IF AT LEAST ONE TEST KIT OF EACH TEST IS VALID/UNEXPIRED | | | | | | |
| 02 | | 1 | 2 | 3 | 4 | 5 | 6 |
| 03 | | 1 | 2 | 3 | 4 | 5 | 6 |
| 04 | | 1 | 2 | 3 | 4 | 5 | 6 |
| 05 | VDRL test for syphilis | 1 | 2 | 3 | 4 | 5 | 6 |
| 1408 | Do ANC providers in this facility routinely diagnose and treat RTIs / STIs, or are RTI/ STI clients referred to another provider or location for diagnosis and treatment? | ROUTINELY DIAGNOSE AND TREAT RTIS/STIS DIAGNOSE BUT REFER ELSEWHERE FOR TREATMENT REFER ELSEWHERE IN FACILITY FOR DIAGNOSIS & TREATMENT | | | | | |
| 1409 | Do you have the ANC guideline/protocol/ manual available in this service area today? | YES NO | | | 1 | ▶ 1411 | |
| 1410 | May I see the ANC guideline/protocol/ manual? ACCEPTABLE IF PART OF OTHER GUIDELINES | OBSERV REPOR | /ED TED NOT SEE | EN | <u>1</u> 2 | ▶ 1415 | |

| 1411 | Do you have any other ANC guideline/ protocol/manual available in this service area today? | YES NO | | 1 2— | ▶ 1415 |
|------|---|--------------------------|----------------------------------|------------------------------|---------------|
| 1412 | May I see the other guideline/protocol/ manual? | OBSERVED | SEEN | 1 2 | |
| 1415 | Do you have visual aids for client education on subjects related to pregnancy or antenatal care available in this service area today? | YES NO | | 1 2— | ▶1417 |
| 1416 | May I see the visual aids for client education? | OBSERVED REPORTED NOT | SEEN | 1 2 | |
| 1417 | Are individual client cards or records (ANC service register) for ANC clients maintained at this service site? | YES NO | | 1 2 <u>—</u> | ▶1419 |
| 1418 | May I see a blank copy of the client records (ANC service register) or cards? | OBSERVED | SEEN | 1 2 | |
| 1419 | Does this facility have a system whereby observation or parameters for ANC clients are routinely carried out before the consultation? | YES NO | ▶ 1421 | | |
| | IF YES, ASK TO SEE THE PLACE WHERE THESE ACTIVITIES TAKE PLACE. | | | | |
| 1420 | OBSERVE IF THE BELOW ACTIVITIES ARE BEING DONE ROUTINELY. IF YOU DO NOT SEE AN ACTIVITY, ASK: IS [ACTIVITY YOU DO NOT SEE] routinely done for all antenatal care clients? | ACTIVITY OBSERVED | ACTIVITY REPORTED NOT SEEN | ACTIVI TY NOT DON E | DON'T KNOW |
| 01 | Weighing of clients | 1 | 2 | 3 | 8 |
| 02 | Taking blood pressure | 1 | 2 | 3 | 8 |
| 03 | Conducting group/individual health education sessions | 1 | 2 | 3 | 8 |
| 04 | Urine test for protein | 1 | 2 | 3 | 8 |
| 05 | Blood test for anemia (hemoglobin) | 1 | 2 | 3 | 8 |
| 08 | Measuring client's height | 1 | 2 | 3 | 8 |
| 09* | Ultrasonography | 1 | 2 | 3 | 8 |
| 10* | Urine test for glucose | 1 | 2 | 3 | 8 |
| 11* | Blood test for glucose | 1 | 2 | 3 | 8 |
| 12* | HBsAg | 1 | 2 | 3 | 8 |

EQUIPMENT AND SUPPLIES FOR ROUTINE ANC

| 1/121 | I would like to know if the following items | (a) A\ | (a) AVAILABLE | | | FUNCTIO | ONING |
|-------|---|---------------|-----------------------|------------------|-----|---------|---------------|
| 1421 | are available in this service area and are functioning. | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01 | DIGITAL BP APPARATUS | 1 - ▶b | 2 _ ▶ b | 3 02 | 1 | 2 | 8 |
| 02 | MANUAL BP APPARATUS | 1 - ▶b | 2 _ ▶ b | 3 03 4 | 1 | 2 | 8 |
| 03 | STETHOSCOPE | 1 - ▶b | 2 _ ▶ b | 3 04 | 1 | 2 | 8 |
| 04 | EXAMINATION LIGHT (FLASHLIGHT OK) | 1 - ▶b | 2 → b | 3 05 🗲 | 1 | 2 | 8 |
| 05 | FETAL STETHOSCOPE/PINNARD/ FETAL DOPPLER | 1 _ ▶b | 2 b | 3 06 ◀ | 1 | 2 | 8 |

| 06 | ADULT WEIGHING SCALE | 1 _ ▶b | 2 b | 3 07 4 | 1 | 2 | 8 |
|-----------------|--------------------------------|---------------|-----------------|------------------|---|---|---|
| 07 | EXAMINATION BED OR COUCH | 1 | 2 | 3 | | | |
| 08 | TAPE MEASURE FOR FUNDAL HEIGHT | 1 | 2 | 3 | | | |
| 09 [‡] | Thermometer | 1 b | 2 > b | 3 10 4 | 1 | 2 | 8 |
| 10 [‡] | Glucometer with strip | 1 - ▶b | 2 — b | 3 1422◀ | 1 | 2 | 8 |

| 1422 | 22 Please tell me if any of the following | | (A) OBSERVED AVAILABLE (B) | | |) NOT OBSERVED | |
|-----------------|---|-----------------------|----------------------------|-----------------------------------|------------------------------|------------------------------|--|
| | medicines are available at this services site today. I would like to see them. CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE NOT SEEN | NOT AVAILABLE TODAY/DK | NO, OR NEVER AVAILABLE | |
| 01 | IRON TABLETS (INDIVIDUAL TABLETS) | 1 | 2 | 3 | 4 | 5 | |
| 02 | FOLIC ACID TABLETS (INDIVIDUAL TABLETS) | 1 | 2 | 3 | 4 | 5 | |
| 03 | COMBINED IRON AND FOLIC ACID TABLETS | 1 | 2 | 3 | 4 | 5 | |
| 07 | MISOPROSTOL TABLET | 1 | 2 | 3 | 4 | 5 | |
| 08 [‡] | CORTICOSTEROIDS FOR PRE-TERM LABOR | 1 | 2 | 3 | 4 | 5 | |

| 1022 | Now I would like to ask you specifically about vaccination services for mothers. For each of the following | | | | | | |
|-------|--|-----------|--------------|---------------|------------------|--------------------|--|
| 1022 | services, please tell me whether the service is offered by your facility, and if so, how many days per month | | | | | | |
| | the service is provided at the facility, if any. | | | | | | |
| | MOTHER VACCINATION SERVICE | | | (a) | | | |
| | | # OF DAY | S PER MONTH | VACCINATION S | ERVICE IS PRO | VIDE TO THE | |
| | TT vessing | | MOT | | ACILITY | | |
| 01 | 11 Vaccine | # OF DAY | s | | | | |
| | | 00=NO SE | RVICE | | | | |
| 1023* | Do you have the <i>national guidelines</i> for mother | YES | | | 1 | | |
| 1025 | vaccinations available in this service area | NO | | | 2 — | → 1025 | |
| | today? | | | | | | |
| 1024* | May I see the guidelines? | OBSERVE | D | | | → 1027 | |
| 1024 | | REPORTE | D, NOT SEEN. | | 2 | | |
| 1025* | Do you have any other guidelines for mother | YES | | | 1 | | |
| 1025 | vaccinations available in this service area | NO | | | 2 — | → 1027 | |
| | today? | | | | | | |
| 1026* | May I see the other guidelines? | OBSERVE | D | | 1 | | |
| 1020 | | REPORTE | D, NOT SEEN. | | 2 | | |
| 1027* | Please tell me if each of the following vaccines | (A) OI | BSERVED | (B) I | NOT OBSER | VED | |
| 1027 | is available in the facility today. If available, I | AVA | ILABLE | | | | |
| | would like to see it. | ATLEAST | AVAILABLE | REPORTED | ΝΟΤ Δναμαρί ε | NEVER AVAILABLE | |
| | IF AVAILABLE, CHECK TO SEE IF AT LEAST ONE IS | ONE VALID | NONE VALID | NOT SEEN | TODAY/DK | | |
| | VALID (NOT EXPIRED, VACCINE VIAL MONITOR | | | | | | |
| | UNCHANGED (NOT FROZEN)) | | | | | | |
| 01 | TT vaccine | 1 | 2 | 3 | 4 | 5 | |
| | | 1 | | | | | |

| 1450 | ASSESS THE ROOM OR AREA FOR THE ITEMS LISTED BELOW. FOR ITEMS THAT YOU DO NOT SEE, ASK YOUR RESPONDENT TO SHOW THEM TO YOU. IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECORDED | GENERAL INFORM CHILD VACCINATI CHILD CURATIVE FAMILY PLANNIN DELIVERY [Q1651 TUBERCULOSIS [C NCD [Q2351] NOT PREVIOUSLY | MATION [Q710] ION [Q1051] CARE [Q1251] NG [Q1351] 21951] Y SEEN | | 2 3 4 7 9 2 4 7 9 2 4 7 7 9 2 4 7 7 9 2 4 7 7 9 2 4 7 7 9 2 4 7 7 8 8 8 7 8 8 8 8 7 8 9 7 8 8 8 9 7 8 9 7 8 9 8 9 |
|------|--|--|--|----------------------|---|
| 1451 | STANDARD PRECAUTIONS AND CONDITIONS EXAMINATION | FOR CLIENT | OBSERVED | REPORTED NOT SEEN | NO/ NOT AVAILABLE |
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POL | JR PITCHER) | 1 | 2 | 3 |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOA | P) | 1 | 2 | 3 |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLAS | 1 06 ◀ | 2 | 3 | |
| 05 | OTHER WASTE RECEPTACLE | 1 | 2 | 3 | |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITA | NE, ALCOHOL] | 1 | 2 | 3 |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH AUTO-DISABLE SYRINGES WITH NEEDLES | H NEEDLES OR | 1 | 2 | 3 |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 |
| 11 | GOWNS | | 1 | 2 | 3 |
| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | 1 | 2 | 3 | |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 |
| 1452 | DESCRIBE THE SETTING OF THE ANC SERVICE ROOM OR AREA. PRIVATE ROOM W PRIVACY VISUAL PRIVACY NO PRIVACY | | I WITH AUDITORY Y ONLY | | L 2 3 4 |

POSTNATAL CARE SERVICES

| 1453 | CHECK Q102.20 PNC SERVICE PROVIDED | PNC SERVICE NOT PROVIDED |
|--------------------|--|--------------------------|
| 1454 | Is there a dedicated room/area for postnatal care (PNC) examination? | YES1 NO2 →1455a |
| 1455 | DESCRIBE THE SETTING OF THE PNC SERVICE ROOM OR AREA. | PRIVATE ROOM |
| 1455a [‡] | Is postnatal care (PNC) provided to mother | YES1 NO2 |

| 1455b [‡] | Is postnatal care (PNC) provided to newborn | YES1 NO2 | |
|--------------------|---|---------------------------------|--------|
| 1456* | How many days each week are PNC service available at this facility? | DAYS PER WEEK | |
| 1457* | Do you have the PNC guideline/protocol/manual available in this service area today? | YES1 NO2 | ▶1459 |
| 1458* | May I see the PNC guideline/protocol/manual? ACCEPTABLE IF PART OF OTHER GUIDELINES | OBSERVED1— REPORTED NOT SEEN | 1461 |
| 1459* | Do you have <i>any other PNC guideline/protocol/</i> <i>manual</i> available in this service area today? | YES1 NO | ▶1461 |
| 1460 | May I see the other guideline/protocol/manual? | OBSERVED1 REPORTED NOT SEEN2 | |
| 1461 | Do you have the postpartum family planning guideline/protocol/manual available in this service area today? | YES1 NO2— | ▶ 1463 |
| 1462 | May I see the postpartum family planning guideline/protocol/manual? | OBSERVED1 REPORTED NOT SEEN2 | |
| 1463 [‡] | Are individual client cards or records (PNC service register) for PNC clients maintained at this service site? | YES1 NO2 | ▶ 1600 |
| 1464 [‡] | May I see a blank copy of the client records (PNC service register) or cards? | OBSERVED1 REPORTED NOT SEEN2 | |

SECTION 16: DELIVERY AND NEWBORN CARE

| 1600 | CHECK Q102.07 | |
|------|--|---|
| | NORMAL DELIVERY AVAILABLE | NORMAL DELIVERY NOT AVAILABLE |
| | | NEXT SECTION OR SERVICE SITE |
| | ASK TO BE SHOWN THE LOCATION IN THE FACILITY WHI FIND THE PERSON MOST KNOWLEDGEABLE AB(INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF TH | ERE NORMAL DELIVERY SERVICES ARE PROVIDED. OUT DELIVERY SERVICES IN THE FACILITY. IE SURVEY AND ASK THE FOLLOWIQUESTIONS. |
| 1601 | Is a person skilled in conducting deliveries present at the facility today or on call at all times (24 hours a day), including weekends, to provide care? Specifically, I am referring to medical specialists, medical officers, nurses, and family welfare visitors and midwives | YES |
| 1602 | Is there a duty schedule or call list for 24-hr staff assignment? | YES1 NO2 1604 |
| 1603 | May I see the duty schedule or call list for 24-HR staff assignment? | OBSERVED1 REPORTED NOT SEEN2 |

EmONC SIGNAL FUNCTION

| 1604 | Please tell me if any of the following | (a) EVER PR | ROVIDED IN FACIL | N FACILITY (b) PROVIDED IN PAST 3 N | | | ONTHS |
|------|--|-------------|------------------|-------------------------------------|-----|---------|---------------------|
| | interventions have ever been carried out by providers as part of their work in this facility, and if so, whether the intervention has been carried out at least once during the past 3 months. | YES | NO | DON'T KNOW | YES | NO | DON'T KNOW |
| 01 | PARENTERAL ADMINISTRATION OF ANTIBIOTICS (IV OR IM) | 1 🛶 | 2 02◀ | 8 02 4 | 1 | 2 | 8 |
| 02 | PARENTERAL ADMINISTRATION OF OXYTOCIC (IV OR IM) | 1 🛶 | 2 03◀ | 8 03◀ | 1 | 2 | 8 |
| 03 | PARENTERAL ADMINISTRATION OF ANTICONVULSANT FOR HYPERTENSIVE DISORDERS OF PREGNANCY /PREVENTION OF ECLAMPSIA (IV OR IM) | 1 🛶 | 2 04 ◀ | 8 04◀ | 1 | 2 | 8 |
| 04 | ASSISTED VAGINAL DELIVERY | 1 🛶 | 2 05 ◀ | 8 05 ↓ | 1 | 2 | 8 |
| 05 | MANUAL REMOVAL OF PLACENTA | 1 🛶 | 2 06◀ | 8 06◀ | 1 | 2 | 8 |
| 06 | REMOVAL OF RETAINED PRODUCTS OF CONCEPTAION | 1 - | 2 07◀ | 8 07 ↓ | 1 | 2 | 8 |
| 07 | NEONATAL RESUSCITATION | 1 — | 2 08◀ | 8 08 ~ | 1 | 2 | 8 |
| 08 | CORTICOSTEROIDS FOR PRE-TERM LABOR NOTE: THIS IS NOT A SIGNAL FUNCTION | 1 🛶 | 2 1605◀ | 8 1605 ◀ | 1 | 2 | 8 |
| 1605 | Do you have the national guideline/protocol/manual for BEmONC a in this service site? | vailable | P YES | | | 1 2— | 1607 |
| 1606 | May I see the guideline/protocol/manual BEmONC? | for | OBSERVED | NOT SEEN | | 1 2 | |
| 1607 | Do you have the national guideline/protocol/manual for CEmONC? ACCEPTABLE IF PART OF ANOTHER GUIDELINE | | YES NO | | | 1 2 | →1608a [‡] |

| 1608 | May I see the national guideline/protocol/manual for CEmONC? | | OBSERVED1 REPORTED NOT SEEN2 | | | | |
|--------------------|---|---------------------------|---------------------------------|-----------------------------------|-------------------------|---------|----------------------|
| 1608a [‡] | Please tell me if you have any of the following materials. IF YES, ASK TO SEE | (A) AV OBSERVED | | /AILABLE REPORTED, NOT SEEN | NOT AVAILABLE | - | |
| 01 | EmONC Register | 1 | | 2 | 3 02 | | |
| 02 | Do you have delivery Register? | 1 | | 2 | 3 1608b [‡] | | |
| 1608b [‡] | Do you maintain documentation of the patier who are referred outside the facility? | nt | YES . NO | | | 1 2— | ► 1608e [‡] |
| 1608c [‡] | Do you write diagnosis in the documentation the patient who are referred outside the facil | of ity? | YES1 NO2 | | | 1 2 | |
| 1608d [‡] | May I see the documentation of the patient w are referred outside the facility? | of the patient who ty? | | ERVED ORTED NOT S | EEN | 1 2 | |

NEWBORN SIGNAL FUNCTION

| 1608e [‡] | Please tell me if any of the following | (a) EVER PROVIDED IN FACILITY (b) | | (b) PROVIDED IN PAST 3 MONTHS | | | |
|--------------------|--|-----------------------------------|------------------|-------------------------------|-----|----|---------------|
| | interventions have ever been carried out by providers as part of their work in this facility, and if so, whether the intervention has been carried out at least once during the past 3 months. | YES | NO | DON'T KNOW | YES | NO | DON'T KNOW |
| 01 | IRON AND FOLIC ACID SUPPLIMENTATION IN PREGNANT WOMEN | 1 b | 2 02 ◀ | 8 02 4 | 1 | 2 | 8 |
| 02 | HAND WASH EACH TIME BEFORE TOUCHING THE BABY | 1 b | 2 03 ◀ | 8 03◀ | 1 | 2 | 8 |
| 03 | DRYING OF THE BABY IMMEDITELY AFTER BIRTH | 1 b | 2 04 | 8 04◀ | 1 | 2 | 8 |
| 04 | DELAYED UMBILICAL CORD CLAMPING APPROXIMATELY 1-3 MINUTES AFTER BIRTH | 1 b | 2 05 4 | 8 05◀ | 1 | 2 | 8 |
| 05 | CLEAN CORD CUTTING STERILE BLADE/SCISSOR | 1 b | 2 06 | 8 06 4 | 1 | 2 | 8 |
| 06 | SINGLE APPLICATON OF CHX 7.1% TO UMBILICAL CORD FOLLOWED BY DRY CORD CARE | 1 — b | 2 07 ◀ | 8 07◀ | 1 | 2 | 8 |
| 07 | PERFORM SKIN TO SKIN CARE IMMEDIATELY AFTER BIRTH FOR HOURS | 1 b | 2 08 ◀ | 8 08 4 | 1 | 2 | 8 |
| 08 | EARLY INITIATION OF BREAST FEEDING WITHIN ONE HOUR OF BIRTH | 1 b | 2 09 4 | 8 09◀ | 1 | 2 | 8 |
| 09 | IMMUNIZE PREGNANT AGAINST TETANUS TOXOID TO PREVENT TETANUS FOR BOTH MOTHER AND NEWBORN | 1 b | 2 10 ◀ | 8 10 4 | 1 | 2 | 8 |
| 10 | ADMINISTER ORAL ANTIBIOTIC FOR PREVENTION OF INFECTION IN NEWBORN | 1 b | 2 11 ◀ | 8 | 1 | 2 | 8 |
| 11 | ADMINISTER INTRAMASCULAR ANTIBIOTICS FOR PREVENTION OF INFECTION IN NEWBORN | 1 b | 2 12 | 8 | 1 | 2 | 8 |
| 12 | ADMINISTER INTRAVENOUS ANTIBIOTICS FOR PREVENTION OF INFECTION IN NEWBORN | 1 b | 2 13 | 8 13 | 1 | 2 | 8 |
| 13 | SUCTION OF AIRWAY OF NEWBORN IF SECRETION OR MECONIUM PRESENT | 1 — b | 2 | 8 | 1 | 2 | 8 |

| | | | 14 | 14 | | | |
|--------------------|--|---------------------|------------------------------|------------------|--------|----------|-------------------------|
| 14 | ADMINISTER ANTIBIOTICS FOR PRE-MATURE RUPTURE OF MEMBRANE (PROM) | 1 b | 2 15 | 8 15 | 1 | 2 | 8 |
| 15 | OXYGEN THERAPY FOR MANAGEMENT OF NEWBORN INFECTION AND RESPIRATORY PROBLEMS | 1 — b | 2 16 | 8 16◀ | 1 | 2 | 8 |
| 16 | KMC FOR PRETERM OR LBW NEWBORN | 1 — b | 2 17 | 8 17 | 1 | 2 | 8 |
| 17 | PHOTOTHERAPY FOR TERM AND PRETERM WITH HYPERBILIRUBINAEMIA | 1 b | 2 18 | 8 18 4 | 1 | 2 | 8 |
| 18 | INCUBATOR SUPPORT FOR SICK TERM BABY, PRETERM BABY OR IF THE BABY RECEIVING KMC BECOMES SICK | 1 b | 2 — 1608f‡◀ | 8 | 1 | 2 | 8 |
| 1608f [‡] | Do you have the national guideline for new sepsis management? | wborn | YES NO | | | 1 2 | ► 1608h [‡] |
| 1608g [‡] | May I see the guideline for newborn sepsi management? | S | OBSERVED REPORTED N | OT SEEN | | 1 | |
| 1608h [‡] | Do you have the national guideline on app 7.1% chlorhexidine? | olication of | YES NO | | | 1 2— | ► 1608j [‡] |
| 1608i [‡] | May I see the national guideline on applic 7.1% chlorhexidine? | ation of | OBSERVED REPORTED N | OT SEEN | | 1 2 | |
| 1608j [‡] | Do you have the national guideline on Anter corticosteroid? | natal | YES NO | | | 1 2— | ► 1609 |
| 1608k [‡] | May I see the national guideline on Antena corticosteroid? | atal | OBSERVED REPORTED N | OT SEEN | | 1 2 | |
| 1609 | Do you have guideline/protocol/manual o management of pre-term labor? | n | YES NO | | | 1 2 | ► 1611 |
| 1610 | May I see the guideline/protocol/manual management of pre-term labor? | on | OBSERVED REPORTED N | OT SEEN | | 1 2 | |
| 1611 | Does this facility practice Kangaroo Mothe low-birth-weight babies? | er Care for | YES NO | | | 1 2 — | → 1613 |
| 1612 | Is there a separate room or space for Kang mother Care or is it integrated into the ma postnatal ward? | garoo ain | YES, SEPARAT YES, INTEGRA | E ROOM TED | | 1 2 | |
| 1613 | Do providers of delivery services in this fa partograph to monitor labor and delivery | cility use | YES NO USE OF PA | ARTOGRAPH | | 1 2— | ► 1615 |
| 1614 | Are partographs used routinely (for all cas selectively (only for some cases) to monito and delivery in this facility? | ses) or or labor | ROUTINELY SELECTIVELY. | | | 1 2 | |
| 1615 | How many dedicated maternity beds are a in this facility? | available | # OF MATERN | IITY BEDs | | | |
| | | | DON'T KNOW | l | | | |
| 1616 | How many dedicated delivery beds are av this facility? | ailable in | # OF DEDICAT BEDs | ED DELIVERY | | | |
| | | | # OF ADJUSTA | ABLE DELIVERY | ' BEDs | | |
| | | | | | | | |
| | | | | 1 | | 998 | |

| 1617 | Does the facility conduct regular reviews of maternal or newborn deaths or "near-misses"? | YES1 NO, DOES NOT PARTICIPATE2 | ► 1622 |
|------|---|-----------------------------------|-----------|
| 1618 | Are reviews done for mothers only, newborns only, or for both mothers and newborns? | FOR MOTHERS ONLY | 1620 |

| 1619 | How often are reviews of <u>maternal deaths</u> or <u>"near</u> <u>misses"</u> carried out? | EVERY WEEKS | |
|------|--|---|-------|
| | | ONLY WHEN CASE OCCURS53 | |
| | | DON'T KNOW98 | |
| 1620 | CHECK Q1618 RESPONSE "3" CIRCLED | RESPONSE "3" NOT CIRCLED | →1622 |
| 1621 | How often are reviews of <u>newborn deaths</u> or <u>"near misses"</u> carried out? | EVERY WEEKS ONLY WHEN CASE OCCURS 53 ALWAYS WITH MATERNAL REVIEWS | |
| | | DON'T KNOW 98 | |

EQUIPMENT AND SUPPLIES FOR ROUTINE DELIVERIES AND NEWBORN CARE

| 1622 | I would like to know if the following items are | (A) AV | /AILABLE? | | (B) | FUNCTI | ONING |
|------|--|------------------|-----------------------|------------------|-----|--------|---------------|
| 1022 | available in this delivery area and are functioning. | OBSERVED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01 | INCUBATOR | 1 b | 2 b | 3 02 | 1 | 2 | 8 |
| 02 | OTHER EXTERNAL HEAT SOURCE/ LED Phototherapy machine | 1 _ ▶b | 2 b | 3 03 ◀ | 1 | 2 | 8 |
| 03 | EXAMINATION LIGHT (FLASHLIGHT OK) | 1 - ▶b | 2 b | 3 04 ◀ | 1 | 2 | 8 |
| 04 | SUCTION APPARATUS WITH CATHETER | 1 _ ▶b | 2_ b | 3 05 4 | 1 | 2 | 8 |
| 05 | SUCTION BULB OR PENGUIN SUCKER | 1 - ▶b | 2 b | 3 06 4 | 1 | 2 | 8 |
| 06 | MANUAL VACUUM EXTRACTOR (FOR VACUUM- ASSISTED DELIVERY) | 1- > b | 2 _ ▶ b | 3 07 ◀ | 1 | 2 | 8 |
| 07 | VACUUM ASPIRATION KIT OR D&C KIT | 1- - b | 2 _ ▶ b | 3 08 4 | 1 | 2 | 8 |
| 08 | NEWBORN BAG & MASK (AMBU BAG & MASK) | 1 b | 2_ b | 3 09 ◀ | 1 | 2 | 8 |
| 09 | THERMOMETER | 1 - ▶b | 2 b | 3 10 | 1 | 2 | 8 |
| 10 | THERMOMETER FOR LOW-BODY TEMPERATURE | 1 b | 2 b | 3 11 4 | 1 | 2 | 8 |
| 11 | INFANT WEIGHING SCALE | 1 - ▶b | 2 _ ▶ b | 3 12 | 1 | 2 | 8 |

| 12 | FETAL STETHOSCOPE/ FETAL DOPPLER | 1 _ ▶b | 2 b | 3 <u> </u> 13 |] 1 | 2 8 |
|---|--|------------------|------------------|--|---|--|
| 13 | DIGITAL BLOOD PRESSURE APPARATUS | 1 - ▶b | 2 -→ b | 3 <u> </u> 14 • |] 1 | 2 8 |
| 14 | MANUAL BLOOD PRESSURE MACHINE | 1 - ▶b | 2 -→ b | 3 — 15 🗲 |] 1 | 2 8 |
| 15 | STETHOSCOPE | 1 _ ▶b | 2 > b | 3 — 16 🗲 |] 1 | 2 8 |
| 16 [‡] | Drip stand | 1 b | 2 b | 3 <u> </u> 17 |] 1 | 2 8 |
| 17 [‡] | Suture, dressing set (forceps, scissors) | 1 _ ▶b | 2_ b | 3 <u> </u> 18 |] 1 | 2 8 |
| 18 [‡] | Pulse oximeter | 1 - ▶b | 2 - ▶ b | 3 <u> </u> 19 |] 1 | 2 8 |
| 20 [‡] | Oxygen cylinder with flowmeter for newborn | 1 | 2- > b | 3 <u></u> 21 - |] 1 | 2 8 |
| 21 [‡] | Nasal catheter | 1 | 2 b | 3 | | |
| 22 [‡] | Nasal Cannula | 1 - ▶b | 2_ b | 3 | | |
| 23 [‡] | Fluorescent tube | 1- > b | 2 → b | 3 <u> </u> 24 |] 1 | 2 8 |
| 24 [‡] | Nasogastric tube | 1 _ ▶b | 2 b | 3 — 1623 4 |] 1 | 2 8 |
| 4 6 9 9 | | | | | | |
| 1623 | Do you have any of the following items? If y them | es, I would | d like to see | OBSERVED | NOT SEEN | NOT AVAILABLE |
| 1623 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU | ves, I would | d like to see | OBSERVED 1 | NOT SEEN | NOT AVAILABLE 3 |
| 1623 1 2 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT | ves, I would | d like to see | OBSERVED 1 1 | REPORTED, NOT SEEN 2 2 | NOT AVAILABLE 3 3 |
| 1623 1 2 3 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD | ves, I would | d like to see | OBSERVED 1 1 1 | REPORTED, NOT SEEN 2 2 2 2 | NOT AVAILABLE 3 3 3 |
| 1623 1 2 3 4 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM | IPS | d like to see | OBSERVED 1 1 1 1 1 1 | 2 2 2 2 2 2 2 | NOT AVAILABLE 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS | IPS | d like to see | OBSERVED 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2 2 2 2 2 2 2 2 2 2 2 2 2 | NOT AVAILABLE 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD | IPS | d like to see | OBSERVED 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 7 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE | IPS | d like to see | OBSERVED 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 | REPORTED, NOT SEEN 2 | NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 7 8 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER | IPS | d like to see | OBSERVED 1 | REPORTED, NOT SEEN 2 | NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 7 8 9 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) | IPS | d like to see | OBSERVED | REPORTED, NOT SEEN 2 | NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 7 8 9 10 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) FORCEPS (MEDIUM) | IPS | d like to see | OBSERVED I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I | REPORTED, NOT SEEN 2 | NOT AVAILABLE 3 |
| 1623 1 2 3 4 5 6 7 8 9 10 11 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) FORCEPS (MEDIUM) SPONGE HOLDER | IPS | d like to see | OBSERVED 1 | REPORTED, NOT SEEN 2 <td>NOT AVAILABLE 3</td> | NOT AVAILABLE 3 |
| 1623 1 2 3 4 5 6 7 8 9 10 11 12 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) FORCEPS (MEDIUM) SPONGE HOLDER BLANK PARTOGRAPH | IPS | d like to see | OBSERVED I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I | REPORTED, NOT SEEN 2 <td>NOT AVAILABLE 3</td> | NOT AVAILABLE 3 |
| 1623 1 2 3 4 5 6 7 8 9 10 11 12 13 | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) FORCEPS (MEDIUM) SPONGE HOLDER BLANK PARTOGRAPH STAIRS (FOR CLIMBING ONTO DELIVERY BED) | IPS | d like to see | OBSERVED 1 | REPORTED, NOT SEEN 2 <td>NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 3</td> | NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 7 8 9 10 11 12 13 14 ‡ | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRU DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) FORCEPS (MEDIUM) SPONGE HOLDER BLANK PARTOGRAPH STAIRS (FOR CLIMBING ONTO DELIVERY BED) TWO OR MORE PIECES OF WARM CLOTH FOR DRY | ING | d like to see | OBSERVED 1 | REPORTED, NOT SEEN 2 <td>NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 </td> | NOT AVAILABLE 3 3 3 3 3 3 3 3 3 3 3 3 |
| 1623 1 2 3 4 5 6 7 8 9 10 11 12 13 14 ‡ | Do you have any of the following items? If y them DELIVERY BED COMPLETE WITH RODS AND STIRRL DELIVERY KIT CORD CLAMP/THREAD SPECULUM EPISIOTOMY SCISSORS SCISSORS OR BLADE TO CUT CORD SUTURE MATERIAL WITH NEEDLE NEEDLE HOLDER FORCEPS (LARGE) FORCEPS (MEDIUM) SPONGE HOLDER BLANK PARTOGRAPH STAIRS (FOR CLIMBING ONTO DELIVERY BED) TWO OR MORE PIECES OF WARM CLOTH FOR DRY STERILE CLOTH FOR WRAPPING | ING | d like to see | OBSERVED 1 | REPORTED, NOT SEEN 2 <td>NOT AVAILABLE 3 3 3 </td> | NOT AVAILABLE 3 3 3 |
| 1624 | Does this facility <i>routinely</i> observe any of the following postpartum or newborns related practices? | | | YES | NO | DON'T KNOW |
|------|--|-----------------------|-------------------------|-----------------------|------------------|-----------------|
| 04 | Routine, complete (head-to-toe) examination of newborn before discharge | | | 1 | 2 | 8 |
| 07 | Weigh the newborn immediately | | | 1 | 2 | 8 |
| 08 | Administer Vitamin K to newborn | | | 1 | 2 | 8 |
| 10 | Give full bath (immerse newborn in water) sho minutes/hours) after birth | rtly (i.e., wil | thin a few | 1 | 2 | 8 |
| 11 | Give the newborn prelacteal liquids | | | 1 | 2 | 8 |
| 14* | Provide maternal Vitamin A | | | 1 | 2 | 8 |
| 1625 | Please tell me if any of the following | (A) OBSERV | ED AVAILABLE | (B) NOT OBSERVED | | |
| | medicines or items are available at this service site today. I would like to see them. | AT LEAST ONE VALID | AVAILABLE NONE VALID | REPORTED AVAILABLE | NOT AVAILABLE | NO, OR NEVER |
| | CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | | | NOTSEEN | TODAY/DK | AVAILABLE |
| 02 | INJECTABLE ANTIBIOTIC (E.G., CEFTRIAXONE) | 1 | 2 | 3 | 4 | 5 |
| 03 | INJECTABLE UTEROTONIC (E.G., OXYTOCIN) | 1 | 2 | 3 | 4 | 5 |
| 04 | MAGNESIUM SULPHATE | 1 | 2 | 3 | 4 | 5 |
| 05 | INJECTABLE DIAZEPAM | 1 | 2 | 3 | 4 | 5 |
| 06 | IV SOLUTION (RINGER LACTATE) WITH INFUSION SET | 1 | 2 | 3 | 4 | 5 |
| 07 | SKIN DISINFECTANT (OTHER THAN CHLORHEXIDINE) | 1 | 2 | 3 | 4 | 5 |
| 08* | 7.1% CHLORHEXIDINE SOLUTION (UMBILICAL CORD CLEANSING) | 1 | 2 | 3 | 4 | 5 |

STANDARD PRECAUTIONS

| 1650 | ASSESS THE ROOM OR AREA FOR THE ITEMS LISTED BELOW. FOR ITEMS THAT YOU DO NOT SEE, ASK YOUR RESPONDENT TO SHOW THEM TO YOU. IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECORDED | ROOM OR AREA FOR THE ITEMS LISTED BELOW. 'HAT YOU DO NOT SEE, ASK YOUR RESPONDENT IEM TO YOU. CHILD VACCINATI CHILD CURATIVE (FAMILY PLANNIN ANTENATAL CAR TUBERCULOSIS [C NCD [Q2351] NOT PREVIOUSLY | | 11 12 13 14 14 15 19 22 22 31 | 'NEXT SECTION / SERVICE SITE |
|------|--|---|-----------------|--|------------------------------------|
| 1651 | STANDARD PRECAUTIONS AND CONDITIONS FOR CLIENT EXAMINATION | | OBSERVED | REPORTED NOT SEEN | NO |
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POUR PITC | CHER) | 1 | 2 | 3 |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOAP) | | 1 | 2 | 3 |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 |
| 04 | 04 WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLASTIC BIN LINER | | 1 | 2 | 3 |
| | | | 06 🗸 | | |
| 05 | OTHER WASTE RECEPTACLE | | 1 | 2 | 3 |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITANE, AL | .COHOL] | 1 | 2 | 3 |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH NEEDLES OR AUTO- DISABLE SYRINGES WITH NEEDLES | | 1 | 2 | 3 |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 |
| 11 | GOWNS | | 1 | 2 | 3 |
| 12 | 2 EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 |
| 1652 | DESCRIBE THE SETTING OF THE DELIVERY SERVICE PRIVATE ROOM ROOM OR AREA. OTHER ROOM WI | | ITH AUDITORY AN | D VISUAL PRIVA | L CY |

| | VISUAL PRIVACY ONLY | |
|--|---------------------|--|
| THANK YOUR RESPONDENT AND MOVE TO YOUR NEXT DATA COLLECTION POINT IF DIFFERENT FROM CURRENT LOCATION. | | |

SECTION 19: TUBERCULOSIS

| 1900 | CHECK Q102.10 TB SERVICE OFFERED IN FACILITY | NO TB SERVICE OFFERED IN FACILITY | |
|------|---|---|--|
| | ASK TO BE SHOWN THE LOCATION IN THE FACIL FIND THE PERSON MOST KNOWLEDGEABLE ABOUT I INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE S | TY WHERE TB SERVICES ARE PROVIDED. PROVISION OF TB SERVICES IN THE FACILITY. URVEY AND ASK THE FOLLOWING QUESTIONS. | |
| 1901 | How many days in a month are tuberculosis services offered at this facility? USE A 4-WEEK MONTH TO CALCULATE # OF DAYS | # OF DAYS /MONTH | |

TB DIAGNOSIS

| 1902 | Do providers in this facility make diagnosis that a client has tuberculosis? | YES1 NO2- | 1 004 |
|--------------------|---|--|---------------|
| 1903 | What is the most common method used by providers in this facility for diagnosing TB? PROBE TO DETERMINE METHOD USED. MULTIPLE | SPUTUM SMEAR MICROSCOPY A X-RAYB CLINICAL SYMPTOM C GeneXpertD | 1904 |
| 1903a [‡] | Is there a record/register of clients who are tested for TB diagnosis? | YES1 REGISTER NOT KEPT2_ | ▶1904 |
| 1903b [‡] | May I see the records or register of clients having tested for TB? CHECK THE RECORDS TO SEE TB TEST RESULTS ARE | REGISTER SEEN (PAPER)1 REGISTER SEEN (ELECTRONIC)2 REGISTER REPORTED, NOT SEEN3 | |
| 1904 | RECORDED Do providers in this facility ever refer clients outside this facility for TB diagnosis? | YES1 NO2- | 1908 |
| 1905 | Does this facility have an agreement with a referral site for TB test results to be returned to the facility either directly or through the client? | YES1 NO2 | |
| 1906 | Is there a record/register of clients who are referred for TB diagnosis? | YES1 REGISTER NOT KEPT2_ | → 1908 |
| 1907 | May I see the records or register of clients referred for TB testing? | REGISTER SEEN (PAPER) 1 REGISTER SEEN (ELECTRONIC) 2 REGISTER REPORTED, NOT SEEN 3 | |
| | CHECK THE RECORDS TO SEE TB DIAGNOSIS RESULTS ARE RECORDED | | |

TB TREATMENT

| 1908 | Do providers at this facility prescribe treatment for TB or manage patients who are on TB treatment | YES1 NO2— | 1910 |
|--------------------|--|--|------|
| 1908a [‡] | Do providers in this facility follow DOTS strategy for treatment of TB patients? | YES | 1910 |
| 1909 | What treatment regimen or approach is followed by providers in this facility for <u>newly diagnosed TB</u> ? i.e., for new patients, not for retreatment? | 2M INTENSIVE PHASE, 4M CONTINUATION PHASE | |
| | PROBE TO ARRIVE AT CORRECT RESPONSE | FOR F/UP4 PROVIDE FULL TREATMENT, WITH NO | |

| | ROUTINE DIRECT OBSERVATION | |
|--|------------------------------|--|
| | PHASE5 | |
| | DIAGNOSE, PRESCRIBE/PROVIDE | |
| | MEDICINES ONLY, NO F/UP6 | |
| | DIAGNOSE ONLY, NO TREATMENT | |
| | OR PRESCRIPTION OF MEDICINE7 | |

| 1910 | CHECK Q1902 AND Q1908 | | |
|---------------------|--|------------------------------|---------|
| | | NO TB DIAGNOSIS OR TREATMENT | |
| | IN FACILITY | NEXT SECTION OR SERVICE SITE | |
| | • | | |
| 1911 | Does this facility have a system for testing TB | YES1 | |
| | patients for HIV infection? | NO SYSTEM2- | 1913 |
| 1912 | May I see the system, or evidence of such a system? | SYSTEM OR REGISTER OBSERVED1 | |
| 1312 | | SYSTEM OR REGISTER REPORTED, | |
| | THE SYSTEM MAY BE IN THE FORM OF A REGISTER | NOT SEEN2 | |
| 1913 | Is HIV rapid diagnostic testing available from this | YES1 | |
| | service site? | NO2— | 1914a |
| 1914 | May I see a sample HIV rapid diagnostic test (RDT) | OBSERVED, AT LEAST 1 VALID1 | |
| | kit? | OBSERVED, NONEVALID2 | |
| | | REPORTED AVAILABLE, NOT SEEN | |
| | CHECK TO SEE IF AT LEAST ONE IS VALID | NOT AVAILABLE TODAY4 | |
| 1914a [‡] | Is follow up sputum microscopy conducted in the | YES1 | |
| 19110 | facility? | NO2— | 1015 |
| 404 41 [‡] | May I see the microscope used for follow up enuture | OBSERVED | 1915 |
| 1914b [.] | test? | REPORTED NOT SEEN2 | |
| 1915 | Do you have the national | YES1 | |
| 1313 | guideline/protocol/manual for the diagnosis and | NO2— | |
| | treatment of TB available in this service area? | | 1917 |
| 1916 | May I see the national guideline/protocol/manual? | OBSERVED1 | |
| | | REPORTED NOT SEEN2 | |
| 1917 | Do you have any guideline/protocol/manual for the | YES1 | |
| | management of HIV and TB co-infection available in this service area? | NO2— | 1919 |
| | | | |
| | THIS MAY BE PART OF OTHER GUIDELINE | | |
| 1918 | May I see the guideline/protocol/manual for the | OBSERVED1 | |
| | management of | REPORTED NOT SEEN2 | |
| | HIV and TB co-infection? | | |
| 1919 | Do you have any guideline/protocol/manual related | YES1 | |
| | to MDR-TB treatment available in this service area? | NO2— | 1920a |
| | | | |
| 1020 | This MAY BE PART OF OTHER GOIDELINE | OBSERVED 1 | |
| 1920 | treatment of MDR-TB? | REPORTED NOT SEEN | |
| 1920a [‡] | Is there a record/register of clients who receive TB | YES1 | |
| 1920a | treatment? | REGISTER NOT KEPT2 | 10200 |
| + | | REGISTER SEEN (DADER) 1 | · 1920C |
| 1920b ⁺ | May I see the records or register of clients who receive TB treatment? | REGISTER SEEN (LECTRONIC) | |
| | | REGISTER REPORTED, NOT SEEN | |
| | CHECK THE RECORDS TO SEE TB DIAGNOSIS RESULTS ARE | | |
| | RECORDED | | 1 |

| 1920c [‡] | Does this facility report TB-10? | YES1 NO2— | 1920e |
|----------------------------|---|--|--------------|
| 1920d [‡] | May I see the last report that was submitted? | OBSERVED1 REPORTED NOT SEEN2 | 15200 |
| 1920e [‡] | Does this facility report TB-11? | YES | 1920g |
| 1920f [‡] | May I see the last report that was submitted? | OBSERVED1 REPORTED NOT SEEN2 | |
| 1920g [‡] | Does this facility report TB-12? | YES1 NO2— | → 1921 |
| 1920h [‡] | May I see the last report that was submitted? | OBSERVED1 REPORTED NOT SEEN2 | |
| 1921 | CHECK Q1903 RESPONSE A OR D CIRCLED | RESPONSE A, OR D NOT CIRCLED | × 10F0 |
| 1922 | | | 1 920 |
| | Do you maintain any sputum containers at this service site for collecting sputum specimen? | YES1 NO2— | |
| 1923 | Do you maintain any sputum containers at this service site for collecting sputum specimen? May I see a sputum container? | YES | 1923a |
| 1923 1923a [‡] | Do you maintain any sputum containers at this service site for collecting sputum specimen? May I see a sputum container? Where the medicines for the treatment of TB generally stored in this facility? | YES 1 NO 2 OBSERVED 1 REPORTED NOT SEEN 2 NOT AVAILABLE TODAY 4 IN FREEZER 1 IN AIR CONDITIONED ROOM 2 NEITHER IN FREEZER NOR IN AIR CONDITIONED 3 | 1923a |

STANDARD PRECAUTIONS

| 1950 | ASSESS THE TB ROOM OR AREA FOR THE ITEMS LISTED BELOW.FOR ITEMS THAT YOU DO NOT SEE, ASK YOUR RESPONDENT TO SHOW THEM TO YOU. IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECORDED. | GENERAL INFORM CHILD VACCINATI CHILD CURATIVE FAMILY PLANNIN ANTENATAL CAR DELIVERY SERVIC NCD [Q2351] NOT PREVIOUSLY | MATION [Q710] ON [Q1051] CARE [Q1251] NG [Q1351] E [Q1451] CES [Q1651] SEEN | 11 12 13 14 14 19 17 17 22 31 | L 3 4 5 7 2 5 5 5 5 5 5 5 5 5 5 5 5 5 |
|------|---|--|---|--|---|
| 1951 | STANDARD PRECAUTIONS AND CONDITIONS EXAMINATION | FOR CLIENT | OBSERVED | REPORTED NOT SEEN | NO/ NOT AVAILABLR |
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POUR PITC | HER) | 1 | 2 | 3 |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOAP) | | 1 | 2 | 3 |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLASTIC BIN LINER | | 1 | 2 | 3 |
| 05 | OTHER WASTE RECEPTACLE | | 1 | 2 | 3 |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITANE, ALCOHOL] | | 1 | 2 | 3 |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH NEEDLES OR AUTO- DISABLE SYRINGES WITH NEEDLES | | 1 | 2 | 3 |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 |
| 11 | GOWNS | | 1 | 2 | 3 |

| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 |
|------|---|---|---------------------|------------|-------------|
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 |
| 1952 | DESCRIBE THE SETTING OF THE ROOM OR AREA | PRIVATE ROOM OTHER ROOM WI VISUAL PRIVACY VISUAL PRIVACY C NO PRIVACY | TH AUDITORY AN | D | 2 3 4 |
| 1953 | CHECK Q214 TB MEDS STORED IN OTHER LOCATION OR NOT STOCKED (RESPONSE 1 NOT CIRCLED) | TB MEDICINES STOF (RESPON | RED IN TB SERVICE A | AREA | 931 |
| THA | NK YOUR RESPONDENT AND MOVE TO YOUR NEXT D | DATA COLLECTION | N POINT IF DIFF | ERENT FROM | /I CURRENT |

SECTION 23: NON-COMMUNICABLE DISEASES

| 2300 | CHECK Q102.14 | | | | | |
|--------|--|---|--|--|--|--|
| | CHRONIC DISEASE SERVICES | | | | | |
| | AVAILABLE FROM FACILITY | | | | | |
| | ↓ | NEXT SECTION OR SERVICE SITE | | | | |
| ASK TO | ASK TO BE SHOWN THE LOCATION IN THE FACILITY WHERE CLIENTS WITH NON-COMMUNICABLE OR CHRONIC CONDITIONS SUCH AS | | | | | |
| DIA | BETES AND CARDIOVASCULAR DISEASES ARE SEEN. FIND THE PE | RSON MOST KNOWLEDGEABLE ABOUT PROVISION OF SUCH | | | | |
| SERV | ICES IN THE FACILITY. INTRODUCE YOURSELF, EXPLAIN THE PUR | POSE OF THE SURVEY AND ASK THE FOLLOWING QUESTIONS. | | | | |
| | D | IABETES | | | | |
| 2301 | Do providers in this facility diagnose and/or manage | YES, DIAGNOSE ONLY1 | | | | |
| | diabetes. | YES, TREAT ONLY2 | | | | |
| | | YES, DIAGNOSE AND TREAT3 | | | | |
| | | NO4 | | | | |
| | | | | | | |
| | | 2310 | | | | |
| 2302 | Do you have the national guidelines for the | YES1 | | | | |
| | diagnosis and management of diabetes available in | NO2-++2304 | | | | |
| | this service area? | 2304 | | | | |
| 2303 | May I see the national guidelines? | OBSERVED1 2310 | | | | |
| | | REPORTED, NOT SEEN2 | | | | |
| 2304 | Do you have any other guidelines for the diagnosis | YES1 | | | | |
| | and management of diabetes available in this service | NO2-> 2310 | | | | |
| | area? | | | | | |
| 2305 | May I see the other guidelines? | OBSERVED1 | | | | |

CARDIO-VASCULAR DISEASES/HYPERTENSION

REPORTED, NOT SEEN2

| 2310 | Do providers in this facility diagnose and/or manage cardio-vascular diseases including coronary heart disease in patients? | YES, DIAGNOSE ONLY | ► 2315 |
|-------|---|-----------------------------------|--------|
| 2311* | Do you have the national guideline/protocol/manual for the diagnosis and management of cardio-vascular disease available in this service area? | YES1 NO2- | 2313 |
| 2312* | May I see the national guideline/protocol/manual for the diagnosis and management of cardio-vascular disease? | OBSERVED1— REPORTED, NOT SEEN2 | ▶2315 |
| 2313* | Do you have any other guideline/protocol/manual for the diagnosis and management of cardio-vascular disease available in this service area? | YES1 NO2- | 2315 |
| 2314* | May I see the other guideline/protocol/ manual? | OBSERVED1 REPORTED, NOT SEEN2 | |
| 2315* | Do providers in this facility diagnose and/or manage hypertension patients? | YES, DIAGNOSE ONLY | |
| | | - | 2320 |
| 2316* | Do you have the national guideline/protocol/manual for the diagnosis and management of hypertension available in this service area? | YES1 NO2- | 2318 |
| 2317* | May I see the national guideline/protocol/manual for the diagnosis and management of hypertension? | OBSERVED1- REPORTED, NOT SEEN2 | ▶2320 |

2305

May I see the other guidelines?

| 2318* | Do you have any other guideline/protocol/manual for the diagnosis and management of hypertension available in this service area? | YES | 2320 |
|-------|---|----------------------------------|------|
| 2319* | May I see the other guideline/protocol/manual? | OBSERVED1 REPORTED, NOT SEEN2 | |

RESPIRATORY

| 2320* | Do providers in this facility diagnose and/or manage asthma/ chronic obstructive pulmonary diseases (COPD) in patients? | YES, DIAGNOSE ONLY | |
|-------|--|-----------------------------------|---------------|
| 2321* | Do you have the national guideline/protocol/manual for the diagno and management of asthma/COPD available in this service area? | YES1 NO2— | 2323 |
| 2322* | May I see the national guideline/protocol/manual for the diagnosis and management of asthma/COPD? | OBSERVED1— REPORTED, NOT SEEN2 | → 2325 |
| 2323* | Do you have any other guideline/protocol/manual for the diagnosis and/ management of asthma/COPD available in this service area? | YES1 NO2— | → 2325 |
| 2324* | May I see the other guideline/protocol/manual? | OBSERVED1 REPORTED, NOT SEEN2 | |

CERVICAL CANCER

| 2325* | Do providers in this facility screen patients for cervical cancer (either through visual inspection with Acetic Acid/Vinegar (VIA), pap smear or Human Papillomavirus (HPV) test)? | YES1 NO2— | 2330 |
|-------|---|-----------------------------------|-----------|
| 2326* | Do you have the national guideline/protocol/manual for cervical cancer screening available in this service area? | YES1 NO2— | → 2328 |
| 2327* | May I see the national guideline/protocol/manual for cervical cancer screening? | OBSERVED1— REPORTED, NOT SEEN2 | 2330 |
| 2328* | Do you have any other guideline/protocol/manual for cervical cancer screening available in this service area? | YES1 NO | 2330 |
| 2329* | May I see the other guideline/protocol/manual? | OBSERVED1 REPORTED, NOT SEEN2 | |

BASIC SUPPLIES AND EQUIPMEMENT

| 2330* | ASSESS THE ROOM OR AREA FOR THE BASIC SUPPLIES AND EQUIPMENT LISTED BELOW. IF THE SAME ROOM OR AREA HAS ALREADY BEEN ASSESSED, INDICATE WHERE THE DATA ARE RECORDED | | GENERAL INFORMATION SECTION (Q700)1- NOT PREVIOUSLY SEEN2 | | | | 1 | ► ₂₃₅₀ |
|-------|--|---------------|--|-----------------------|------------------|-----|-------|-------------------|
| 2331* | I would like to know if the following items are available today | (a) | AVAIL | ABLE? | | (b) | FUNCT | IONING |
| 2551 | in the main service area and are functioning ASK TO SEE ITEMS. | OBSERVE | D R | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01 | ADULT WEIGHING SCALE | 1 → b | | 2 > b | 3 02 | 1 | 2 | 8 |
| 02 | CHILD WEIGHING SCALE [250 GRAM GRADATION] | 1 - ▶b | | 2 — ▶b | 3 03 4 | 1 | 2 | 8 |
| 03 | INFANT WEIGHING SCALE [100 GRAM GRADATION] | 1 - ▶b | | 2 — ▶b | 3 04 | 1 | 2 | 8 |
| 04 | STADIOMETER [OR HEIGHT ROD] FOR MEASURING HEIGHT | 1 → b | | 2 > b | 3 05 4 | 1 | 2 | 8 |
| 05 | MEASURING TAPE [FOR CIRCUMFERENCE] | 1 | | 2 | 3 | | | |

| 06 | THERMOMETER | 1 → b | 2 → b | 3 07 4 | 1 | 2 | 8 |
|-----------------|---|---------------|-----------------|---------------------|---|---|---|
| 07 | STETHOSCOPE | 1 - ▶b | 2 > b | 3 08 4 | 1 | 2 | 8 |
| 08 | DIGITAL BP APPARATUS | 1 - ▶b | 2 — ▶b | 3 09 4 | 1 | 2 | 8 |
| 09 | MANUAL BP APPARATUS | 1 - ▶b | 2 → b | 3 10 | 1 | 2 | 8 |
| 10 | LIGHT SOURCE (FLASHLIGHT ACCPTABLE) | 1 - ▶b | 2 → b | 3 11 4 | 1 | 2 | 8 |
| 11 | SELF-INFLATING BAG AND MASK [ADULT] | 1 → b | 2 → b | 3 12 | 1 | 2 | 8 |
| 12 | SELF-INFLATING BAG AND MASK [PEDIATRIC] | 1 → b | 2 → b | 3 13 4 | 1 | 2 | 8 |
| 13 | MICRONEBULIZER | 1 - ▶b | 2 > b | 3 14 | 1 | 2 | 8 |
| 14 | SPACERS FOR INHALERS | 1 | 2 | 3 | | | |
| 15 | PEAK FLOW METERS | 1 - ▶b | 2 _ ▶b | 3 16 ◀ | 1 | 2 | 8 |
| 16 | PULSE OXIMETER | 1 - ▶b | 2 > b | 3 17 | 1 | 2 | 8 |
| 17 | OXYGEN CONCENTRATORS | 1 - ▶b | 2 > b | 3 18 4 | 1 | 2 | 8 |
| 18 | FILLED OXYGEN CYLINDER | 1 - ▶b | 2 — ▶b | 3 19 4 | 1 | 2 | 8 |
| 19 | OXYGEN DISTRIBUTION SYSTEM | 1 - ▶b | 2 → b | 3 20 4 | 1 | 2 | 8 |
| 20 | INTRAVENOUS INFUSION KITS - ADULT | 1 | 2 | 3 | | | |
| 21 | INTRAVENOUS INFUSION KITS - PEDIATRIC | 1 | 2 | 3 | | | |
| 22 [‡] | Examination table | 1 | 2 | 3 | | | |
| 23 [‡] | Glucometer with strip | 1 → b | 2 - | 3 <u></u> 2350 ◀ | 1 | 2 | 8 |

STANDARD PRECAUTIONS

| 2350 | ASSESS THE ROOM OR AREA FOR THE ITEMS LISTED GENERAL INFORMATION BELOW. FOR ITEMS THAT YOU DO NOT SEE. ASK YOUR CHILD VACCINATION | | | 11 | | | |
|------|---|----------------|-------------------------|-----------|----------------|--|--|
| | RESPONDENT TO SHOW THEM TO YOU. | CHILD CURATIVE | CARE [Q1251] | 13 | 3 | | |
| | | FAMILY PLANNIN | NG [Q1351] | | 1 - | | |
| | IF THE SAME ROOM OR AREA HAS ALREADY BEEN | ANTENATAL CAR | E [Q1451] ES [01651] | V 'NEXT | | | |
| | | TUBERCULOSIS [| Q1951] | | • SERVICE SITE | | |
| | | NOT PREVIOUSLY | SEEN | 31 | L L | | |
| 2351 | STANDARD PRECAUTIONS AND CONDITIONS | FOR CLIENT | OBSERVED | REPORTE | NO/ NOT | | |
| | EXAMINATION | | | D NOT | AVAILABLE | | |
| | | | | JLLN | | | |
| 01 | RUNNING WATER (PIPED, BUCKET WITH TAP OR POUR PITC | CHER) | 1 | 2 | 3 | | |
| 02 | ANY HAND-WASHING SOAP (MAY BE LIQUID SOAP) | | 1 | 2 | 3 | | |
| 03 | ALCOHOL-BASED HAND RUB | | 1 | 2 | 3 | | |
| 04 | WASTE RECEPTACLE (PEDAL BIN) WITH LID AND PLASTIC BI | N LINER | 1 | 2 | 3 | | |
| | | 06 🗲 | | | | | |
| 05 | OTHER WASTE RECEPTACLE | | 1 | 2 | 3 | | |
| 06 | SHARPS CONTAINER ("SAFETY BOX") | | 1 | 2 | 3 | | |
| 07 | DISPOSABLE LATEX/OTHER GLOVES | | 1 | 2 | 3 | | |
| 08 | DISINFECTANT/ANTISEPTICS [E.G., CHLORINE, HIBITANE, AL | _COHOL] | 1 | 2 | 3 | | |
| 09 | SINGLE-USE STANDARD DISPOSABLE SYRINGES WITH NEE DISABLE SYRINGES WITH NEEDLES | DLES OR AUTO- | 1 | 2 | 3 | | |
| 10 | MEDICAL MASKS | | 1 | 2 | 3 | | |
| 11 | GOWNS | | 1 | 2 | 3 | | |
| 12 | EYE PROTECTION [GOGGLES OR FACE PROTECTION] | | 1 | 2 | 3 | | |
| 13 | GUIDELINES FOR STANDARD PRECAUTIONS | | 1 | 2 | 3 | | |
| 2352 | DESCRIBE THE SETTING OF THE ROOM OR AREA | PRIVATE ROOM | | 1 | L | | |
| | | VISUAL PRIVACY | ITH AUDITORY AND | | , | | |
| | | VISUAL PRIVACY | ONLY | | 3 | | |
| | | NO PRIVACY | | | t I | | |
| THA | NK YOUR RESPONDENT AND MOVE TO YOUR NEXT D | | N POINT IF DIFFE | RENT FROM | /I CURRENT | | |
| | LOCATION. | | | | | | |

| | SECTION | 24: COVID-19 |
|-------------------|--|---|
| 2400 [‡] | CHECK Q102. 27 [‡] COVID-19 SERVICES DONE IN FACILITY | COVID-19 SERVICES NOT DONE NOT DONE IN FACILITY A NEXT SECTION OR SERVICE SITE |
| ASK T | O BE SHOWN THE LOCATION IN THE FACILITY WHERE COVID-19 SE ABOUT PROVISION OF SUCH SER INTRODUCE YOURSELF, EXPLAIN THE PURPOSE OF THE S | RVICES ARE DONE. FIND THE PERSON MOST KNOWLEDGEABLE VICES IN THE FACILITY. URVEY AND ASK THE FOLLOWING QUESTIONS. |
| 2401* | Does this facility provide counselling regarding COVID-19? | YES1 NO2 2403 |
| 2402* | What are the counselling this facility provides regarding COVID-19? | Keeping Social DistanceA Maintaining Respiratory- Cough EtiquetteB Use of Soap-water or, Alcohol based Hand Rub for Hand WashingC Not Touching Eyes-Nose- Mouth by handsD Using Medical Masks Environmental Disinfection Waste Management |
| 2403‡ | Does the facility collect sample for COVID-19 cases? | YES1 NO2 |
| 2404‡ | Does the facility do test for COVID-19? | YES1 NO2 → 2406 |
| 2405* | What are the types of test this facility do for COVID-19? | RT PCR1 RAPID ANTIGEN TEST2 BOTH RT PCR & RAPID ANTIGEN TEST3 |
| 2406 [‡] | Does this facility manage COVID-19 patient? | YES1 NO2 2500 |
| 2407* | Does this facility have separate room/corner for COVID- 19 patient? | YES1 NO2 |
| 2408* | Does this service area have Oxygen Provision for COVID-19 patient? | YES |
| 2409* | What kind of Oxygen Provision does this service area have? | OXYGEN CYLINDER |

| 2/10‡ | I would like to know if the following items are | (a |) A\ | /AILABLE? | | (b) | FUNC | FIONING |
|-------------------|---|---------------------|------------|-------------------------|------------------|-----|----------|---------------|
| 2410 | available in this COVID-19 service area and are functioning. | OBSERV | /ED | REPORTED, NOT SEEN | NOT AVAILABLE | YES | NO | DON'T KNOW |
| 01 | NASAL CANNULA | 1 | | 2 | 3 | | | |
| 02 | OXYGEN MASK | 1 | | 2 | 3 | | | |
| 03 | PPE | 1 | | 2 | 3 | | | |
| 04 | PULSE OXIMETER | 1- | b | 2 > b | 3 2411◀ | 1 | 2 | 8 |
| 2411* | Do you have the clinical case management guideline/protocol/manual for COVID-19 in th service area? | is | YES NO | | | | 1 2 | 2413 |
| 2412 [‡] | May I see the <i>clinical case management guideline/protocol/manual</i> for COVID-19? | | OBS REP | SERVED PORTED, NOT S | EEN | | 1 2 | |
| 2413 [‡] | Do you have the IPC guideline/protocol/manu COVID-19 in this service area? | al for | YES NO | | | | 1 2 — | ► 2415 |
| 2414 [‡] | May I see the <i>IPC guideline/protocol/manual</i> COVID-19? | for | OBS REP | SERVED PORTED, NOT S | EEN | | 1 2 | |
| 2415* | Do you have the "How to Run COVID-19 Hosp i guideline/protocol/manual in this service area | i tal" a? | YES NO | | | | 1 2 - | ► 2500 |
| 2416‡ | May I see the "How to Run COVID-19 Hospital guideline/protocol/manual? | " | OBS REP | SERVED PORTED, NOT S | EEN | | 1 2 | |

| SECTION 25: CESAREAN DELIVER |
|------------------------------|
|------------------------------|

| 2500 | CHECK Q102.16 | | | | | | | |
|--------------------|---|-----------------|---|------------------------------|-------------------------|------------|---------|--------|
| | CESAREAN SECTION DONE |] | (| CESAREAN DELIV | ERY NOT DONE | , E | | |
| | IN FACILITY | | | NEXT SECTION | OR SERVICE SITE | · | ┥ | |
| ASK TO I | ASK TO BE SHOWN THE LOCATION IN THE FACILITY WHERE CESAREAN DELIVERIES ARE DONE. FIND THE PERSON MOST KNOWLEDGEABLE ABOUT PROVISION OF SUCH SERVICES IN THE FACILITY. | | | | | | | |
| | INTRODUCE YOURSELF, EXPLAIN THE PURPOSI | E OF THE | SURVI | EY AND ASK THE | FOLLOWING QU | JESTIONS. | | |
| 2501 | Does the facility have a health worker who car perform Cesarean delivery (section) present at facility or on call 24 hours a day (including wee and on public holidays)? | the kends | YES NO | 5 | | | 1 2 | 2504 |
| 2502 | Is there a duty schedule or call list for 24-hr sta assignment? | aff | YES1 24-HOUR DUTY SCHEDULE NOT MAINTAINED2 | | | | | |
| 2503 | May I see the duty schedule or call list for 24-H assignment? | IR staff | SCH SCH | HEDULE OBSER | VED TED, NOT SEEN | | 1 2 | |
| 2504 | Does this facility have an anesthetist present in facility or on call 24 hours a day (including wee and on public holidays)? | n the ekends | YES NO | 5) | | | 1 2 | 2507 |
| 2505 | Is there a duty schedule or call list? | | YES 24- | HOUR DUTY SCI | HEDULE NOT MA | INTAINED. | 1 2 | ▶ 2507 |
| 2506 | May I see the duty schedule or call list? | | SCH SCH | HEDULE OBSER HEDULE REPOR | VED TED, NOT SEEN | | 1 2 | |
| 2507 | Have Cesarean deliveries been performed in the facility during the past 3 months? | nis | is YES1 NO2 | | | | | |
| 2507a [‡] | Please tell me if you have any of the following materials. | (| a) A' | VAILABLE | | | | |
| | IF YES, ASK TO SEE | OBSEF | RVED | REPORTED, NOT SEEN | NOT AVAILABLE | | | |
| 01 | EmONC Register | 1 | | 2 | 3 02 ◀ | | | |
| 02 | Do you have Operation Register? | 1 | | 2 | 3 2507b [‡] | | | |
| 2507b [‡] | Do you maintain documentation of the patier who are referred outside the facility? | nt | YES NO. | | | | 1 2 | 2510 |
| 2507c [‡] | Do you write diagnosis in the documentation the patient who are referred outside the facil | of ity? | YES NO. | | | | 1 2 | |
| 2507d [‡] | May I see the documentation of the patient w are referred outside the facility? | /ho | OBS REP | ERVED ORTED NOT S | EEN | | 1 2 | |
| | ASK TO SEE THE ROOM OR AREA WHERE CESARE | AN DELI | /ERIES | | DASK TO SEE TH | E ITEMS BE | LOW | |
| 2510 | available at this site today and is functioning. I would like to see them | OPCE | | (A) AVAILA | BLE? | VEC | (B) FUN | |
| | | OBSER | VED | NOT SEEN | AVAILABLE | YES | NO | KNOW |
| 01 | ANESTHESIA MACHINE | 1- | ►b | 2 -> b | 3 02 ◀ | 1 | 2 | 8 |
| 02 | TUBINGS AND CONNECTORS (TO CONNECT ENDOTRACHEAL TUBE) | 1- | ►b | 2 -> b | 3 03 4 | 1 | 2 | 8 |
| 03 | OROPHARYNGEAL AIRWAY (ADULT) | 1- | ►b | 2 -> b | 3 04 4 | 1 | 2 | 8 |

| 04 | OROPHARYNGEAL AIRWAY (PEDIATRIC) | 1 → b | 2 -> b | 3 05 • | 1 | 2 | 8 |
|-----------------|--|------------------|------------------|------------------|---|---|---|
| 05 | MAGILLS FORCEPS - ADULT | 1- > b | 2 -> b | 3 06 4 | 1 | 2 | 8 |
| 06 | MAGILLS FORCEPS - PEDIATRIC | 1 - ▶b | 2 -> b | 3 07 4 | 1 | 2 | 8 |
| 07 | ENDOTRACHEAL TUBE CUFFED SIZES 3.0 - 5.0 | 1 - ▶b | 2 -> b | 3 08 4 | 1 | 2 | 8 |
| 08 | ENDOTRACHEAL TUBE CUFFED SIZES 5.5 - 9.0 | 1 - ▶b | 2 -> b | 3 09 4 | 1 | 2 | 8 |
| 09 | INTUBATING STYLET | 1 - ▶b | 2 -> b | 3 10 4 | 1 | 2 | 8 |
| 10 | SPINAL NEEDLE | 1 → b | 2 -> b | 3 11 4 | 1 | 2 | 8 |
| 11* | OT TABLE | 1- > b | 2 -> b | 3 12 | 1 | 2 | 8 |
| 12* | OT LIGHT | 1 - ▶b | 2 -> b | 3 13 4 | 1 | 2 | 8 |
| 13* | IV STAND | 1 → b | 2 -> b | 3 14 4 | 1 | 2 | 8 |
| 14* | EMERGENCY POWER SUPPLY | 1 - ▶b | 2 → b | 3 15 • | 1 | 2 | 8 |
| 15* | INSTRUMENT SET FOR CESAREAN DELIVERY | 1 | 2 | 3 | | | |
| 16* | AIR CONDITIONER | 1 - ▶b | 2 -> b | 3 17 ↓ | 1 | 2 | 8 |
| 17* | OXYGEN CYLINDER WITH FLOWMETER | 1 - ▶b | 2 -> b | 3 18 4 | 1 | 2 | 8 |
| 18* | OXYGEN CYLINDER WITHOUT FLOWMETER | 1 - ▶b | 2 -> b | 3 – 18a ◀ | 1 | 2 | 8 |
| 18a | OXYGEN CONCENTRATORS | 1 - ▶b | 2 -> b | 3 18b ◀ | 1 | 2 | 8 |
| 18b | OXYGEN DISTRIBUTION SYSTEM | 1 - ▶b | 2 -> b | 3 19 4 | 1 | 2 | 8 |
| 19* | STERILE GLOVES | 1 | 2 | 3 | | | |
| 20* | DISINFECTANT | 1 | 2 | 3 | | | |
| 21 [‡] | Electro-surgical unit | 1 - ▶b | 2 -> b | 3 22 ↓ | 1 | 2 | 8 |
| 22 [‡] | Laparotomy set | 1 - ▶b | 2 -> b | 3 23 4 | 1 | 2 | 8 |
| 23 [‡] | Laryngoscopes | 1 - ▶b | 2 -> b | 3 24 | 1 | 2 | 8 |
| 24 [‡] | Pulse Oximeter | 1 - ▶b | 2 -> b | 3 25 • | 1 | 2 | 8 |
| 25 [‡] | Adult Resuscitation Trolley (Ambu with adult mask) | 1 → b | 2 -> b | 3 26 ◀ | 1 | 2 | 8 |
| 26 [‡] | Coagulating Cautery | 1 - ▶b | 2 -> b | 3 27 4 | 1 | 2 | 8 |
| 27 [‡] | Suction machine manual/electric | 1 - ▶b | 2 -> b | 3 28 4 | 1 | 2 | 8 |

| 28 [‡] | Trolleys | 1 - ▶b | 2 > b | 3 29 4 | 1 | 2 | 8 |
|---|--|---------------|------------------|------------------|---|---|---|
| 29 [‡] | Suture, dressing set (forceps, scissors) | 1 → b | 2 -> b | 3 2600 ◀ | 1 | 2 | 8 |
| THANK YOUR RESPONDENT AND MOVE TO YOUR NEXT DATA COLLECTION POINT IF DIFFERENT FROM CURRENT LOCATION. | | | | | | | |

SECTION 26: BLOOD TYPING AND COMPATIBILITY TESTING

| 2600 | CHECK Q102.18 | | | | | |
|------|--|------------|---|---|-----------------|-----------|
| | BLOOD TYPING SERVICES AVAILABLE FROM FACILITY |] | BLOOD TYPING AVAILABLE F NEXT SECTION | G SERVICES NOT FROM FACILITY OR SERVICE SITE | | |
| 2601 | Please tell me if any of the following reagents | (A) OBSERV | ED AVAILABLE | (| B) NOT OBSERVED | |
| | or equipment is available at this services site | AT LEAST | AVAILABLE | REPORTED | ΝΟΤ | NO, OR |
| | today. I would like to see them. | ONE VALID | NONE VALID | NOT SEEN | TODAY/DK | AVAILABLE |
| | CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | | | | | |
| 01 | Anti-A Reagent | 1 | 2 | 3 | 4 | 5 |
| 02 | Anti-B Reagent | 1 | 2 | 3 | 4 | 5 |
| 03 | Anti-D Reagent | 1 | 2 | 3 | 4 | 5 |
| 04 | COOMB'S REAGENT | 1 | 2 | 3 | 4 | 5 |
| 05 | Anti-A, B Reagent | 1 | 2 | 3 | 4 | 5 |

SECTION 27: BLOOD TRANSFUSION SERVICES

| 2700 | CHECK Q102.19 | | | | |
|--|---|---------------------------------|---------------------------------------|-------------------|-------------|
| | BLOOD TRANSFUSION AVAILABLE | BLOOD | D TRANSFUSION NO | г 🗌 | |
| | FROM FACILITY | | | | |
| | | | | | |
| ASK TO B | E SHOWN THE LOCATION IN THE FACILITY WHERE BLOOD IS COLLECTED PERSON MOST KNOWLEDGEABLE ABOUT PROVIS | ION OF BLOOD TRA | NSFUSION SERVICES | UK TU TKANSFUSION | I. FIND THE |
| | IN THE FACILITY INTRODUCE YOURSELF, EXPLAIN THE PURPOSE O | OF THE SURVEY AN | ID ASK THE FOLLOWIN | IG QUESTIONS. | |
| 2701 | What is the source of the blood that is transfused in | MEDICAL COLL | EGE HOSPITAL BLO | OD | |
| | this facility? | | | A | |
| | | BLOOD BANK | PITAL BLOOD BANK . IN THE FACILITY | В | |
| | PROBE FOR A COMPLETE LIST OF SOURCES OF | | NATING DIRECTLY | D | |
| | BLOOD. | RED CRESCENT | г | E | |
| | | OTHER | | x | |
| 2702 | | VES | (SPECIFY) | 1 | |
| 2702 | has blood transfusion been done in this facility in an obstetric context (i.e., for maternal care) during the | NO | | | |
| | past 3 months? | _ | | | |
| 2702a [‡] | Is there any register/record maintained for | YES | | 1 | |
| | pregnant women who required blood transfusion? | NO | | 2 _ | ▶2710 |
| 2702b [‡] | May I see that register/record? | OBSERVED | | 1 | |
| | | REPORTED, NO | OT SEEN | 2 | |
| | SCREENING FOR INFEC | CTIOUS DIS | EASES | | |
| 2710 | Is blood that is transfused in this facility screened, | YES | | 1 | |
| either in this facility or externally, for any | | NO2 | | | |
| 0=44 | infectious diseases prior to transfusion? | | | 1 | |
| 2/11 | Is the blood that is transfused screened only in the facility only at an external facility or both? | ONLY AT AN E | XTERNAL FACILITY | 2 | |
| | raemty, only at an external raemty, or both. | BOTH INTERNALLY AND EXTERNALLY3 | | | |
| 2712 | Is the blood that is transfused in the facility screened, | ALWAYS | SOMETIMES | RARELY | NO |
| | either in this facility or externally, for any of the | | | | |
| | following infectious diseases? | | | | |
| | IF YES, ASK: Is the blood "always", "sometimes", or | | | | |
| | "rarely" screened? | | | | |
| 01 | HIV | 1 | 2 | 3 | 4 |
| 02 | SYPHILIS | 1 | 2 | 3 | 4 |
| 03 | HEPATITIS B | 1 | 2 | 3 | 4 |
| 04 | HEPATITIS C | 1 | 2 | 3 | 4 |
| 05 | MALARIA | 1 | 2 | 3 | 4 |
| 2713 | Do you ever send blood sample outside the facility | YES | | 1 | |
| 2,15 | for screening for any of the tests mentioned above? | NO | | 2— | >7720 |
| 2714 | For which of the following tests do you could blood | (A) SEND S | PECIMEN OUT | (B) RECORD OF O | |
| 2714 | sample outside the facility for screening? | YES | NO | YES | NO |
| | , | | | | _ |
| | ASK TO SEE DOCUMENTATION | 4 - 1- | | 1 | 2 |
| 01 | HIV | 1−►p | 2 | 1 | 2 |
| | | 1 b h | 02 | 1 | |
| 02 | SYPHILIS | 1 − ∎0 | | 1 | 2 |
| 02 | | 1_ > h | | 1 | 2 |
| 03 | HEPATTIS B | 1 0 | 2 <u>-</u> | - | 2 |

| | | | 04 | | |
|----|-------------|---------------|-----------|---|---|
| 04 | HEPATITIS C | 1 b | 2 05 | 1 | 2 |
| 05 | MALARIA | 1 - ▶b | 2 2720 | 1 | 2 |

BLOOD STORAGE

| 2720 | 20 Has the facility run out of blood for more than one day | | YE | S | | 1 | |
|-------|--|-----------|-----------------------------|----------------|------------|----------------|-----------|
| | anytime during the past 3 months? | | N | D | | 2 | |
| 2721 | Is there a blood bank fridge or other refrigerator | | YE | S | | 1 | |
| | available for blood storage in this service area? |) | N | D | | 2- | 2724 |
| 2722 | May I see the blood bank fridge or other refrige | erator? | OE | SERVED | | 1 | |
| | | | RE | PORTED, NOT SI | EEN | 2— | 2724 |
| 2723 | WHAT IS THE TEMPERATURE IN THE BLOOD BANK FRI | IDGE OR | BE | TWEEN +2 AND | +6 DEGREES | 1 | |
| | OTHER REFRIGERATOR? | | AB | OVE +6 DEGREE | S | 2 | |
| | | | BE | LOW +2 DEGREE | ES | 3 | |
| | | | THERMOMETER NOT FUNCTIONAL4 | | | | |
| 2724 | Do you have any guideline/protocol/manual on the | | YES1 | | | | |
| | appropriate use of blood and safe transfusion | | NO2- | | | | |
| | practices? | ractices? | | | | | 2725A |
| 2725 | May I see the guideline/protocol/manu | ial on | OBSERVED1 | | | | |
| _ | appropriate use of blood and safe blood transfu | usion? | REPORTED, NOT SEEN2 | | | | |
| 2725A | Please tell me if any of the following items is | (A) OBS | SERV | ED AVAILABLE | (В |) NOT OBSERVED |) |
| | available at this services site today. I would | | т | AVAILABLE | REPORTED | NOT | NO, OR |
| | like to see them. | | | NONE VALID | AVAILABLE | AVAILABLE | NEVER |
| | CHECK TO SEE IF AT LEAST ONE IS VALID (NOT EXPIRED) | VALID | | | NOT SEEN | TODAY/DK | AVAILABLE |
| 01 | Disposable blood transfusion set | 1 | | 2 | 3 | 4 | 5 |
| 02 | Blood bags | 1 | | 2 | 3 | 4 | 5 |

| 3000 | ASSESS GENERAL CLEANLINESS / CONDITIONS OF FACILITY | YES | NO | |
|--|---|-------|---------|--|
| 01 | FLOOR: SWEPT, NO OBVIOUS DIRT OR WASTE | 1 | 2 | |
| 02 | COUNTERS/TABLES/CHAIRS: WIPED CLEAN- | 1 | 2 | |
| | NO OBVIOUS DUST OR WASTE | | | |
| 03 | NEEDLES, SHARPS OUTSIDE SHARPS BOX | 1 | 2 | |
| 04 | SHARPS BOX OVERFLOWING OR TORN/PIERCED | 1 | 2 | |
| 05 | BANDAGES/INFECTIOUS WASTE LYING UNCOVERED | 1 | 2 | |
| 06 | WALLS: SIGNIFICANT DAMAGE | 1 | 2 | |
| 07 | DOORS: SIGNIFICANT DAMAGE | 1 | 2 | |
| 08 | CEILING: WATER STAINS OR DAMAGE | 1 | 2 | |
| 09 | WASTE SEGREGATED ACCORDING TO COLOR BIN | 1 | 2 | |
| | AT THE SOURCE OF WASTE GENERATION? | | | |
| | INTERVIEW END TIME | Hours | Minutes | |
| THANK YOUR RESPONDENT AND MOVE TO YOUR NEXT DATA COLLECTION POINT IF DIFFERENT FROM CURRENT LOCATION | | | | |

SECTION 30: GENERAL FACILITY LEVEL CLEANLINESS

INTERVIEWER'S OBSERVATIONS

TO BE FILLED IN AFTER COMPLETING INTERVIEW

COMMENTS ABOUT RESPONDENT:

COMMENTS ON SPECIFIC QUESTIONS:

ANY OTHER COMMENTS:

SUPERVISOR'S OBSERVATIONS

| NAME OF THE SUPERVISOR: | DATE: |
|-------------------------|-------|
|-------------------------|-------|

2022 BANGLADESH HEALTH FACILITY SURVEY

HEALTH PROVIDER QUESTIONNAIRE

| HFALTH | PROVIDER | INTERVIEW |
|--------|----------|-----------|
| | | |

| FACILITY NUMBER | | | | |
|--|---------------------------------|--|--|--|
| PROVIDER SERIAL NUMBER | | | | |
| PROVIDER SEX (1= MALE, 2= Female) | | | | |
| PROVIDER Status (1=Assigned; 2=Seconded/deputed) | | | | |
| Interviewer Code | | | | |
| INDICATE IF PROVIDER WAS PREVIOUSLY INTERVIEWED | YES, Previously Interviewed1 | | | |
| IN ANOTHER FACILITY. | | | | |
| HE/SHE WAS INTERVIEWED | NAME AND NUMBER OF FACILITY | | | |
| | No, Not Previously Interviewed2 | | | |

TOTAL NUMBER OF PROVIDER INTERVIEWS

TOTAL NUMBER OF PROVIDERS INTERVIEWED.....

READ THE FOLLOWING CONSENT

Good day! My name is ______. We are here on behalf of the National Institute of Population Research and Training (NIPORT), Medical Education Family Welfare Division (ME&FWD) and the Ministry of Health and Family Welfare conducting a survey of health facilities to assists the Government in Knowing more about the health services in Bangladesh.

Now I will read a statement explaining the study.

Your facility was selected to participate in this study. We will be asking you several questions about the types of services that you personally provide, as well as questions about training you have received.

The information you provide us may be used by NIPORT, the MOHFW, or other organizations or researchers, for planning service improvement or further studies of services.

Neither your name nor that of any other health worker respondents participating in this study will be included in the data set or any report, however, there is a small chance that any of the respondents may be identified later. Still, we are asking for your help to ensure that the information we collect is accurate.

You may refuse to answer any question or choose to stop the interview at any time. However, we hope you will collaborate with the study. Do you have any questions about the study? Do I have your agreement to proceed?

| Interviewer' | s signature | Day | Month | Year | |
|--------------|---|-----------|-------|---------|--|
| SIGNATURE | OF INTERVIEWER INDICATES INFORMED CONSENT | PROVIDED | | | |
| 100 | May I begin the interview? | Yes No | | | |
| 101 | INTERVIEW START TIME | | | | |
| | | | Hours | Minutes | |

1. EDUCATION AND EXPERIENCE

| | - | |
|-------|--|---|
| 102. | I would like to ask you some questions | |
| | about your educational background | |
| | completed in total starting from your | MONTHS |
| | primary, secondary and further | |
| | education? | |
| 102a. | What professional degree/ | Bachelor of Medicine, Bachelor of Surgery |
| | basic training obtained? (please | (MBBS) A |
| | mention degree/course name) | Fellow of College of Physicians and Surgeons |
| | Multiple Answer can be obtained | (FCPS)B |
| | | |
| | | MD/MS C |
| | | Masters of Public Health D |
| | | B.Sc in Nursing (4years) E |
| | | Diploma in Nursing Science and Midwifery (3 years) F |
| | | B.Sc in Midwifery (post basic- |
| | | Zyears)G |
| | | Diploma course in Medical Laboratory Tecnology (3 |
| | | years) H |
| | | Medical Assistant Training CoursesI |
| | | Basic Training for FWV (18 months)J |
| | | Diploma in Medical Technology (BCG- EPI) K |
| | | CHCP basic training)L |
| | | Community Paramedic course |
| | | (2years) M |
| | | Others (Specific) |
| 103 | What is your current occupational | SPECIALIST/CONSULTANT MEDICINE [INCLUDING CARDIOLOGY]01 |
| 105. | category? | SPECIALIST/CONSULTANT |
| | | GENERAL SURGERY |
| | BOTH SENIOR AND JUNIOR | OBSTETRICS / GYNECOLOGY |
| | CONSULTANTS ARE INCLUDED IN | SPECIALIST/CONSULTANT PEDIATRICS04 |
| | CONSULTANTS CTEGORY | SPECIALIST/CONSULTANT PSYCHIATRY05 |
| | | SPECIALIST/CONSULTANT ANESTHESIA06 |
| 1 | | ANY OTHER SPECIALIST/CONSULTANT |

| | | NOT LISTED ABOVE 07 |
|------|--|---------------------------------------|
| | | MEDICAL OFFICER (MRBS) (ANV |
| | | |
| | | IMO MCH/ED |
| | | RMO, REGARDLESS OF DESIGNATION |
| | | |
| | | |
| | | DENTAL SUDCEON |
| | | |
| | | SACINO / INIEDICAL ASSISTANT |
| | | |
| | | |
| | | |
| | | ASSISTANT NURSE / STAFF NURSE |
| | | |
| | | |
| | | FAMILY WELFARE ASSISTANT (FWA) |
| | | |
| | | COMMUNITY HEALTH CARE PROVIDER (CHCP) |
| | | HEALTH INSPECTOR |
| | | ASSISTANT HEALTH INSPECTOR |
| | | NUTRITIONIST |
| | | HEALTH EDUCATOR |
| | | MEDICAL TECHNOLOGIST - LABORATORY25 |
| | | MEDICAL TECHNOLOGIST - EPI |
| | | MIDWIFE |
| | | PARAMEDICS |
| | | NURSE/MIDWIFE |
| | | TB LEPROSY CONTROL ASSISTANT (TLCA) |
| | | Other non-clinical/technical staff95 |
| | | OTHER CLINICAL STAFF NOT LISTED ABOVE |
| | | (Specify) |
| 104. | What year did you graduate (or | |
| | complete) with the qualification? | YEAR |
| | IF NO TECHNICAL QUALIFICATION ASK | |
| | What year did you complete any basic | |
| | training for your current occupational | |
| | category2 | |
| 4.05 | | |
| 105. | In what year did you start working in | |
| | this facility? | |
| | YOU MAY PROBE BY ASKING "HOW LONG | |
| | HAVE YOU WORKED IN THIS FACILITY"? | |
| 108 | Are you a manager or in charge for any | Yes 1 |
| 100. | Are you a manager or in-charge for any | No 2 |
| | clinical services? | NO Z |

| | 2. GENERAL TRAINING/NON-COMMUNICABLE DISEASES | | | | |
|---|---|--|--|---|--|
| 200 | I will like to ask you a few questions about in-service training you have received related to your work. In-service training refers to training you have received related to your work since you started working. I will start with some general topics. Note that the training topics I will mention may have been covered as standalone trainings, or they may have been covered under another training topic. | | | | |
| | Have you received any in-service training, training updates or refresher training in any of the following topics [READ TOPIC] IF YES, ASK: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES | |
| 01 | Standard precautions, including hand hygiene, cleaning and disinfection waste management, needle stick and sharp injury prevention? | 1 | 2 | 3 | |
| 02 | Any specific training related to injection safety practices or safe injection practices? | 1 | 2 | 3 | |
| 03 | Health Management Information Systems (HMIS) or reporting requirements for any service? | 1 | 2 | 3 | |
| 04 | Confidentiality and rights to non-discrimination practices for people living with HIV/AIDS | 1 | 2 | 3 | |
| 05* | Any specific training related to adolescent health services? | | 2 | 3 | |
| 06* | * Any specific training related to nutrition services? | | 2 | 3 | |
| 07 | Any specific training related to observation of acetic acid/pap smear/Human Papolloma Virus (HPV) test for screening of cervical cancer? | 1 | 2 | 3 | |
| 201 | CHECK Q103 FOR PROVIDER OCCUPATIONAL CATEGORY / QUALIFICATION CODE 25 (MEDICAL TECHNOLOGIST LABORATORY) CIRCLED CODE 25 (MEDICAL TECHNILOGIST LABORATORY) NOT CIRCLED | | | | |
| I will no training, t services ye | w ask you a few questions about services you personally provide in your current por raining updates or refresher trainings you may have received related to that service ou provide in your current position in this facility. The training topics I will mention training, or covered as part of another training topic. | sition in this e. Please rem may have be | facility and ar ember we are en covered as | ny in-service e talking about a stand-alone | |
| 202. | In your current position, and as a part of your work for this facility, do you personally provide any services that are designed to be youth or adolescent friendly? i.e., designed with the specific aim to encourage | | 1 2 | | |
| | youth or adolescent utilization? | | | | |
| 203. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics specific to youth or adolescent friendly services? | ONTHS GO GOR UPDATES | 1 2 3 | | |
| | IF YES: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | | | | |

DIABETES

| 207. | In your current position, and as a part of your work for this facility, do you personally diagnose and/or manage diabetes? | YES 1 NO | |
|------|--|----------------------------|--|
| 208. | Have you received any in-service training, training updates or refresher training on topics specific to the diagnosis and/or management of diabetes? | YES, WITHIN PAST 24 MONTHS | |
| | IF YES: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | | |

CARDIO-VASCULAR DISEASES

| 209. | In your current position, and as a part of your work for this facility, do you personally diagnose and/or manage cardio-vascular diseases such as hypertension? | YES 1 NO 2 | |
|------|---|--|--|
| 210. | Have you received any in-service training, trainin updates or refresher training on the diagnosis and/or management of cardio-vascular diseases? IF YES: Was the training, training update o refresher training within the past 24 months or more than 24 months ago? | YES, WITHIN PAST 24 MONTHS 1 Yes, OVER 24 MONTHS AGO 2 NO IN SERVICE TRAINING OR UPDATES 3 | |

CHRONIC RESPIRATORY DISEASES

| 211. | In your current position, and as a part of your work for this facility, do you personally diagnose and/or manage chronic respiratory conditions such as chronic obstructive pulmonary disease (COPD)? | YES | |
|------|---|----------------------------|--|
| 212. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on the diagnosis and/or management of chronic respiratory diseases? | YES, WITHIN PAST 24 MONTHS | |
| | IF YES: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | | |

3. CHILD HEALTH SERVICES

| 300. | In your current position, and as a part of your work for this facility, do you personally provide any child vaccination services? | YES | |
|------|--|-----|-------|
| 301. | In your current position, and as a part of your work for this facility, do you personally provide any child growth monitoring services? | YES | |
| 302. | In your current position, and as a part of your work for this facility, do you personally provide any child curative care services? | YES | |
| 303. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics related to child health or childhood illnesses? | YES | ▶ 400 |

| 304. | Have you received any <i>in-service training or training updates</i> In any of the following topics [READ TOPIC] | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES |
|------|---|-------------------------------------|-------------------------------|---|
| | IF YES: Was the training, training update or refresher training within the nast 24 months or more than 24 months ago? | | | |
| 01 | EPI OR COLD CHAIN MONITORING | 1 | 2 | 3 |
| 02 | INTEGRATED MANAGEMENT OF CHILDHOOD ILLNESSES | 1 | 2 | 3 |
| 06 | DIAGNOSIS AND/OR TREATMENT OF ACUTE RESPIRATORY INFECTIONS | 1 | 2 | 3 |
| 07 | DIAGNOSIS AND/OR TREATMENT OF DIARRHEA | 1 | 2 | 3 |
| 08 | MICRONUTRIENT DEFICIENCIES AND/OR NUTRITIONAL ASSESSMENT | 1 | 2 | 3 |
| 09 | BREASTFEEDING | 1 | 2 | 3 |
| 10 | COMPLIMENTARY FEEDING IN INFANTS | 1 | 2 | 3 |
| 13* | EARLY CHILDHOOD DEVELOPMENT | 1 | 2 | 3 |
| 14 | INFANT AND YOUNG CHILD FEEDING TRAINING (IYCF TRAINING) | 1 | 2 | 3 |
| 15 | SAM and MAM TRAINING | 1 | 2 | 3 |
| 16 | OTHER TRAINING ON CHILD HEALTH(SPECIFY) | 1 | 2 | 3 |

4. FAMILY PLANNING SERVICES

| 400. | In your current position, and as a part of your work for this facility, do you personally provide any family planning services? | YES1 NO2 | | | |
|------|---|---------------------|----------------------------------|-------------------------------|--|
| 401. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics related to family planning? | YES | | | |
| 403. | Have you received any <i>in-service training or training</i> any of the following topics [READ TOPIC] | g updates In | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES |
| | IF YES: Was the training, training update or refresher the past 24 months or more than 24 months ago? | r training within | | | |
| 01 | GENERAL COUNSELING FOR FAMILY PLANNING | | 1 | 2 | 3 |
| 02 | IUCD INSERTION AND/OR REMOVAL | | 1 | 2 | 3 |
| 03 | IMPLANT INSERTION AND/OR REMOVAL | | 1 | 2 | 3 |
| 04 | PERFORMING VASECTOMY | | 1 | 2 | 3 |
| 05 | PERFORMING TUBAL LIGATION | | 1 | 2 | 3 |
| 06 | CLINICAL MANAGEMENT OF FP METHODS, INCLUDING MANAGING SIDE EFFECTS | | 1 | 2 | 3 |
| 07 | FAMILY PLANNING FOR HIV POSITIVE WOMEN | | 1 | 2 | 3 |
| 08 | POST-PARTUM FAMILY PLANNING | | 1 | 2 | 3 |
| 09 | INJECTABLE CONTRACEPTIVES | | 1 | 2 | 3 |
| 10 | EMERGENCY CONTRACEPTIVE PILL | | 1 | 2 | 3 |
| 11 | Post Abortion/ MR | | 1 | 2 | 3 |
| 12 | OTHER TRAINING ON FAMILY PLANNING (SPECIFY) | | 1 | 2 | 3 |

5. MATERNAL HEALTH SERVICES

ANC - PNC

| 500. | In your current position, and as a part of your work for this facility, do you personally provide any antenatal care or postnatal care services? | YES, ANTENATAL YES, POSTNATAL YES, BOTH NO, NEITHER | | 1 2 3 4 | |
|-------|---|--|-------------------------------------|----------------------------------|---|
| | IF YES, PROBE AND INDICATE WHICH SERVICES ARE PROVIDED | | | | |
| 501. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics related to antenatal care or postnatal care? | YES NO | | 1 2- | ▶ 503 |
| 502. | Have you received any <i>in-service training or training</i> the following topics [READ TOPIC] [IF YES: Was the training, training update or refreshed past 24 months or more than 24 months ago? | g updates In any of r training within the | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES |
| 01 | ANC screening (e.g., blood pressure, sugar & albumin, and | emia)? | 1 | 2 | 3 |
| 02 | Counseling for ANC (e.g., nutrition, birth planning, FP and | newborn care)? | 1 | 2 | 3 |
| 03 | Complications of pregnancy and their management? | | 1 | 2 | 3 |
| 04 | Nutritional assessment of the pregnant woman, such as Body Mass Index calculation and Mid-Upper Arm circumference measurement? | | 1 | 2 | 3 |
| 06* | Postnatal care (PNC) (blood pressure checkup, examine perineum and abdomen, test urine for albumin) | | 1 | 2 | 3 |
| 07* | Postnatal care counselling (breast feeding, nutrition, FP, anemia) | | 1 | 2 | 3 |
| 08* | Management of pre-eclampsia/eclampsia | | 1 | 2 | 3 |
| 09* | Antenatal corticosteroids for threatened preterm labor | | 1 | 2 | 3 |
| 10* | Prevalence of postpartum hemorrhage (management of F | PH) | 1 | 2 | 3 |
| 503. | Do you <i>personally</i> provide any services that are specifically geared toward preventing mother-to- child transmission of HIV? IF YES, ASK: Which specific services do you provide? INDICATE WHICH OF THE LISTED SERVICES ARE PROVIDED AND PROBE: Anything else? | PREVENTIVE COUNSELING | | A B C D D E Y | |
| 504*. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics related to maternal and/or newborn health? | YES NO | | 1 2_ | ▶506 |
| 505. | Have you received any <i>in-service training or training updates</i> In any of the following topics [READ TOPIC] IF YES: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES |
| 02 | Newborn nutrition counseling of mother with HIV? | | 1 | 2 | 3 |
| 03 | Infant and young child feeding | | 1 | 2 | 3 |
| 04 | Modified obstetric practices as relates to HIV (e.g., not rupturing membranes)? | | 1 | 2 | 3 |

| 506. | In your current position, and as a part of your work for this facility, do you personally provide delivery services ? By that I mean conducting the actual delivery of newborns? | YES NO | | 509 — |
|------|--|---|----------------------------|---|
| 507. | During the past 6 months, approximately how many deliveries have you conducted as the <i>main provider (include deliveries conducted for private practice and for facility)?</i> | Total deliveries | | |
| 508. | When was the last time you used a partograph? | NEVER WITHIN PAST WE WITHIN PAST MO WITHIN PAST 6 N OVER 6 MONTHS | 0 EEK | |
| 509. | Have you received any <i>in-service training, training updates</i> <i>or refresher training</i> on topics related to delivery care? | YES NO | | 511 — |
| 510. | Have you received any <i>in-service training or training updates</i> In any of the following topics [READ TOPIC] IF YES: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES |
| 01 | Integrated Management of Pregnancy and Childbirth (IMPAC) | 1 | 2 | 3 |
| 02 | Comprehensive Emergency Obstetric Care (CEmOC) | 1 | 2 | 3 |
| 03 | Routine care for labor and normal vaginal delivery | 1 | 2 | 3 |
| 04 | Active Management of Third Stage of Labor (AMTSL) | 1 | 2 | 3 |
| 05 | Emergency obstetric care (EmOC)/Life-saving skills (LSS) - in general | 1 | 2 | 3 |
| 06 | Post abortion care | 1 | 2 | 3 |
| 07 | Special delivery care practices for preventing mother-to-child transmission of HIV? | 1 | 2 | 3 |

DELIVERY SERVICES

NEWBORN CARE SERVICES

| 511. | In your current position, and as a part of your work | YES1 | | | |
|------|--|----------|-------------------|--------------|----------|
| | for this facility, do you personally provide care for | NO | | 2 | |
| | the newborn? | | | | |
| 512. | Have you received any <i>in-service training, training</i> | YES | | 1 | |
| | updates or refresher training on topics related to | NO | | 2- | ▶600 |
| | newborn care? | | | | |
| 513. | Have you received any <i>in-service training or training updates</i> In any of | | Yes, | Yes, OVER | NO IN |
| | the following topics [READ TOPIC] | | WITHIN PAST 24 | 24 MONTHS | TRAINING |
| | | | MONTHS | AGO | OR |
| | months or more than 24 months ago? | | | | UPDATES |
| 01 | Neonatal resuscitation using bag and mask (Helping Babies Breath, HBB) | | 1 | 2 | 3 |
| 02 | Early and exclusive breastfeeding | | 1 | 2 | 3 |
| 03 | Newborn infection management (including injectable anti | biotics) | 1 | 2 | 3 |
| 04 | Thermal care (including immediate drying and skin-to-skir | n care) | 1 | 2 | 3 |
| 05 | Sterile cord cutting and appropriate cord care | | 1 | 2 | 3 |
| 06 | Kangaroo Mother Care (KMC) for low birth weight babies | | 1 | 2 | 3 |
| 08* | Essential Newborn Care | | 1 | 2 | 3 |
| 09* | Umblical cord care (use of 7.1% chlorhexidine) | | 1 | 2 | 3 |
| 10* | Emergency traige assessment training (ETAT) | | 1 | 2 | 3 |

| 11* | IMCI guidelines (0-59 days) | 1 | 2 | 3 |
|-----|-----------------------------------|---|---|---|
| 12* | Special care newborn unit (SCANU) | 1 | 2 | 3 |
| 13* | Comprehensive on newborn care | 1 | 2 | 3 |

6. SEXUALLY TRANSMITTED INFECTIONS -TB - COVID

SEXUALLY TRANSMITTED INFECTIONS

| 600. | In your current position, and as part of your work for this facility, do you personally provide any STI services? | YES NO | | 1 2 | |
|------|--|-----------|----------------------------------|----------------------------------|--|
| 601. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics related to | YES NO | | 1 2- | ▶603 |
| | STI services? | | | | |
| 602. | Have you received any <i>in-service training or training updates</i> In any of the following topics [READ TOPIC] IF YES: Was the training, training update or refresher training within the past 24 months or more than 24 months ago? | | Yes, WITHIN PAST 24 MONTHS | Yes, OVER 24 MONTHS AGO | NO IN SERVICE TRAINING OR UPDATES |
| 01 | Diagnosing and treating sexually transmitted infections (S | TIS) | 1 | 2 | 3 |
| 02 | The syndromic management for STIs | | 1 | 2 | 3 |
| 03 | Drug resistance to STI treatment medications | | 1 | 2 | 3 |

TUBERCULOSIS

| 603. | Now I will ask if you provide certain TB-related services. For each service, regardless of whether you currently | Do you pi [READ SE | rovide RVICE] | Have you received training or training update on [SERVICE]? IF YES, within 24 months or over? | | | | |
|------|--|-----------------------|------------------|---|-----------------|----------------|--|--|
| | provide it, I will also ask if you have received related in- service training, training undates or refresher training | YES | NO | YES, WITHIN 24 MONTHS | YES, OVER 24 | NO TRAINING | | |
| | READ THE QUESTIONS FROM COLUMNS A AND B | | | | MONTHS | | | |
| 01 | Diagnosis of tuberculosis based on sputum tests using AFB Smear Microscopy | 1 | 2 | 1 | 2 | 3 | | |
| 02 | Diagnosis of tuberculosis based on clinical symptoms or TB Diagnostic Algorithm | 1 | 2 | 1 | 2 | 3 | | |
| 03 | Treatment prescription for tuberculosis | 1 | 2 | 1 | 2 | 3 | | |
| 04 | Treatment follow-up services for tuberculosis | 1 | 2 | 1 | 2 | 3 | | |
| 05 | Direct Observation Treatment Short-course (DOTS) strategy | 1 | 2 | 1 | 2 | 3 | | |
| 06 | Management of TB - HIV co-infection | 1 | 2 | 1 | 2 | 3 | | |
| 07 | Management of MDR-TB or identification and referral of MDR-TB suspects | 1 | 2 | 1 | 2 | 3 | | |

COVID-19

| 604.‡ | Now I will ask if you have obtained these Covid-19 regarded in-service training, training updates or refresher training or not. | Have you receive training name) tr IF YES, within 24 | (state the g update? | |
|-------|---|--|-------------------------|-------------|
| | | YES, WITHIN 24 | YES, OVER | NO TRAINING |
| | | MONTHS | 24 MONTHS | |
| 1. | ADVANCED COVID-19 CLINICAL MANAGEMENT | 1 | 2 | 3 |
| | | | _ | - |
| 2. | IPC trainings on COVID-19 management | 1 | 2 | 3 |
| 3. | WASH (water, sanitation and hygiene) training | 1 | 2 | 3 |
| 4. | Comprehensive Training Course on COVID-19 for Doctors | 1 | 2 | 3 |
| 5. | COVID-19 Awareness and Prevention | 1 | 2 | 3 |

| 6. | COVID-19: Special Course for Informal Healthcare Providers | 1 | 2 | 3 |
|-----|--|---|---|---|
| 7. | Online Course on COVID-19 for Doctors | 1 | 2 | 3 |
| 8. | Training Course on COVID-19 for Healthcare Providers | 1 | 2 | 3 |
| 9. | Training Course on COVID-19 and Diabetes | 1 | 2 | 3 |
| 10. | Training on Clinical case management of Covid-19 | 1 | 2 | 3 |
| 11. | Training on 'How to Run COVID-19 Hospital' | 1 | 2 | 3 |

7. DIAGNOSTIC SERVICES

| | | - | | | | | | | | |
|-----------------|--|-----------|--|---|--|-------------------------------|--|--|--|--|
| 700. | In your current position, and as a part of your work for this facility, do you personally conduct laboratory tests? | YES NO | YES1 NO2 | | | | | | | |
| | CIRCLE 'NO' IF THE PROVIDER ONLY COLLECTS SPECIMENS. | | | | | | | | | |
| 701. | Please tell me if you personally conduct any of the following tests as part of your work in this facility | YES | NO | | | | | | | |
| 01 | Microscopic examining of sputum for diagnosing tuberculosis | | 1 | 2 | | | | | | |
| 02 | HIV rapid testing | | 1 | 2 | | | | | | |
| 04 | Hematology testing, such as anemia testing | | 1 | 2 | | | | | | |
| 06 | Malaria microscopy | | 1 | 2 | | | | | | |
| 07 | Malaria rapid diagnostic test (mRDT) | | 1 | 2 | | | | | | |
| 08 [‡] | RT-PCR test for COVID-19 | | 1 | 2 | | | | | | |
| 09 [‡] | Antigen test for COVID-19 | | 1 | 2 | | | | | | |
| 702. | Have you received any <i>in-service training, training</i> <i>updates or refresher training</i> on topics related to the different diagnostic tests you conduct? | | | | | | | | | |
| 703. | Have you received any <i>in-service training, training update</i> <i>refresher training</i> in any of the following topics? | es or | Yes, WITHIN Yes, OVER 24 NO II PAST 24 MONTHS AGO TRAI MONTHS UPD, | | | N SERVICE INING OR ATES | | | | |
| | [READ TOPIC] | | | | | | | | | |
| | IF YES: Was the training, training update or refresher train within the past 24 months or more than 24 months ago? | ning | | | | | | | | |
| 01 | Microscopic examination of sputum for diagnosing tuberculosis | | 1 2 | | | 3 | | | | |
| 02 | HIV testing | | 1 | 2 | | 3 | | | | |
| 04 | Blood screening for HIV prior to transfusion? | | 1 | 2 | | 3 | | | | |
| 05 | Blood screening for Hepatitis B prior to transfusion? | | 1 | 2 | | 3 | | | | |
| 06 | Tests for monitoring ART such as TLC and serum creatinine. | | 1 | 2 | | 3 | | | | |
| 07 | Malaria microscopy | | 1 | 2 | | 3 | | | | |
| 08 | Malaria rapid diagnostic test (mRDT) | | 1 | 2 | | 3 | | | | |
| 09 [‡] | Specimen collection for diagnosing COVID-19 test | | 1 | 2 | | 3 | | | | |
| 10* | RT-PCR Test for diagnosing COVID-19 test | | 1 2 | | | 3 | | | | |
| 111 | | | 1 | 2 | | 3 | | | | |

8. WORKING CONDITIONS IN FACILITY

| 800. | Now I want to ask you a few more questions about your work in this facility In an average week, how many hours do you work in this facility? | AVERAGE HOURS PER WEEK | |
|------|---|----------------------------|--|
| | IF WEEKS ARE NOT CONSISTENT, ASK THE RESPONDENT TO AVERAGE OUT HOW MANY HOURS PER MONTH AND THEN DIVIDE THIS BY 4 | | |
| 801. | Now I would like to ask you some questions about | YES, IN THE PAST 3 MONTHS1 | |

| | supervision you have personally received. This YES, IN THE PAST 4-6 MONTHS | | | | | | | | | | | | |
|------|--|--|--|---|--|---------------------------|-------|--|--|--|--|--|--|
| 802. | How many times in the past six month | How many times in the past six months has your NUMBER OF TIMES | | | | | | | | | | | |
| 803. | The last time you were personally supervised, did your supervisor do any of the following: [READ TOPIC] | | EVERY DAY | YES | NO | 96 DON'T KNOW | | | | | | | |
| C | D1 Check your records or reports? | CHECKED R | ECORD | 1 | 2 | 8 | 1 | | | | | | |
| C | 02 Observe your work? | OBSERVED | WORK | 1 | 2 | 8 | | | | | | | |
| C | Provide any feedback (either positive or negative) | FEEDBACK | | 1 | 2 05 ◀ | 8 − 05 ← | | | | | | | |
| C | Give you verbal or written feedback that you were doing your work well? | VERBAL PRA | AISE | 1 | 2 | 8 | | | | | | | |
| C | D5 Provide updates on administrative or technical issues related to your work? | PROVIDED | JPDATES | 1 | 2 | 8 | | | | | | | |
| C | Discuss problems you have encountered? | PROBLEMS | 1 | 2 | 8 | | | | | | | | |
| 804. | Do you have a written job description of your current job or position in this facility? YES, OBSERVED | | | | | | | | | | | | |
| | IF YES, ASK: May I see it? | | | | | | | | | | | | |
| 805. | Are there any opportunities for promote current job? | tion in your | NO UNCERTAIN/D | ON'T KNOW | | 1 2 8 | | | | | | | |
| 806. | Which type(s) of salary supplement do y if any? PROBE: Anything else? | you receive, | MONTHLY OR PERDIEM WHI DUTY ALLOW/ PAYMENT FOF (NOT ROUT OTHER (SI NONE | DAILY SALARY EN ATTENDING ANCE R EXTRA ACTIVI INELY PROVIDI PECIFY) | ARY SUPPLEMENTA ING TRAININGB C TIVITIES VIDED)D X | | | | | | | | |
| 807. | In your current position, what non- incentives have you received for the w if any? PROBE BY ASKING: Anything else? options? | n-monetary ork you do, Any other | TIME OFF / VACATIONS A UNIFORMS, BACKPACKS< CAPS, etc. | | | | | | | | | | |
| 808. | Among the various things related to working situation that you would like t improved; can you tell me the three tha think would most improve your abili provide good quality of care services? P rank them in order of importance, w being the most important. | your SU o see MO at you TF ity to MO Please BET vith 1 SU BET FL MO IE 1ST PF | RE SUPPORT FR JPERVISOR RE KNOWLEDG &AINING RE SUPPLIES/ST TER QUALITY EC JPPLIES S WORKLOAD (I TER WORKING EXIBLE TIMES RE INCENTIVES &OMOTION HOL | OM E / UPDATES OCK QUIPMENT/ .e. MORE STA HOURS / (SALARY, .IDAYS) | \FF) | A B C E Ran F | nking | | | | | | |

| MENTIONED INTO THE 1ST BOX, AND REPEAT | TRANSPORTATION FOR REFERRAL |
|--|--------------------------------------|
| WITH THE 2ND AND 3RD | PATIENTS H |
| | PROVIDING ARTI |
| | PROVIDING PEP J |
| IF THE PROVIDER ONLY MENTIONS 1 OR 2 TIEMS | INCREASED SECURITYK |
| PUT CODE "Y" IN THE REMAINING BOX OR BOXES | BETTER FACILITY |
| | INFRASTRUCTUREL |
| DO NOT READ CHOICES TO YOUR RESPONDENT | MORE AUTONOMY / |
| | INDEPENDENCE M |
| | STAFF (COUNSELING / SOCIAL |
| | ACTIVITIES)N |
| | OTHERX |
| | (SPECIFY) |
| | NONEY |
| THANK YOUR RESPONDENT AND MC | VE TO THE NEXT DATA COLLECTION POINT |

| STAFF LISTING FORM: HEALTH WORKERS AVAILABLE ON DAY OF VISIT | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|--------|-------|--|-----------------------------|----|--------|---------------|----------------------------|---------------------------|--------------------|------------|------|-----|----------------|-----------|----------|-----------------|--------------|---------|-------------------------|-----------------------|---------------------------|---|
| FACILIT | FACILITY NUMBER I TOTAL NUMBER OF PROVIDERS LISTED ON ALL 5 SHEETS INTERVIEWER CODE | | | | | | | | | | | | | | | | | | | | | | | |
| LIST ALL CLINICAL STAFF/PROVIDERS WHO ARE PRESENT TODAY IN THIS FACILITY. COMPILE THIS LIST AS THE TEAM MOVES FROM ONE SERVICE AREA (OR DEPARTMENT) TO ANOTHER OBTAINING INFORMATION ON THE SERVICES THAT THE FACILITY PROVIDES AND FOR WHICH INVENTORY SECTIONS ARE BEING COMPLETED, AND/OR FOR WHICH CLIENT-PROVIDER OBSERVATIONS ARE BEING DONE. WRITE THE HEALTH WORKER'S QUALIFICATION CODE IN COLUMN 3 "PROVIDER QUALIFICATION CODE" , AND THE PROVIDER'S GENDER UNDER COLUMN 4 "GENDER" . PUT CHECK MARKS IN THE APPROPRIATE HEADINGS UNDER COLUMN 5 "SERVICES PROVIDED IN FACILITY" TO INDICATE THE SERVICE THAT THE PROVIDER PROVIDES IN THE FACILITY. IN COLUMN 6 "INTERVIEWED FOR INVENTORY" CIRCLE THE LINE NUMBER IF THE PROVIDER WAS INTERVIEWED FOR ANY SECTION OF THE INVENTORY QUESTIONNAIRE. FINALLY, IN COLUMN 7 "SELECTED FOR HEALTH WORKER INTERVIEW" CIRCLE THE LINE NUMBER IF THE PROVIDER IS SELECTED TO BE INTERVIEWED WITH THE INDIVIDUAL HEALTH WORKER QUESTIONNAIRE. | | | | | | | | | | | | | | | | | | | | | | | | |
| (1) | | (2) | | | (3 | 3) | (4) | | | | | | | | | (5) | | | | | | | (6) | (7) |
| | NAME C | OF PRC | VIDER | | | | | | | 1 | | | | SER | VICES PRO | OVIDED IN | FACILITY | , | 1 | | 1 | 1 | | |
| | | | | | PROVIDER QUALIFICATION CODE | | Gender | PRESCRIBE ART | HIV COUNSELING AND TESTING | DIAGI HIV/AIDS RELATED | MALARIA MALARIA | REATM 8 | IENT | NCD | ANTENATAL CARE | PMTCT | DELIVERY | FAMILY PLANNING | CHILD HEALTH | SURGERY | CONDUCT LABORATORY TEST | OTHER CLIENT SERVICES | INTERVIEWED FOR INVENTORY | SELECTED FOR HEALTH WORKER INTERVIEW |
| 1 | | | | | | | | | | | | | | | | | | | | | | | 1 | 1 |
| 2 | | | | | | | | | | | | | | | | | | | | | | | 2 | 2 |
| 3 | | | | | | | | | | | | | | | | | | | | | | | 3 | 3 |
| 4 | | | | | | | | | | | | | | | | | | | | | | | 4 | 4 |
| 5 | | | | | | | | | | | | | | | | | | | | | | | 5 | 5 |
| 6 | | | | | | | | | | | | | | | | | | | | | | | 6 | 6 |
| 7 | | | | | | | | | | | | | | | | | | | | | | | 7 | 7 |
| 8 | | | | | | | | | | | | | | | | | | | | | | | 8 | 8 |
| 9 | | | | | | | | | | | | | | | | | | | | | | | 9 | 9 |
| 10 | | | | | | | | | | | | | | | | | | | | | | | 10 | 10 |

PROVIDER QUALIFICATION CATEGORY

| SPECIALIST/CONSULTANT MEDICINE (INCLUDING CARDIOLOGY) 01 |
|--|
| SPECIALIST /CONSULTANT GENERAL SURGERY02 |
| SPECIALIST/CONSULTANT OBSTETRICS / GYNOCOLOGY |
| SPECIALIST/CONSULTANT PEDIATRICS04 |
| SPECIALIST /CONSULTANT PSYCHIATRY 05 |
| SPECIALIST/CONSULTANT ANESTHESIA |
| ANY OTHER SPECIALIST/CONSULTANT NOT LISTED ABOVE07 |
| MEDICAL OFFICER (MBBS) (ANY NON-SPECIALIST DOCTOR, |
| INCLUDING ASSISTANT SURGEON, EMO, IMO, MCH/FP, RMO |
| REGARDLESS OF DESIGNATION OR TITLE |
| MEDICAL OFFICER - ANESTHETIST |

| DENTAL SURGON 10 |
|---|
| SACMO / MEDICAL ASSISTANT11 |
| MATRON12 |
| NURSING SUPERVISOR13 |
| SENIOR STAFF NURSE14 |
| ASSISTANT NURSE/STAFF NURSE (IN PRIVATE) 15 |
| FAMILY WELFARE VISITOR (FWV)16 |
| FAMILY WELFARE ASSISTANT (FWA)17 |
| HEALTH ASSISTANT18 |
| COMMUNITY HEALTH CARE PROVIDER 19 |
| HEALTH INSPECTOR 20 |

| ASSISTANT HEALTH INSPECTOR | 21 |
|-----------------------------------|----|
| NUTRITIONIST | 22 |
| HEALTH EDUCATOR | 23 |
| MEDICAL TECHNOLOGIST - LABORATORY | 25 |
| MEDICAL TECHNOLOGIST - EPI | 26 |
| MIDWIFE | 30 |
| PARAMEDICS (INPRIVATE/NON) | 31 |
| NURSE MIDWIFE | 32 |
| TB LEPROSY CONTROL ASSISTANT(TLC) | 33 |
| OTHER CLINICAL STAFF NOT | |
| LISTED ABOVE | 96 |

(SPECIFY)

| STAFF LISTING FORM: HEALTH WORKERS AVAILABLE ON DAY OF VISIT | | | | | | | | | | | | | |
|--|--|--|---|--|--|--|------------------|--|--|--|--|--|--|
| FACILITY NUMBER | | | Τ | | | TOTAL NUMBER OF PROVIDERS LISTED ON ALL 5 SHEETS | INTERVIEWER CODE | | | | | | |

LIST ALL CLINICAL STAFF/PROVIDERS WHO ARE PRESENT TODAY IN THIS FACILITY. COMPILE THIS LIST AS THE TEAM MOVES FROM ONE SERVICE AREA (OR DEPARTMENT) TO ANOTHER OBTAINING INFORMATION ON THE SERVICES THAT THE FACILITY PROVIDES AND FOR WHICH INVENTORY SECTIONS ARE BEING COMPLETED, AND/OR FOR WHICH CLIENT-PROVIDER OBSERVATIONS ARE BEING DONE. WRITE THE HEALTH WORKER'S QUALIFICATION CODE IN **COLUMN 3 "PROVIDER QUALIFICATION CODE"**, AND THE PROVIDER'S GENDER UNDER **COLUMN 4 "GENDER"**. PUT CHECK MARKS IN THE APPROPRIATE HEADINGS UNDER **COLUMN 5 "SERVICES PROVIDED IN FACILITY"** TO INDICATE THE SERVICE THAT THE PROVIDER PROVIDES IN THE FACILITY. **IN COLUMN 6 "INTERVIEWED FOR INVENTORY"** CIRCLE THE LINE NUMBER IF THE PROVIDER WAS INTERVIEWED FOR ANY SECTION OF THE INVENTORY QUESTIONNAIRE. FINALLY, IN **COLUMN 7 "SELECTED FOR HEALTH WORKER INTERVIEW"** CIRCLE THE LINE NUMBER IF THE PROVIDER IS SELECTED TO BE INTERVIEWED WITH THE INDIVIDUAL HEALTH WORKER QUESTIONNAIRE.

| (1) | (2) | (3 | 3) | (4) | (5) | | | | | | | | | | | | | | (6) | (7) | |
|-----|------------------|--|------------------|--------|---------------|---------------------------|---------------------|---------|----|-----|-----|----------------|-------|----------|-----------------|--------------|---------|-------------------------|-----------------------|-------------------------|--|
| | NAME OF PROVIDER | NAME OF PROVIDER SERVICES PROVIDED IN FACILITY | | | | | | | | | | | | 1 | | | | | | | |
| | | | L C C C | Gender | | Ŋ | DIAGNOSIS/TREATMENT | | | | | | | | | | L | | ۲ | ĸ | |
| | | PROVIDER QUALIFICATION CC | | | PRESCRIBE ART | HIV COUNSELING AND TESTIN | HIV/AIDS RELATED | MALARIA | TB | ILS | NCD | ANTENATAL CARE | PMTCT | DELIVERY | FAMILY PLANNING | СНІГР НЕАГТН | SURGERY | CONDUCT LABORATORY TESI | OTHER CLIENT SERVICES | INTERVIEWED FOR INVENTO | SELECTED FOR HEALTH WORKE INTERVIEW |
| 11 | | | | | | | | | | | | | | | | | | | | 11 | 11 |
| 12 | | | | | | | | | | | | | | | | | | | | 12 | 12 |
| 13 | | | | | | | | | | | | | | | | | | | | 13 | 13 |
| 14 | | | | | | | | | | | | | | | | | | | | 14 | 14 |
| 15 | | | | | | | | | | | | | | | | | | | | 15 | 15 |
| 16 | | | | | | | | | | | | | | | | | | | | 16 | 16 |
| 17 | | | | | | | | | | | | | | | | | | | | 17 | 17 |
| 18 | | | | | | | | | | | | | | | | | | | | 18 | 18 |
| 19 | | | | | | | | | | | | | | | | | | | | 19 | 19 |
| 20 | | | | | | | | | | | | | | | | | | | | 20 | 20 |
| SPECIALIST/CONSULTANT MEDICINE (INCLUDING CARDIOLOGY) 01 |
|--|
| SPECIALIST /CONSULTANT GENERAL SURGERY02 |
| SPECIALIST/CONSULTANT OBSTETRICS / GYNOCOLOGY |
| SPECIALIST/CONSULTANT PEDIATRICS04 |
| SPECIALIST /CONSULTANT PSYCHIATRY |
| SPECIALIST/CONSULTANT ANESTHESIA06 |
| ANY OTHER SPECIALIST/CONSULTANT NOT LISTED ABOVE07 |
| MEDICAL OFFICER (MBBS) (ANY NON-SPECIALIST DOCTOR, |
| INCLUDING ASSISTANT SURGEON, EMO, IMO, MCH/FP, RMO |
| REGARDLESS OF DESIGNATION OR TITLE |
| MEDICAL OFFICER - ANESTHETIST |

| DENTAL SURGON 10 |
|---|
| SACMO / MEDICAL ASSISTANT 11 |
| MATRON12 |
| NURSING SUPERVISOR13 |
| SENIOR STAFF NURSE 14 |
| ASSISTANT NURSE/STAFF NURSE (IN PRIVATE) 15 |
| FAMILY WELFARE VISITOR (FWV)16 |
| FAMILY WELFARE ASSISTANT (FWA) 17 |
| HEALTH ASSISTANT |
| COMMUNITY HEALTH CARE PROVIDER 19 |
| HEALTH INSPECTOR 20 |

| ASSISTANT HEALTH INSPECTOR | 21 |
|-----------------------------------|----|
| NUTRITIONIST | 22 |
| HEALTH EDUCATOR | 23 |
| MEDICAL TECHNOLOGIST - LABORATORY | 25 |
| MEDICAL TECHNOLOGIST - EPI | 26 |
| MIDWIFE | 30 |
| PARAMEDICS (INPRIVATE/NON) | 31 |
| NURSE MIDWIFE | 32 |
| TB LEPROSY CONTROL ASSISTANT(TLC) | 33 |
| OTHER CLINICAL STAFF NOT | |
| LISTED ABOVE | 96 |

| | | | STAFF LISTING FORM: HEALTH WORKERS AVAILABLE ON DA | AY OF | VIS | IT | | |
|-----------------|--|--|--|-------|-----|----|--|--|
| FACILITY NUMBER | | | TOTAL NUMBER OF PROVIDERS LISTED ON ALL 5 SHEETS INTERVIEWER | CODE | | | | |

LIST ALL CLINICAL STAFF/PROVIDERS WHO ARE PRESENT TODAY IN THIS FACILITY. COMPILE THIS LIST AS THE TEAM MOVES FROM ONE SERVICE AREA (OR DEPARTMENT) TO ANOTHER OBTAINING INFORMATION ON THE SERVICES THAT THE FACILITY PROVIDES AND FOR WHICH INVENTORY SECTIONS ARE BEING COMPLETED, AND/OR FOR WHICH CLIENT-PROVIDER OBSERVATIONS ARE BEING DONE. WRITE THE HEALTH WORKER'S QUALIFICATION CODE IN **COLUMN 3 "PROVIDER QUALIFICATION CODE"**, AND THE PROVIDER'S GENDER UNDER **COLUMN 4 "GENDER"**. PUT CHECK MARKS IN THE APPROPRIATE HEADINGS UNDER **COLUMN 5 "SERVICES PROVIDED IN FACILITY"** TO INDICATE THE SERVICE THAT THE PROVIDER PROVIDES IN THE FACILITY. **IN COLUMN 6 "INTERVIEWED FOR INVENTORY"** CIRCLE THE LINE NUMBER IF THE PROVIDER WAS INTERVIEWED FOR ANY SECTION OF THE INVENTORY QUESTIONNAIRE. FINALLY, IN **COLUMN 7 "SELECTED FOR HEALTH WORKER INTERVIEW"** CIRCLE THE LINE NUMBER IF THE PROVIDER IS SELECTED TO BE INTERVIEWED WITH THE INDIVIDUAL HEALTH WORKER QUESTIONNAIRE.

| (1) | (2) | (3 | 3) | (4) | | | | | | | | | (5) | | | | | | | (6) | (7) |
|-----|------------------|---------------------------|----|--------|---------------|---------------------------|------------------|--------------------|----|-----|-----|----------------|-------|----------|-----------------|--------------|---------|-------------------------|-----------------------|--------------------------|--|
| | NAME OF PROVIDER | | | | | | | | | | SER | VICES PRO | | FACILITY | , | | | r | 1 | | |
| | | DDE | | | | ŋ | DIAG | IAGNOSIS/TREATMENT | | | | | | 24 | ¥. | | | | | | |
| | | PROVIDER QUALIFICATION CO | | Gender | PRESCRIBE ART | HIV COUNSELING AND TESTIN | HIV/AIDS RELATED | MALARIA | ТВ | STI | NCD | ANTENATAL CARE | РМТСТ | DELIVERY | FAMILY PLANNING | СНІГР НЕАГТН | SURGERY | CONDUCT LABORATORY TEST | OTHER CLIENT SERVICES | INTERVIEWED FOR INVENTOF | SELECTED FOR HEALTH WORKE INTERVIEW |
| 21 | | | | | | | | | | | | | | | | | | | | 21 | 21 |
| 22 | | | | | | | | | | | | | | | | | | | | 22 | 22 |
| 23 | | | | | | | | | | | | | | | | | | | | 23 | 23 |
| 24 | | | | | | | | | | | | | | | | | | | | 24 | 24 |
| 25 | | | | | | | | | | | | | | | | | | | | 25 | 25 |
| 26 | | | | | | | | | | | | | | | | | | | | 26 | 26 |
| 27 | | | | | | | | | | | | | | | | | | | | 27 | 27 |
| 28 | | | | | | | | | | | | | | | | | | | | 28 | 28 |
| 29 | | | | | | | | | | | | | | | | | | | | 29 | 29 |
| 30 | | | | | | | | | | | | | | | | | | | | 30 | 30 |

| SPECIALIST/CONSULTANT MEDICINE (INCLUDING CARDIOLOGY) 01 | |
|--|--|
| SPECIALIST /CONSULTANT GENERAL SURGERY02 | |
| SPECIALIST/CONSULTANT OBSTETRICS / GYNOCOLOGY | |
| SPECIALIST/CONSULTANT PEDIATRICS04 | |
| SPECIALIST /CONSULTANT PSYCHIATRY 05 | |
| SPECIALIST/CONSULTANT ANESTHESIA06 | |
| ANY OTHER SPECIALIST/CONSULTANT NOT LISTED ABOVE07 | |
| MEDICAL OFFICER (MBBS) (ANY NON-SPECIALIST DOCTOR, | |
| INCLUDING ASSISTANT SURGEON, EMO, IMO, MCH/FP, RMO | |
| REGARDLESS OF DESIGNATION OR TITLE | |
| MEDICAL OFFICER - ANESTHETIST | |

| DENTAL SURGON 10 |
|---|
| SACMO / MEDICAL ASSISTANT 11 |
| MATRON12 |
| NURSING SUPERVISOR13 |
| SENIOR STAFF NURSE14 |
| ASSISTANT NURSE/STAFF NURSE (IN PRIVATE) 15 |
| FAMILY WELFARE VISITOR (FWV)16 |
| FAMILY WELFARE ASSISTANT (FWA)17 |
| HEALTH ASSISTANT |
| COMMUNITY HEALTH CARE PROVIDER 19 |
| HEALTH INSPECTOR 20 |

| ASSISTANT HEALTH INSPECTOR | 21 |
|-----------------------------------|----|
| NUTRITIONIST | 22 |
| HEALTH EDUCATOR | 23 |
| MEDICAL TECHNOLOGIST - LABORATORY | 25 |
| MEDICAL TECHNOLOGIST - EPI | 26 |
| MIDWIFE | 30 |
| PARAMEDICS (INPRIVATE/NON) | 31 |
| NURSE MIDWIFE | 32 |
| TB LEPROSY CONTROL ASSISTANT(TLC) | 33 |
| OTHER CLINICAL STAFF NOT | |
| LISTED ABOVE | 96 |

| | | | STAFF LISTING FORM: HEALTH WORKERS A | AVAILABLE ON DAY OF | VISI | T | | |
|-----------------|--|--|--|---------------------|------|---|--|--|
| FACILITY NUMBER | | | TOTAL NUMBER OF PROVIDERS LISTED ON ALL 5 SHEETS | INTERVIEWER CODE | | | | |

LIST ALL CLINICAL STAFF/PROVIDERS WHO ARE PRESENT TODAY IN THIS FACILITY. COMPILE THIS LIST AS THE TEAM MOVES FROM ONE SERVICE AREA (OR DEPARTMENT) TO ANOTHER OBTAINING INFORMATION ON THE SERVICES THAT THE FACILITY PROVIDES AND FOR WHICH INVENTORY SECTIONS ARE BEING COMPLETED, AND/OR FOR WHICH CLIENT-PROVIDER OBSERVATIONS ARE BEING DONE. WRITE THE HEALTH WORKER'S QUALIFICATION CODE IN **COLUMN 3 "PROVIDER QUALIFICATION CODE"**, AND THE PROVIDER'S GENDER UNDER **COLUMN 4 "GENDER"**. PUT CHECK MARKS IN THE APPROPRIATE HEADINGS UNDER **COLUMN 5 "SERVICES PROVIDED IN FACILITY"** TO INDICATE THE SERVICE THAT THE PROVIDER PROVIDES IN THE FACILITY. **IN COLUMN 6 "INTERVIEWED FOR INVENTORY"** CIRCLE THE LINE NUMBER IF THE PROVIDER WAS INTERVIEWED FOR ANY SECTION OF THE INVENTORY QUESTIONNAIRE. FINALLY, IN **COLUMN 7 "SELECTED FOR HEALTH WORKER INTERVIEW"** CIRCLE THE LINE NUMBER IF THE PROVIDER IS SELECTED TO BE INTERVIEWED WITH THE INDIVIDUAL HEALTH WORKER QUESTIONNAIRE.

| (1) | (2) | (3 | 3) | (4) | | | | | | | | | (5) | | | | | | | (6) | (7) |
|-----|------------------|---------------------------|----|--------|---------------|---------------------------|------------------|---------|------|---------------------|-----|----------------|-----------|------------|-----------------|--------------|---------|-------------------------|-----------------------|---------------------------|--|
| | NAME OF PROVIDER | | | | | | | | | | SER | VICES PRO | OVIDED IN | I FACILITY | , | | | | | | |
| | | DE | DE | | | | | ŋ | DIAG | DIAGNOSIS/TREATMENT | | | | | | | | | | | ¥. |
| | | PROVIDER QUALIFICATION CC | | Gender | PRESCRIBE ART | HIV COUNSELING AND TESTIN | HIV/AIDS RELATED | MALARIA | ЯД | STI | NCD | ANTENATAL CARE | РМТСТ | DELIVERY | FAMILY PLANNING | СНІГР НЕАГТН | SURGERY | CONDUCT LABORATORY TEST | OTHER CLIENT SERVICES | INTERVIEWED FOR INVENTORY | SELECTED FOR HEALTH WORKE INTERVIEW |
| 31 | | | | | | | | | | | | | | | | | | | | 31 | 31 |
| 32 | | | | | | | | | | | | | | | | | | | | 32 | 32 |
| 33 | | | | | | | | | | | | | | | | | | | | 33 | 33 |
| 34 | | | | | | | | | | | | | | | | | | | | 34 | 34 |
| 35 | | | | | | | | | | | | | | | | | | | | 35 | 35 |
| 36 | | | | | | | | | | | | | | | | | | | | 36 | 36 |
| 37 | | | | | | | | | | | | | | | | | | | | 37 | 37 |
| 38 | | | | | | | | | | | | | | | | | | | | 38 | 38 |
| 39 | | | | | | | | | | | | | | | | | | | | 39 | 39 |
| 40 | | | | | | | | | | | | | | | | | | | | 40 | 40 |

| SPECIALIST/CONSULTANT MEDICINE (INCLUDING CARDIOLOGY) 01 | |
|--|--|
| SPECIALIST /CONSULTANT GENERAL SURGERY02 | |
| SPECIALIST/CONSULTANT OBSTETRICS / GYNOCOLOGY | |
| SPECIALIST/CONSULTANT PEDIATRICS04 | |
| SPECIALIST /CONSULTANT PSYCHIATRY 05 | |
| SPECIALIST/CONSULTANT ANESTHESIA06 | |
| ANY OTHER SPECIALIST/CONSULTANT NOT LISTED ABOVE07 | |
| MEDICAL OFFICER (MBBS) (ANY NON-SPECIALIST DOCTOR, | |
| INCLUDING ASSISTANT SURGEON, EMO, IMO, MCH/FP, RMO | |
| REGARDLESS OF DESIGNATION OR TITLE | |
| MEDICAL OFFICER - ANESTHETIST | |

| DENTAL SURGON 10 |
|---|
| SACMO / MEDICAL ASSISTANT 11 |
| MATRON12 |
| NURSING SUPERVISOR13 |
| SENIOR STAFF NURSE14 |
| ASSISTANT NURSE/STAFF NURSE (IN PRIVATE) 15 |
| FAMILY WELFARE VISITOR (FWV)16 |
| FAMILY WELFARE ASSISTANT (FWA)17 |
| HEALTH ASSISTANT |
| COMMUNITY HEALTH CARE PROVIDER 19 |
| HEALTH INSPECTOR 20 |

| ASSISTANT HEALTH INSPECTOR | 21 |
|-----------------------------------|----|
| NUTRITIONIST | 22 |
| HEALTH EDUCATOR | 23 |
| MEDICAL TECHNOLOGIST - LABORATORY | 25 |
| MEDICAL TECHNOLOGIST - EPI | 26 |
| MIDWIFE | 30 |
| PARAMEDICS (INPRIVATE/NON) | 31 |
| NURSE MIDWIFE | 32 |
| TB LEPROSY CONTROL ASSISTANT(TLC) | 33 |
| OTHER CLINICAL STAFF NOT | |
| LISTED ABOVE | 96 |

| STAFF LISTING FORM: HEALTH WORKERS AVAILABLE ON DAY OF VISIT | | | | | | | | | | | | | | |
|--|-----------------|--|--|--|--|--|--|------------------|--|--|--|--|-----------|--|
| | FACILITY NUMBER | | | | | | TOTAL NUMBER OF PROVIDERS LISTED ON ALL 5 SHEETS | INTERVIEWER CODE | | | | | \square | |

LIST ALL CLINICAL STAFF/PROVIDERS WHO ARE PRESENT TODAY IN THIS FACILITY. COMPILE THIS LIST AS THE TEAM MOVES FROM ONE SERVICE AREA (OR DEPARTMENT) TO ANOTHER OBTAINING INFORMATION ON THE SERVICES THAT THE FACILITY PROVIDES AND FOR WHICH INVENTORY SECTIONS ARE BEING COMPLETED, AND/OR FOR WHICH CLIENT-PROVIDER OBSERVATIONS ARE BEING DONE. WRITE THE HEALTH WORKER'S QUALIFICATION CODE IN **COLUMN 3 "PROVIDER QUALIFICATION CODE"**, AND THE PROVIDER'S GENDER UNDER **COLUMN 4 "GENDER"**. PUT CHECK MARKS IN THE APPROPRIATE HEADINGS UNDER **COLUMN 5 "SERVICES PROVIDED IN FACILITY"** TO INDICATE THE SERVICE THAT THE PROVIDER PROVIDES IN THE FACILITY. **IN COLUMN 6 "INTERVIEWED FOR INVENTORY"** CIRCLE THE LINE NUMBER IF THE PROVIDER WAS INTERVIEWED FOR ANY SECTION OF THE INVENTORY QUESTIONNAIRE. FINALLY, IN **COLUMN 7 "SELECTED FOR HEALTH WORKER INTERVIEW"** CIRCLE THE LINE NUMBER IF THE PROVIDER IS SELECTED TO BE INTERVIEWED WITH THE INDIVIDUAL HEALTH WORKER QUESTIONNAIRE.

| (1) | (2) | (3 | 3) | (4) | | | | | | | | | (5) | | | | | | | (6) | (7) |
|-----|------------------|--------------------------|-----|--------|---------------|-------------------------------|------------------|---------|-------|------|-----|----------------|-------|----------|-----------------|--------------|---------|-------------------------|-----------------------|---------------------------|--|
| | NAME OF PROVIDER | | | | | SERVICES PROVIDED IN FACILITY | | | | | | | | | | | | | | | |
| | | ODE | ODE | | | ۶ | DIAG | NOSIS/1 | REATN | IENT | | | | | | | | L | | | ĸ |
| | | PROVIDER QUALIFICATION C | | Gender | PRESCRIBE ART | HIV COUNSELING AND TESTII | HIV/AIDS RELATED | MALARIA | Я | ILS | NCD | ANTENATAL CARE | РМТСТ | DELIVERY | FAMILY PLANNING | СНІГР НЕАГТН | SURGERY | CONDUCT LABORATORY TEST | OTHER CLIENT SERVICES | INTERVIEWED FOR INVENTORY | SELECTED FOR HEALTH WORKE INTERVIEW |
| 41 | | | | | | | | | | | | | | | | | | | | 41 | 41 |
| 42 | | | | | | | | | | | | | | | | | | | | 42 | 42 |
| 43 | | | | | | | | | | | | | | | | | | | | 43 | 43 |
| 44 | | | | | | | | | | | | | | | | | | | | 44 | 44 |
| 45 | | | | | | | | | | | | | | | | | | | | 45 | 45 |
| 46 | | | | | | | | | | | | | | | | | | | | 46 | 46 |
| 47 | | | | | | | | | | | | | | | | | | | | 47 | 47 |
| 48 | | | | | | | | | | | | | | | | | | | | 48 | 48 |
| 49 | | | | | | | | | | | | | | | | | | | | 49 | 49 |
| 50 | | | | | | | | | | | | | | | | | | | | 50 | 50 |

| SPECIALIST/CONSULTANT MEDICINE (INCLUDING CARDIOLOGY) 01 |
|--|
| SPECIALIST /CONSULTANT GENERAL SURGERY02 |
| SPECIALIST/CONSULTANT OBSTETRICS / GYNOCOLOGY |
| SPECIALIST/CONSULTANT PEDIATRICS04 |
| SPECIALIST /CONSULTANT PSYCHIATRY |
| SPECIALIST/CONSULTANT ANESTHESIA06 |
| ANY OTHER SPECIALIST/CONSULTANT NOT LISTED ABOVE07 |
| MEDICAL OFFICER (MBBS) (ANY NON-SPECIALIST DOCTOR, |
| INCLUDING ASSISTANT SURGEON, EMO, IMO, MCH/FP, RMO |
| REGARDLESS OF DESIGNATION OR TITLE |
| MEDICAL OFFICER - ANESTHETIST |

| DENTAL SURGON 10 |
|---|
| SACMO / MEDICAL ASSISTANT 11 |
| MATRON12 |
| NURSING SUPERVISOR13 |
| SENIOR STAFF NURSE14 |
| ASSISTANT NURSE/STAFF NURSE (IN PRIVATE) 15 |
| FAMILY WELFARE VISITOR (FWV)16 |
| FAMILY WELFARE ASSISTANT (FWA)17 |
| HEALTH ASSISTANT18 |
| COMMUNITY HEALTH CARE PROVIDER 19 |
| HEALTH INSPECTOR 20 |

| ASSISTANT HEALTH INSPECTOR | 21 |
|-----------------------------------|----|
| NUTRITIONIST | 22 |
| HEALTH EDUCATOR | 23 |
| MEDICAL TECHNOLOGIST - LABORATORY | 25 |
| MEDICAL TECHNOLOGIST - EPI | 26 |
| MIDWIFE | 30 |
| PARAMEDICS (INPRIVATE/NON) | 31 |
| NURSE MIDWIFE | 32 |
| TB LEPROSY CONTROL ASSISTANT(TLC) | 33 |
| OTHER CLINICAL STAFF NOT | |
| LISTED ABOVE | 96 |

